

Di-hadron SIDIS measurements at CLAS

Silvia Pisano^{1,a} for the CLAS Collaboration

¹INFN - Istituto Nazionale di Fisica Nucleare
Via Enrico Fermi, 40, 00044 Frascati, Rome, Italy.

Abstract. Semi-inclusive deep-inelastic scattering (SIDIS) is an essential tool to probe nucleon internal structure. Through single hadron SIDIS processes, indeed, it is possible to access the TMDs, containing information on both the longitudinal and transverse motion of the partons. In recent years, moreover, an increasing attention has been devoted to dihadron SIDIS. It constitutes the golden channel to access the higher-twist collinear Parton Distribution Functions $e(x)$ and $h_L(x)$, so far only marginally known, whose extraction will complete the collinear description of the nucleon at the twist-3 level. The CLAS detector in the Hall-B at JLab, thanks to its large acceptance, is particularly suited for such measurements. Analyses aiming at the extraction of dihadron SIDIS Beam and Target-Spin Asymmetries are presently in progress. In these proceedings, preliminary results for the Beam-Spin Asymmetry are reported, together with a summary of the dihadron SIDIS experimental program at JLab.

1 Introduction

Despite many years of investigation, both theoretical and experimental, the collinear picture of the proton is still far from being complete. At the twist-3 level, the nucleon is described through six Parton Distribution Functions (PDFs), that depend on the fraction x of the proton total momentum carried by the parton: the *leading twist* unpolarized $f_1(x)$, the helicity distribution $g_1(x)$ and the so-called transversity $h_1(x)$ and the three *higher twist* ones, $e(x)$, $h_L(x)$ and $g_T(x)$ [1, 2]. Among them, $f_1(x)$ and $g_1(x)$ are pretty well known, having been extracted through Inclusive Deep Inelastic Scattering experiments. $h_1(x)$, on the other hand, due to its *chiral-odd* nature, appears in observables only through the coupling with a second *chiral-odd* object, and can only be accessed through Semi-Inclusive DIS processes, where at least one hadron is detected in the final state together with the outgoing electron. In these processes, indeed, the transversity distribution appears coupled to the Fragmentation Function, another *chiral-odd* object describing the hadronization of the struck quark that leads to the final state hadrons. As to the three *higher – twist* PDFs, $e(x)$ and $h_L(x)$ are almost unknown, while $g_T(x)$ has been also extracted through inclusive DIS measurements. These *higher-twist* PDFs encodes important information on the physics of the largely unexplored quark-gluon correlations. Furthermore, the x -integral of $e(x)$ is related to the poorly known *scalar charge* of the nucleon, a fundamental quantity sensible to parity-violating components of the nucleon-nucleon interactions, while the x -integral of $h_1(x)$ is connected to the nucleon *tensor charge*, so far only poorly constrained by the available data.

^ae-mail: silvia.pisano@lnf.infn.it

The cleanest process to access the unknown PDFs is represented by the Di-hadron Semi-Inclusive Deep Inelastic Scattering (di-hadron SIDIS), where, together with the outgoing electron, at least two hadrons are detected in the final state. The cross-section of such a process, indeed, is described in terms of specific Structure Functions (SF) [3, 4], where the PDFs of interest appear coupled to the so-called Interference Fragmentation Functions, i.e. functions specifically introduced to describe the hadronization of the struck quark into two hadrons. With respect to the single-hadron case, furthermore, the PDFs appear coupled to the FF only through a simple product, and not through a convolution over the quark transverse momentum, making the actual extraction of the parton distribution from di-hadron SIDIS data fully model independent. The CEBAF *Large Acceptance Spectrometer* (CLAS) [5] installed in the Hall-B at the Thomas Jefferson Accelerator Facility, is particularly suited for the extraction of these quantities. Its large acceptance implies a good capability in the reconstruction of multi-particle final states, allowing the investigation of the aforementioned process in a wide range of kinematics.

2 CLAS data sets

The CLAS experiment performed two analysis devoted to the extraction of di-hadron SIDIS related observables. One has been performed on a data set collected in 2003 by impinging a 5.5 GeV longitudinally-polarized electron beam - provided by the CEBAF - on an unpolarized liquid-hydrogen target 2H , for a total integrated luminosity of 21 fb $^{-1}$. The second data set has been collected in 2009 with a slightly larger electron energy (5.967 GeV), and by exploiting a longitudinally-polarized 3NH target. Due to the polarization degrees of freedom active in the process, the first data set allows only the extraction of the longitudinal Beam-Spin Asymmetry A_{LU} , while the presence of a polarized target in the second data set provides access also to the Target-Spin Asymmetry A_{UL} and to the Double-Spin Asymmetry A_{LL} . In this proceedings, preliminary results for A_{LU} will be shown, both on the 2H and the 3NH target. Extraction of the Target and of the Double-Spin Asymmetries is in progress on the 3NH data.

3 Beam-Spin Asymmetry extraction

A_{LU} turns out to be proportional to the following structure function:

$$F_{LU}^{\sin\phi_R} = -x \frac{|R| \sin\theta}{Q} \left[\frac{M}{M_h} x e^q(x) H_1^{\leftarrow q}(z, \cos\theta, M_h) + \frac{1}{z} f_1^q(x) \tilde{G}^{\leftarrow q}(z, \cos\theta, M_h) \right] \quad (1)$$

where $e(x)$ appears coupled to the Interference Fragmentation Function $H_1^{\leftarrow q}$, recently extracted [6] through the *Belle* data [7]. The channel of interest in these analyses is the semi-inclusive electro-production of two hadrons, i.e. $ep \rightarrow e\pi^+\pi^-X$. The final di-hadron sample has been selected by identifying final states where an electron and at least one π^+ and one π^- are present. When more than one positive or negative pion is present, all the possible combinations are considered and treated as possible candidate events. The right kinematic region is then selected by applying: *i*) standard DIS cuts: $Q^2 > 1\text{GeV}^2$ (where Q^2 is the photon virtuality), to be in the deep-inelastic scattering regime, and $W > 2\text{GeV}$ (where W is the centre-of-mass energy of the system composed of the proton and the virtual-photon), to avoid the resonance region; *ii*) a cut on the missing mass of the system $m_X > 1.05\text{GeV}$, to eliminate contamination from exclusive channels; *iii*) the cut $x_F = 2 \frac{p_{\perp}}{W} > 0$, where p_{\perp} is the longitudinal momentum of the final hadron in the $p - \gamma^*$ reference frame, to select hadrons actually coming from the quark fragmentation (current-fragmentation region, CFR) and not from the




Figure 1. A_{LU} dependence on x_B (left plot), z (middle plot) and $m_{\pi^+\pi^-}$ (right plot), as extracted from the 2H data (yellow squares) and from the 3NH data (magenta triangles).

target. A_{LU} is then defined as

$$A_{LU} = \frac{1}{P_B} \frac{N^+ - N^-}{N^+ + N^-} \quad (2)$$

where N^\pm represents the number of countings in the positive/negative helicity state, and P_B is the beam polarization. Both the 2H and 3NH data sets are divided in three sets of one-dimensional bins, in order to explore the dependence of the $\sin\phi$ moment of A_{LU} ($A_{LU}^{\sin\phi}$) of x_B , the Bjorken variable, $m_{\pi^+\pi^-}$, the two-pion invariant mass, and z , the fraction of the virtual photon energy carried by the final hadron pairs. In Fig. 1 the results obtained on the two data sets are shown. $A_{LU}^{\sin\phi}$ turns out to be significantly non zero in all the kinematic bins explored. Furthermore, the results from the two sets show consistent magnitude for the moments extracted, suggesting that no effect from the nuclear background present in the 3NH target can be appreciated.

4 Di-hadron in the 12-GeV era

The upgraded CEBAF will provide the four experimental halls an electron beam with an increased energy, that, for Hall-B, can arrive at 11GeV . Hall-B equipment is presently being upgraded to a new detector, *CLAS12*, meant to perform high-acceptance, high-luminosity measurements. Furthermore, more targets will be available for CLAS12 operation. In particular, beyond the standard unpolarized and longitudinally-polarized targets already exploited in the 6 GeV measurements, a transversely-polarized target will be available. Di-hadron SIDIS is already part of the JLab12 physics program. In particular, the simultaneous presence of an unpolarized, a longitudinally-polarized and a transversely-polarized target, operating with the longitudinally-polarized CEBAF electron beam, will allow to access the full set of Structure Functions describing the di-hadron SIDIS cross-section. Di-hadron SIDIS A_{UT} is of particular interest, since it provides an access to the transversity $h_1(x)$ complementary to the single-hadron one, and one proposal has been devoted to the extraction of this observables, whose projections are reported in Fig. 2. The projections show the statistical power of the CLAS12 data, that will be able to explore the A_{UT} dependence on x_B , z and $m_{\pi^+\pi^-}$ in a wide kinematical range and with a high accuracy. It is important to stress here that CLAS12 will extend the constraint on $h_1(x)$ to the *high-x* (valence) region, where no data are presently available, that is also essential to constraint the nucleon tensor charge.




Figure 2. Projections for CLAS12 for the A_{UT} dependence on z (left plot), $m_{\pi^+\pi^-}$ (middle plot) and x_B (right plot). These projections corresponds to 110 days of data taking.

5 Conclusions

Di-hadron SIDIS processes constitute the easiest channel to access the leading-twist pdf $h_1(x)$ and the higher-twist pdfs $e(x)$, $h_L(x)$. These last two PDFs can be accessed by extracting longitudinal Beam and Target-Spin Asymmetries. CLAS preliminary measurements show a significantly non-zero A_{LU} both on 2H and 3NH . This extraction represents the first measurement of A_{LU} and, combining the results presented here with the recently extracted $H_1^{\zeta q}$ [6] is possible, in principle, to get information on $e(x)$. Measurements of the Target and of the Double-Spin Asymmetries are also in progress, together with the extension of the analysis to channels with different pion combinations in the final state, *e. g.* $ep \rightarrow e\pi^+\pi^0X$. Di-hadron SIDIS is an important part of the 12 GeV physics program of JLab, not only in the Hall-B, where measurements aiming at the extraction of the full set of Structure Functions for this process have been proposed, but also in Hall-A, where a letter of intent has been submitted by the SoLID Collaboration, proposing a measurement of A_{UT} on a neutron target. The combination of the CLAS12 (proton) and SoLID (neutron) measurements will allow to operate a flavour separation, leading, eventually, to the extraction of $h_1(x)$ for the u and d flavours.

References

- [1] Jaffe R. L. *et al.*, *Phys. Rev. Lett.* **67**, 552 (1991).
- [2] Jaffe R. L. *et al.*, *Nucl. Phys. B* **375**, 527 (1992).
- [3] A. Bacchetta, *Ph. D. thesis*, hep-ph/0212025.
- [4] A. Bacchetta *et al.*, *JHEP* 02 (2007) 093, hep-ph/0611265.
- [5] B. A. Mecking *et al.*, *Nucl. Instr. Meth. A* **503** 513 (2003).
- [6] Courtoy, Aureore and Bacchetta, Alessandro and Radici, Marco and Bianconi, Andrea *First extraction of interference fragmentation functions from e^+e^- data* *Phys. Rev. D* **85**, 114023 2012.
- [7] The BELLE Collaboration, A. Vossen *et al.*, *Phys. Rev. Lett.* **107**, 072004, (2011) (arXiv:1104.2425).