

Electroexcitation of Nucleon Resonances

Ralf W. Gothe

UNIVERSITY OF
SOUTH CAROLINA

Baryons 2016

International Conference on the Structure of Baryons
May 16 - 20, Tallahassee, Florida


- **γ NN* Vertexcouplings:** A unique exploration of baryon and quark structure?
- **Analysis and New Results:** Phenomenological but consistent!
- **Outlook:** New experiments with extended scope and kinematics!
- **QCD based Theory:** Can we solve non-perturbative QCD and confinement?

Spectroscopy

Build your Mesons and Baryons ...

Three Generations of Matter (Fermions)


	I	II	III	
mass →	2.4 MeV	1.27 GeV	171.2 GeV	0
charge →	$\frac{2}{3}$	$\frac{2}{3}$	$\frac{2}{3}$	0
spin →	$\frac{1}{2}$	$\frac{1}{2}$	$\frac{1}{2}$	1
name →	u up	c charm	t top	γ photon
	4.8 MeV	104 MeV	4.2 GeV	0
	$-\frac{1}{3}$	$-\frac{1}{3}$	$-\frac{1}{3}$	0
	$\frac{1}{2}$	$\frac{1}{2}$	$\frac{1}{2}$	1
Quarks	d down	s strange	b bottom	g gluon
	<2.2 eV	<0.17 MeV	<15.5 MeV	91.2 GeV
	0	0	0	0
	$\frac{1}{2}$	$\frac{1}{2}$	$\frac{1}{2}$	1
	ν_e electron neutrino	ν_μ muon neutrino	ν_τ tau neutrino	Z⁰ weak force
	0.511 MeV	105.7 MeV	1.777 GeV	80.4 GeV
	-1	-1	-1	± 1
	$\frac{1}{2}$	$\frac{1}{2}$	$\frac{1}{2}$	1
Leptons	e electron	μ muon	τ tau	W[±] weak force




N and Δ Excited Baryon States ...

Simon Capstick

➤ Orbital excitations
(two distinct kinds in contrast to mesons)



➤ Radial excitations
(also two kinds in contrast to mesons)




Quark Model Classification of N*

Lowest Baryon Supermultiplets
 SU(6)xO(3) Symmetry

Dietmar Menze

Particle Data Group
 ■ ****
 ■ ***
 ■ **



$\Delta(1232)$

$D_{13}(1520)$
 $S_{11}(1535)$


Roper $P_{11}(1440)$

+ q^3g
 + $q^3q\bar{q}$
 + N-Meson
 + ...

- q^2q
 - ...

Quark Model Classification of N^*


BnGa energy-dependent coupled-channel PWA of CLAS $K^+\Lambda$ and other data



N/Δ Spectrum in RPP 2012

N^*	$J^P (L_{2I,2J})$	2010	2012	Δ	$J^P (L_{2I,2J})$	2010	2012
p	$1/2^+ (P_{11})$	****	****	$\Delta(1232)$	$3/2^+ (P_{33})$	****	****
n	$1/2^+ (P_{11})$	****	****	$\Delta(1600)$	$3/2^+ (P_{33})$	***	***
$N(1440)$	$1/2^+ (P_{11})$	****	****	$\Delta(1620)$	$1/2^- (S_{31})$	****	****
$N(1520)$	$3/2^- (D_{13})$	****	****	$\Delta(1700)$	$3/2^- (D_{33})$	****	****
$N(1535)$	$1/2^- (S_{11})$	****	****	$\Delta(1750)$	$1/2^+ (P_{31})$	*	*
$N(1650)$	$1/2^- (S_{11})$	****	****	$\Delta(1900)$	$1/2^- (S_{31})$	**	**
$N(1675)$	$5/2^- (D_{15})$	****	****	$\Delta(1905)$	$5/2^+ (F_{35})$	****	****
$N(1680)$	$5/2^+ (F_{15})$	****	****	$\Delta(1910)$	$1/2^+ (P_{31})$	****	****
$N(1685)$			*				
$N(1700)$	$3/2^- (D_{13})$	***	**	$\Delta(1920)$	$3/2^+ (P_{33})$	***	**
$N(1710)$	$1/2^+ (P_{11})$	***	**	$\Delta(1930)$	$5/2^- (D_{35})$	***	**
$N(1720)$	$3/2^+ (P_{13})$	****	****	$\Delta(1940)$	$3/2^- (D_{33})$	*	**
$N(1860)$	$5/2^+$		**				
$N(1875)$	$3/2^-$		***				
$N(1880)$	$1/2^+$		**				
$N(1895)$	$1/2^-$		**				
$N(1900)$	$3/2^+ (P_{13})$	**	***	$\Delta(1950)$	$7/2^+ (F_{37})$	****	****
$N(1990)$	$7/2^+ (F_{17})$	**	**	$\Delta(2000)$	$5/2^+ (F_{35})$	**	**
$N(2000)$	$5/2^+ (F_{15})$	**	**	$\Delta(2150)$	$1/2^- (S_{31})$	*	*
$N(2080)$	D_{13}	**		$\Delta(2200)$	$7/2^- (G_{37})$	*	*
$N(2090)$	S_{11}	*		$\Delta(2300)$	$9/2^+ (H_{39})$	**	**
$N(2040)$	$3/2^+$		*				
$N(2060)$	$5/2^-$		**				
$N(2100)$	$1/2^+ (P_{11})$	*	*	$\Delta(2350)$	$5/2^- (D_{35})$	*	*
$N(2120)$	$3/2^-$		**				
$N(2190)$	$7/2^- (G_{17})$	****	****	$\Delta(2390)$	$7/2^+ (F_{37})$	*	*
$N(2200)$	D_{15}	**		$\Delta(2400)$	$9/2^- (G_{39})$	**	**
$N(2220)$	$9/2^+ (H_{19})$	****	****	$\Delta(2420)$	$11/2^+ (H_{3,11})$	****	****
$N(2250)$	$9/2^- (G_{19})$	****	****	$\Delta(2750)$	$13/2^- (I_{3,13})$	**	**
$N(2600)$	$11/2^- (I_{1,11})$	***	**	$\Delta(2950)$	$15/2^+ (K_{3,15})$	**	**
$N(2700)$	$13/2^+ (K_{1,13})$	**	**				


High-statistics and high-precision photoproduction data from JLAB, MAMI, ELSA, GRAAL



Are we observing parity doublets with the new states or not?

V. Crede & W. Roberts, Rep. Prog. Phys. 76 (2013)

New FROST Results from $\vec{\gamma}\vec{p} \rightarrow \pi^+n$




- FROST experiment produced 900 data points of the **double-polarization observable E** in π^+ photoproduction with circularly polarized beam on longitudinally polarized protons for $W = 1240 - 2260$ MeV.
- Significant improvements of the description of the data in SAID, Jülich, and BnGa partial-wave analyses after fitting.
- **New evidence found in this data for a $\Delta(2200)7/2^-$ resonance (BnGa analysis).**

S. Strauch *et al.*, Phys. Let. B **750** (2015) 53 and A.V. Anisovich *et al.*, arXiv:1503.05774

N* Spectrum in LQCD

The strong interaction physics is encoded in the nucleon excitation spectrum that spans the degrees of freedom from meson-baryon and dressed quarks to elementary quarks and gluons.




LQCD predicts states with the same quantum numbers as CQMs with underlying $SU(6) \times O(3)$ symmetry.


R. Edwards *et al.*,
arXiv:1104.5152, 1201.2349

Transition Form Factors


Hadron Structure with Electromagnetic Probes




- Study the structure of the nucleon spectrum in the domain where dressed quarks are the major active degree of freedom.
- Explore the formation of excited nucleon states in interactions of dressed quarks and their emergence from QCD.




Hadron Structure with Electromagnetic Probes





- Study the structure of the nucleon spectrum in the domain where dressed quarks are the major active degree of freedom.
- Explore the formation of excited nucleon states in interactions of dressed quarks and their emergence from QCD.




Baryon Excitations and Quasi-Elastic Scattering





hard and
confined



quasi-elastic



hard



soft


Deep Inelastic Scattering
S. Stein et al., PR **D22** (1975) 1884

Structure Analysis of the Baryon


Demolition of a chimney at the "Henninger Brewery" in Frankfurt am Main, Germany, on 2 December 2006




hard and confined



hard and soft



quasi-elastic



$\gamma_{\nu} \text{NN}^*$


Extraction

Data-Driven Data Analyses

Consistent Results





- Single meson production:
Unitary Isobar Model (UIM)
Fixed- t Dispersion Relations (DR)
- Double pion production:
Unitarized Isobar Model (JM)
- Coupled-Channel Approach:
EBAC \Rightarrow Argonne-Osaka
JAW \Rightarrow Jülich-Athens-Washington
BoGa \Rightarrow Bonn-Gatchina



Int. J. Mod. Phys. E, Vol. 22, 1330015 (2013) 1-99

Electrocouplings of $N(1440)P_{11}$ from CLAS Data




□ PDG
 ● $N\pi$ (UIM, DR)
 ▲ $N\pi\pi$ (JM) 2012
 ■ $N\pi\pi$ (JM) preliminary

Consistent results obtained in the low-lying resonance region by independent analyses in the exclusive $N\pi$ and $p\pi^+\pi^-$ final-state channels – that have fundamentally different mechanisms for the nonresonant background – underscore the capability of the reaction models to extract reliable resonance electrocouplings.


Phys. Rev. C 80, 055203 (2009) 1-22 and Phys. Rev. C 86, 035203 (2012) 1-22

Transition Form Factors and QCD Models

Roper resonance $P_{11}(1440)$



PDG 2013 update





- $A_{1/2}$ has zero-crossing near $Q^2=0.5$ and becomes dominant amplitude at high Q^2 .
- Consistent with radial excitation at high Q^2 and large meson-baryon coupling at small Q^2 .
- Eliminates gluonic excitation (q^3G) as a dominant contribution.

Nick Tyler closes the 1-2 GeV^2 gap for single pion production.

Transition Form Factors and QCD Models

Roper resonance $P_{11}(1440)$

Si-xue Qin *et al.*, Phys. Rev. C 84 (2011) 042202(R)




- $A_{1/2}$ has zero-crossing near $Q^2=0.5$ and becomes dominant amplitude at high Q^2 .
- Consistent with radial excitation at high Q^2 and large meson-baryon coupling at small Q^2 .
- Eliminates gluonic excitation (q^3G) as a dominant contribution.

New proposal on electroexcited gluon hybrids to be submitted to PAC44

N* Spectrum in LQCD

The strong interaction physics is encoded in the nucleon excitation spectrum that spans the degrees of freedom from meson-baryon and dressed quarks to elementary quarks and gluons.




LQCD predicts hybrid baryon states replicating the negative parity multiplet structure.

New proposal on electroexcited gluon hybrids submitted to PAC44


Evidence for the Onset of Precocious Scaling?

I. G. Aznauryan *et al.*, Phys. Rev. C80, 055203 (2009)




➤ $A_{1/2} \propto 1/Q^3$

➤ $A_{3/2} \propto 1/Q^5$



Evidence for the Onset of Precocious Scaling?


Ye Tian



V. Mokeev, userweb.jlab.org/~mokeev/resonance_electrocouplings/ (2016)




Evidence for the Onset of Precocious Scaling?

Ye Tian



V. Mokeev, userweb.jlab.org/~mokeev/resonance_electrocouplings/ (2016)

Electrocouplings of $N(1520)D_{13}$ and $N(1535)S_{11}$



— Argonne Osaka / EBAC DCC MB dressing
(absolute values)


— E. Santopinto, M. Giannini, hCQM
PRC 86, 065202 (2012)

— S. Capstick, B.D. Keister (rCQM)
PRD51, 3598 (1995)

■ $\pi^+\pi^-p$ 2012 ▲ $\pi^+\pi^-p$ 2010 ● $N\pi$ 2009

▲ ηp
CLAS/Hall-C


Evidence for the Onset of Precocious Scaling?



➤ $A_{1/2} \propto 1/Q^3$


➤ $A_{3/2} \propto 1/Q^5$

➤ $G_M^* \propto 1/Q^4$




$q^2 q$


I. G. Aznauryan *et al.*, Phys. Rev. C80, 055203 (2009)



N → Δ Multipole Ratios R_{EM} , R_{SM}



Phys. Rev. Lett. 97, 112003 (2006)



➤ New trend towards pQCD behavior **does not** show up

➤ $R_{EM} \rightarrow +1$ $R_{SM} \rightarrow \text{const}$


➤ $G_{M,J-S}^* \rightarrow 1/Q^4$ $G_{M,Ash}^* \rightarrow 1/Q^5$

➤ CLAS12 can measure G_M^* , R_{EM} , and R_{SM} up to $Q^2 \sim 12 \text{ GeV}^2$

N(1520)D₁₃ Helicity Asymmetry


L. Tiator

$$A_{\text{hel}} = \frac{A_{1/2}^2 - A_{3/2}^2}{A_{1/2}^2 + A_{3/2}^2}$$



γNN^* Helicity Asymmetries


Ye Tian




V. Mokeev, userweb.jlab.org/~mokeev/resonance_electrocouplings/ (2016)

Interplay between Meson-Baryon Cloud and Quark Core

N(1520)3/2-



N(1675)5/2-



..... Argonne-Osaka MB dressing (absolute values)

— E. Santopinto and M. Giannini, PRC 86 (2012) 065202

The almost direct access to

- quark core from the data on N(1520)3/2-
- meson-baryon cloud from the data on N(1675)5/2-


sheds light on the transition from the confined quark to the colorless meson-baryon structure and its dependents on the N* quantum numbers.

New Experimental Results & Approaches

Higher-Lying Resonance Electrocouplings

N(1680)F₁₅

Kijun Park



▲ RPP (PDG) Phys. Rev. D 86 (2012)

◻ M. Dugger Phys. Rev. C 76 (2007)

◻ I.G. Aznauryan, Phys. Rev. C 72 (2005)

△ $N\pi\pi$: V. Mokeev (JM)

● $N\pi$: I.G. Aznauryan (UIM & DR)

K. Park *et al.*, Phys. Rev. C 91, 045203 (2015)

--- D. Merten, U. Löring *et al.*


- · - · - Z. Lee and F. Close

— E. Santopinto and M.M. Gianini

Higher-Lying Resonance Electrocouplings

$N(1675)D_{15}$

Kijun Park




K. Park *et al.*, Phys. Rev. C **91**, 045203 (2015)

- ▲ RPP (PDG) Phys. Rev. D **86** (2012)
- M. Dugger Phys. Rev. C **76** (2007)
- $N\pi$: I.G. Aznauryan (UIM & DR)

- D. Merten, U. Löring et al.
- · - · - B. Julia-Diaz, T.-S.H. Lee et al.
- E. Santopinto and M.M. Gianini

Higher-Lying Resonance Electrocouplings

Viktor Mokeev



Independent fits in different W-intervals

green: $1.46 < W < 1.56 \text{ GeV}$

magenta: $1.56 < W < 1.66 \text{ GeV}$

red: $1.61 < W < 1.71 \text{ GeV}$

blue: $1.66 < W < 1.76 \text{ GeV}$

black: $1.71 < W < 1.81 \text{ GeV}$

result in consistent electrocouplings and hence offer sound evidence for their reliable extraction.

The $\pi^+\pi^-p$ electroproduction channel provides first preliminary results on the $\Delta(1620)1/2^-$, $N(1650)1/2^-$, $N(1680)5/2^+$, $\Delta(1700)3/2^-$, and $N(1720)3/2^+$ electrocouplings with good accuracy.

V. Mokeev *et al.*, Phys. Rev. C **93**, 025206

New $N'(1720)3/2^+$ State and its Properties

N^* hadronic decays from JM15 that incorporates $N'(1720)3/2^+$

Resonance	BF($\pi\Delta$), %	BF(ρp), %
$N'(1720)3/2^+$ electroproduction photoproduction	47-64 46-62	3-10 4-13
$N(1720)3/2^+$ electroproduction photoproduction	39-55 38-53	23-49 31-46
$\Delta(1700)3/2^-$ electroproduction photoproduction	77-95 78-93	3-5 3-6

A successful description of $\pi^+\pi^-p$ photo- and electro-production cross sections at $Q^2=0, 0.65, 0.95,$ and 1.30 GeV^2 has been achieved by implementing a new $N'(1720)3/2^+$ state with Q^2 -independent hadronic decay widths of all resonances that contribute at $W \sim 1.7 \text{ GeV}$, that allows us to claim the existence of a new $N'(1720)3/2^+$ state.

Mass: 1.715-1.735 GeV




Width: 120 6 MeV

■ $N'(1720)3/2^+$

Mass: 1.743-1.753 GeV


Width: 112 8 MeV

▲ $N(1720)3/2^+$



ϕ -dependent $N_{\pi\pi}$ Single-Differential Cross Sections

Q^2, W bin = $[1.25, 1.75) \text{ GeV}^2, [1.625, 1.650) \text{ GeV}$ Arjun Trivedi




$$\left(\frac{d^2\sigma}{dX^{ij}d\phi^j} \right) = R2_T X^{ij} + R2_I X^{ij} + R2_{LT} X^{ij} \cos \phi_j + R2_{TT} X^{ij} \cos 2\phi_j$$

Chris McLauchlin extracts the **beam helicity dependent** differential cross sections.

ϕ -dependent $N\pi\pi$ Single-Differential Cross Sections





Q^2, W bin = $[1.25, 1.75)\text{GeV}^2, [1.625, 1.650)\text{GeV}$ Arjun Trivedi



$$\left(\frac{d^2\sigma}{dX^{ij}d\phi^j} \right) = R2_T^{X_{ij}} + R2_L^{X_{ij}} + \underline{R2_{LT}^{X_{ij}} \cos \phi_j} + \underline{R2_{TT}^{X_{ij}} \cos 2\phi_j}$$


Single π Electroproduction off the Deuteron

Ye Tian



Single π Electroproduction off the Deuteron


Ye Tian



Below a missing momentum of 0.2 GeV the **measured data** coincides with the **resolution smeared theoretical Fermi momentum distribution**.

Single π Electroproduction off the Deuteron

Ye Tian



Gary Hollis inclusive of the **proton** in the Deuteron **with correction of Fermi smearing**.

FSI for $\gamma n \rightarrow \pi^- p$

[V. Tarasov, A. Kudryavtsev, W. Briscoe, H. Gao, IS, Phys Rev C 84, 035203 (2011)]

$$R_{FSI} = (d\sigma/d\Omega_{\pi p}) / (d\sigma^{IA}/d\Omega_{\pi p})$$


Cuts:

$p_s > 200 \text{ MeV}/c$
 $p_f > 200 \text{ MeV}/c$

CLAS data:

$E > 1 \text{ GeV}$
 $\theta > 32 \text{ deg}$

• There is no large sensitivity to cuts.



• For CLAS data

- The FSI correction factor $R < 1$.
- The behavior is smooth vs. θ .
- The effect $\Delta\sigma/\sigma \leq 10\%$.

• There is a sizeable FSI effect from S-wave part of pp-FSI at small angles.

• This region narrows as the E_γ increases.

• Our estimation of the Glauber FSI corrections gives the value of 5%.

• Previous estimations gave the order of 15-30%.




--- $[IA + NN_{fsi}] / IA$
 — $[IA + (NN + \pi N)_{fsi}] / IA$



Single π Electroproduction off the Deuteron

Ye Tian




Single π Electroproduction off the Deuteron

Ye Tian

$W = 1125$ MeV

$\Delta W = 25$ MeV



$Q^2 = 0.7$ GeV²

$\Delta Q^2 = 0.2$ GeV²

$W = 1685$ MeV

Single π Electroproduction off the Deuteron

Ye Tian


$W = 1125$ MeV

$\Delta W = 25$ MeV

$W = 1685$ MeV

$Q^2 = 0.5$ GeV²

$Q^2 = 0.9$ GeV²



$Q^2 = 0.7$ GeV²

$\Delta Q^2 = 0.2$ GeV²



**Kinematic FSI
Corrections**

Single π^- Electroproduction off the Deuteron

Ye Tian



$W = 1212 \text{ MeV}$

$\Delta W = 25 \text{ MeV}$



$Q^2 = 0.5 \text{ GeV}^2$



$\Delta Q^2 = 0.2 \text{ GeV}^2$



$\cos(\theta) = -0.7$

$\Delta \cos(\theta) = 0.2$



$\cos(\theta) = 0.7$



$\phi = 15^\circ$

$\Delta \phi = 30^\circ$


$\phi = 345^\circ$



Double π Electroproduction off the Deuteron

Iuliia Skorodina


P_X of $ep(n) \rightarrow e'p'(n)\pi^+\pi^-$



Double π Electroproduction off the Deuteron

Iuliia Skorodina

P_X of $ep(n) \rightarrow e'p'(n)\pi^+\pi^-$




QCD-Based Models and Theory

For some highlighted examples see posted presentation or
Int. J. Mod. Phys. E, Vol. 22, 1330015 (2013)

DSE and EBAC/ANL-Osaka Approaches

Semi-quantitative agreement with the first DSE 3-dressed-quark core radial excitation




Int. J. Mod. Phys. E, Vol. 22, 1330015 (2013)

Re (E) (MeV)

Dyson-Schwinger Equation (DSE) Approach

DSE approaches provide links between dressed quark propagators, form factors, scattering amplitudes, and QCD.



N* electrocouplings can be determined by applying Bethe-Salpeter / Faddeev equations to 3 dressed quarks while the properties and interactions are derived from QCD.

DSE calculations of elastic and transition form factors are very sensitive to the momentum dependence of the dressed-quark propagator.


I.C. Cloet *et al.*, Phys. Rev. Lett. **111**, 101803

DSE electrocouplings of several excited nucleon states will become available as part of the commitment of the Argonne NL.

Int. J. Mod. Phys. E, Vol. 22, 1330015 (2013) 1-99

Dyson-Schwinger Equation (DSE) Approach

DSE approaches provide links between dressed quark propagators, form factors, scattering amplitudes, and QCD.



N^* electrocouplings can be determined by applying Bethe-Salpeter / Faddeev equations to 3 dressed quarks while the properties and interactions are derived from QCD.

Impact of a modified momentum dependence of the dressed-quark propagator.


I.C. Cloet *et al.*, Phys. Rev. Lett. **111**, 101803

DSE electrocouplings of several excited nucleon states will become available as part of the commitment of the Argonne NL.

Int. J. Mod. Phys. E, Vol. 22, 1330015 (2013) 1-99

Anomalous Magnetic Moment in DSE Approach


J. Segovia



- contact interaction
- sophisticated interaction
- - - with momentum dependent κ
- == renormalized at real photon point

J. Segovia *et al.*, FBS 55 (2014) 1185-1222



The DSE calculation of R_{EM} zero crossing is sensitive to the momentum dependent anomalous magnetic moment of the dressed-quark.



Roper Transition Form Factors in DSE Approach


$N(1440)P_{11}$

J. Segovia *et al.*, Phys. Rev. Lett. **115**, 171801



DSE Contact $x=Q^2/m_N^2$
 DSE Realistic
 Inferred meson-cloud contribution
 Anticipated complete result



Importantly, the existence of a zero in F_2 is not influenced by meson-cloud effects, although its precise location is.



Roper Transition Form Factors in LQCD

$N(1440)P_{11}$

Huey-Wen Lin and S.D. Cohen





Lattice QCD calculations of the $p(1440)P_{11}$ transition form factors have been carried out with various pion masses, $m_\pi = 390$, 450 , and 875 MeV. Particularly remarkable is the zero crossing in F_2 that appears at the current statistics in the unquenched but not in the quenched calculations. This might suggest that at low Q^2 the pion-cloud dynamics are significant in full QCD.

LQCD calculations of N^* electrocouplings will be extended to $Q^2 = 10$ GeV² near the physical π -mass as part of the commitment of the JLab LQCD and EBAC groups in support of this proposal.

Int. J. Mod. Phys. E, Vol. 22, 1330015 (2013) 1-99

LQCD & Light Cone Sum Rule (LCSR) Approach

N(1535)S₁₁



LQCD is used to determine the moments of N* distribution amplitudes (DA) and the N* electrocouplings are determined from the respective DAs within the LCSR framework.

Calculations of N(1535)S₁₁ electrocouplings at Q^2 up to 12 GeV² are already available and shown by shadowed bands on the plot.


LQCD & LCSR electrocouplings of others N* resonances will be evaluated as part of the commitment of the University of Regensburg group.

Int. J. Mod. Phys. E, Vol. 22, 1330015 (2013) 1-99

CLAS12



- Luminosity $> 10^{35} \text{ cm}^{-2}\text{s}^{-1}$
- Hermeticity
- Polarization


- Baryon Spectroscopy
- Elastic Form Factors
- N to N* Form Factors
- GPDs and TMDs
- DIS and SIDIS
- Nucleon Spin Structure
- Color Transparency
- ...




New Forward Time of Flight Detector for CLAS12

ToF12 Time Resolution Measurements






Torus, LTCC, and FTOF Fully Installed



Anticipated N^* Electrocouplings from Combined Analyses of $N\pi/N\pi\pi$



Open circles represent projections and all other markers the available results with the 6-GeV electron beam

➤ Examples of **published and projected results** obtained within 60d for three prominent excited proton states from analyses of $N\pi$ and $N\pi\pi$ electroproduction channels. Similar results are expected for many other resonances at higher masses, e.g. $S_{11}(1650)$, $F_{15}(1685)$, $D_{33}(1700)$, $P_{13}(1720)$, ...

➤ The approved CLAS12 experiments E12-09-003 (NM, $N\pi\pi$) and E12-06-108A (KY) are currently **the only experiments** that can provide data on $\gamma_v NN^*$ electrocouplings for almost all well established excited proton states at the highest photon virtualities ever achieved in N^* studies up to Q^2 of 12 GeV^2 , see <http://boson.physics.sc.edu/~gothe/research/pub/whitepaper-9-14.pdf>.

Summary

- First high precision photo- and electroproduction data have become available and led to a new wave of significant developments in reaction and QCD-based theories.
- New high precision hadro-, photo-, and electroproduction data off the proton and the neutron will stabilize coupled channel analyses and expand the validity of reaction models, allowing us to
 - investigate and search for baryon hybrids,
 - establish a repertoire of high precision spectroscopy parameters, and
 - measure light-quark-flavor separated electrocouplings over an extended Q^2 -range, both to lower and higher Q^2 , for a wide variety of N^* states.
- Comparing these results with DSE, LQCD, LCSR, and rCQM will build further insights into
 - the strong interaction of dressed quarks and their confinement,
 - the emergence of bare quark dressing and dressed quark interactions from QCD, and
 - the QCD β -function and the origin of 98% of nucleon mass.
- A close collaboration of experimentalists and theorists has formed and is needed to push these goals, see Review Article *Int. J. Mod. Phys. E*, Vol. 22, 1330015 (2013) 1-99, that shall lead to a QCD theory that describes the strong interaction from current quarks to nuclei. **ECT*2015 and INT2016.**

