


Radiative and Hadronic Decay modes of the η -Meson with CLAS and WASA-at-COSY

17.05.2016 | Baryons 2016

The η -Meson: A unique tool

- $m_\eta = 0.5478 \text{ GeV}/c^2$
 - $\Gamma_\eta = (1.31 \pm 0.05) \text{ keV}$
 - $\bar{\tau} \approx 5 \cdot 10^{-19} \text{ s}$
 - $J^{PC} = 0^{-+} \implies \eta\text{-meson is:}$
 C -, P -, G - and CP - eigenstate
 - All strong and electromagnetic decays are forbidden to first order
- ⇒ **Access to rare decay processes**



Focus of this talk:


- 1 $\eta \rightarrow \pi^+ \pi^- \pi^0$ || Measured in $\gamma p \rightarrow p \eta [\eta \rightarrow \dots]$ with CLAS
- 2 $\eta \rightarrow \pi^+ \pi^- \gamma$ || Measured in $p p \rightarrow p p \eta [\eta \rightarrow \dots]$ with WASA

η -Decays with CLAS and WASA-at-COSY

Decay mode	Issue
$\eta' \rightarrow \pi^+ \pi^- \eta$	Dalitz plot analysis
$\eta \rightarrow \pi^+ \pi^- \pi^0$	Dalitz plot analysis
$\eta^{(')} \rightarrow \pi^+ \pi^- \gamma$	Box anomaly, $\pi^+ \pi^-$ FSI
$\eta^{(')} \rightarrow e^+ e^- \gamma$	Single-off-shell transition form factor
$\eta^{(')} \rightarrow \pi^+ \pi^- e^+ e^-$	CP-Violation
$\eta \rightarrow e^+ e^- e^+ e^-$	Double-off-shell transition form factor
$\eta \rightarrow \pi^0 e^+ e^-$	C-Violation

- CLAS: ([Thomas Jefferson Laboratory, Newport News, USA](#))
 - Access to variety of light meson decays via: $\gamma p \rightarrow pX$
 - Light Meson Decay program with focus on: π^0 , η , ω , η' and Φ -decays
- WASA-at-COSY: ([Forschungszentrum Jülich, Germany](#))
 - η -Decay program with data sets: $pd \rightarrow {}^3\text{He}\eta$ and $pp \rightarrow pp\eta$
 - Analyses of π^0 - and ω -decays were also performed

$\eta \rightarrow \pi^+ \pi^- \pi^0$ The Dalitz Plot



(a) KLOE coll., JHEP, 05, (2008)

Dalitz plot variables:

$$X = \sqrt{3} \frac{T_{\pi^+} - T_{\pi^-}}{T_{\pi^+} + T_{\pi^-} + T_{\pi^0}}$$

$$Y = \frac{3T_{\pi^0}}{T_{\pi^+} + T_{\pi^-} + T_{\pi^0}}$$

- Decay via strong isospin violation: $\Gamma_{meas} = \left(\frac{Q_D}{Q}\right)^4 \bar{\Gamma}$
 - $Q^2 = \frac{m_s^2 - \hat{m}^2}{m_d^2 - m_u^2}$, $\hat{m} = \frac{1}{2}(m_u + m_d)$
 - $\bar{\Gamma}$ calculated with ChPT at Dashen limit: $Q_D = 24.2$
- Dalitz plot analysis: $\frac{d^2\Gamma}{dXdY} \propto (1 + aY + bY^2 + dX^2 + fY^3 + gX^2Y + \dots)$
 $\rightarrow c, e$ and h would imply C-violation

$\eta \rightarrow \pi^+ \pi^- \pi^0$ Recent Measurements
and Theoretical Predictions

Parameter:		-a	b	d	f
Theor.	ChPT (NNLO) ^(b)	1.271(75)	0.394(102)	0.055(57)	0.025(160)
	NREFT ^(c)	1.213(14)	0.308(23)	0.050(3)	0.083(19)
	PWA ^(e)	1.116(32)	0.188(12)	0.063(4)	0.091(3)
Exp.	KLOE ^(a)	1.090(5)(⁺⁸ ₋₁₉)	0.124(6)(10)	0.057(6)(⁺⁷ ₋₁₆)	0.14(1)(2)
	WASA ^(d)	1.144(18)	0.219(19)(47)	0.086(18)(15)	0.115(37)

(a) KLOE coll., *JHEP*, 05, (2008)


(b) J. Bijnens and K. Ghorbani., *JHEP*, 11, (2007)

(c) S- P. Schneider et al., *JHEP*, 028, (2011)


(d) WASA-at-COSY coll., *Phys. Rev.*, C90(045207), 2014

(e) Peng Guo et al., *Phys. Rev.*, D92(05016), (2015)


- $\approx 120\text{ k}$ $\eta \rightarrow \pi^+ \pi^- \pi^0$ events in the final WASA-at-COSY event sample
- Partial Wave Analysis from JPAC group:
 $Q = 21.4 \pm 0.4_{\text{stat}}^{(\text{e})}$
- Dalitz plot analysis for
 $\gamma p \rightarrow p \eta [\eta \rightarrow \pi^+ \pi^- \pi^0]$ with CLAS



CEBAF Large Acceptance Spectrometer - CLAS





The CLAS $\gamma p \rightarrow pX$ Data Set



- Photon beam: $E_{\gamma,\text{beam}} \in [1.1 \text{ GeV}, 5.45 \text{ GeV}]$
- (Main) Contributions from:
 - Direct pion production (e.g. $\gamma p \rightarrow p\pi^+\pi^-\pi^0$)
 - π^0 , η , ω and ρ decays
- Use missing mass: $|\mathbf{P}_{\gamma,\text{beam}} + \mathbf{P}_{\text{target}} - \mathbf{P}_{\text{proton}}|$ to monitor analysis

$\eta \rightarrow \pi^+ \pi^- \pi^0$ Analysis and Results




→ Use kinematic fit and kinematic limits of $M(\pi^+, \pi^-)$ to suppress background

- $\approx 770 \text{ k}$ $\eta \rightarrow \pi^+ \pi^- \pi^0$ events reconstructed for 2/3 of CLAS data set
- Performed side-band subtraction for each Dalitz Plot bin
- To do:
 - Efficiency correction and correction for $\eta \rightarrow \pi^+ \pi^- \gamma$ events
 - Determine Dalitz Plot parameters and Q

Small Outlook: Dalitz Plot Analysis for $\omega \rightarrow \pi^+ \pi^- \pi^0$ with WASA-at-COSY

- Theoretical description of this decay:
VMD Model, Lagrangian Approach^(a), Dispersive Analysis^{(b),(c)}
 \rightarrow Input from experiment needed
- Look at: $\frac{d^2\Gamma}{dZd\Phi} \propto (1 + 2\alpha Z + 2\beta Z^{3/2} \sin^3 \Phi + \mathcal{O}(Z^2) + \dots)$
- Analysis ongoing^{(d),(e)} for the reaction:
 $pd \rightarrow {}^3\text{He}\omega[\omega \rightarrow \pi^+ \pi^- \pi^0]$ and $pp \rightarrow pp\omega[\omega \rightarrow \pi^+ \pi^- \pi^0]$



(a) S. Leupold et al., Eur. Phys. J. **A** 39, 205-212, (2009)

(b) N. Niemann et al., Eur. Phys. J. **C** 39, 2014, (2012)

(c) I.V.Danilkin et al., Phys. Rev. **D91**, 094029, (2015)


(d) PhD.-Project of Lena Heijkenskjöld

(e) PhD.-Project of Siddhesh Sawant

$\eta \rightarrow \pi^+ \pi^- \gamma$

The Box Anomaly and $\pi^+ \pi^-$ FSI

Chiral limit: (a),(b)



- Wess-Zumino-Witten Lagrangian


(a) Wess, Zumino, *Phys. Lett.*, B37(95), 1971

(b) Witten, *Nucl. Phys.*, B223:422-432, 1983

- Decay amplitude $A_{\eta \rightarrow \pi^+ \pi^- \gamma}$ is sensitive to box anomaly^(c):

$$A_{\eta \rightarrow \pi^+ \pi^- \gamma} \propto \frac{e}{4\sqrt{3}\pi^2 F_\pi^3} \left(\frac{F_\pi}{F_8} \cos \theta - \sqrt{2} \frac{F_\pi}{F_0} \sin \theta \right)$$


- $\Gamma^{\text{Theory}}(\eta \rightarrow \pi^+ \pi^- \gamma) = 35.7 \text{ eV}^{(c)}$
- $\Gamma^{\text{Exp.}}(\eta \rightarrow \pi^+ \pi^- \gamma) = (55.3 \pm 2.4) \text{ eV}^{(d)}$
- (c) B.R. Holstein, *Phys. Scripta*, T99:55-67, 2002
- (d) PDG, *Chin. Phys.*, 090001, 2014
- Photon energy distribution E_γ :^(e)
- (e) WASA-at-COSY coll. *Phys. Lett.*, B707:243-249, 2012



$$E_\gamma(s_{\pi\pi}) = \frac{1}{2} \cdot \left(m_\eta - \frac{s_{\pi\pi}}{m_\eta} \right)$$

$\eta \rightarrow \pi^+ \pi^- \gamma$ The Box Anomaly and $\pi^+ \pi^-$ FSI

Beyond chiral limit:



- Wess-Zumino-Witten Lagrangian & $\pi^+ \pi^-$ Final State Interactions
- Modification of decay amplitude:^(a)


(a) F.Stollenwerk et al., *Phys. Lett.*, B707:184-190, 2012

$$A_{\eta \rightarrow \pi^+ \pi^- \gamma} \times [F_{PV}(s_{\pi\pi}) \cdot (1 + \alpha s_{\pi\pi})]$$

$$\Rightarrow \text{Description of FSI: } \begin{cases} \text{by } F_{PV} & \alpha = 0 \\ \text{reaction specific*} & \alpha \neq 0 \end{cases}$$

*Input from theory

- $\Gamma^{\text{Theory}}(\eta \rightarrow \pi^+ \pi^- \gamma) = 35.7 \text{ eV}^{(b)}$
- $\Gamma^{\text{Exp.}}(\eta \rightarrow \pi^+ \pi^- \gamma) = (55.3 \pm 2.4) \text{ eV}^{(c)}$
- (b) B.R. Holstein, *Phys. Scripta*, T99:55-67, 2002
- (c) PDG, *Chin. Phys.*, 090001, 2014
- Photon energy distribution E_γ :^(d)
- (d) WASA-at-COSY coll. *Phys. Lett.*, B707:243-249, 2012



$$E_\gamma(s_{\pi\pi}) = \frac{1}{2} \cdot \left(m_\eta - \frac{s_{\pi\pi}}{m_\eta} \right)$$

$\eta \rightarrow \pi^+ \pi^- \gamma$ Recent Measurements
and Theoretical Predictions

		$\Gamma(\eta \rightarrow \pi^+ \pi^- \gamma) / \Gamma(\eta \rightarrow \pi^+ \pi^- \pi^0)$	$\alpha [\text{GeV}^{-2}]$
Experiment	Gormley et al.	0.202 ± 0.006	1.8 ± 0.4
	Thaler et al.	0.209 ± 0.004	-
	Layter et al.	-	-0.9 ± 0.1
	GAMS-200*	-	2.7 ± 0.1
	CRYSTAL BARREL*	-	1.8 ± 0.53
	CLEO	0.175 ± 0.013	-
	WASA-at-COSY	Preliminary: 0.206 ± 0.011	1.89 ± 0.86
	KLOE	0.1856 ± 0.003	1.32 ± 0.2
	CLAS	Analysis ongoing for η and η'	-
Theory	BESIII	Analysis ongoing for η and η'	-
	N/D	0.2188 ± 0.0088	0.64 ± 0.02
	HLS	0.1875 ± 0.0094	0.23 ± 0.01
	($O(p^5)$) + 1-loop	0.1565 ± 0.0063	-0.7 ± 0.1
	Box anomaly	0.119 ± 0.0048	-1.7 ± 0.02

*Measured $\eta' \rightarrow \pi^+ \pi^- \gamma$

⇒ Determine $\frac{\Gamma(\eta \rightarrow \pi^+ \pi^- \gamma)}{\Gamma(\eta \rightarrow \pi^+ \pi^- \pi^0)}$ and α via E_γ -distribution in $pp \rightarrow pp\eta[\eta \rightarrow \pi^+ \pi^- \gamma]$

$\eta \rightarrow \pi^+ \pi^- \gamma$ Recent Measurements
and Theoretical Predictions


		$\Gamma(\eta \rightarrow \pi^+ \pi^- \gamma) / \Gamma(\eta \rightarrow \pi^+ \pi^- \pi^0)$	$\alpha [\text{GeV}^{-2}]$
Experiment	<i>Phys. Rev.</i> , D2:501-505, 1970	0.202 ± 0.006	1.8 ± 0.4
	<i>Phys. Rev.</i> , D7:2569-2571, 1973	0.209 ± 0.004	-
	<i>Phys. Rev.</i> , D7:2565-2568, 1973	-	-0.9 ± 0.1
	<i>Phys.</i> , C50:451-454, 1991 *	-	2.7 ± 0.1
	<i>Phys. Lett.</i> , B402:195, 1997 *	-	1.8 ± 0.53
	<i>Phys. Rev. Lett.</i> , 99(122001), 2007	0.175 ± 0.013	-
	<i>Phys. Rev. Lett.</i> , B707:243-249, 2013	-	1.89 ± 0.86
	<i>Phys. Lett.</i> , B718:910-914, 2013	0.1856 ± 0.003	1.32 ± 0.2
	-	-	-
	-	-	-
Theory	<i>Phys. Scripta</i> , T99:55-67, 2002	0.2188 ± 0.0088	0.64 ± 0.02
	<i>Europ. Phys. Journal</i> , C31:525-547, 2003	0.1875 ± 0.0094	0.23 ± 0.01
	<i>Phys. Lett.</i> , B237:488-494, 1990	0.1565 ± 0.0063	-0.7 ± 0.1
	<i>Phys. Scripta</i> , T99:55-67, 2002	0.119 ± 0.0048	-1.7 ± 0.02

* Measured $\eta' \rightarrow \pi^+ \pi^- \gamma$

⇒ Determine $\frac{\Gamma(\eta \rightarrow \pi^+ \pi^- \gamma)}{\Gamma(\eta \rightarrow \pi^+ \pi^- \pi^0)}$ and α via E_γ -distribution in $pp \rightarrow pp\eta[\eta \rightarrow \pi^+ \pi^- \gamma]$


Wide Angle Shower Apparatus - WASA

Example: $pp \rightarrow pp\eta [\eta \rightarrow \pi^+\pi^-\gamma]$





The WASA-at-COSY $pp \rightarrow pp\eta$ Data Set

- $\sigma(\eta) = (9.8 \pm 1)\mu\text{b}$ at $T_{beam} = 1.4 \text{ GeV}$
- Background contributions from direct pion production:
 - $pp \rightarrow pp\pi^+\pi^-$ [$\sigma = (660 \pm 100)\mu\text{b}$]
 - $pp \rightarrow pp\pi^0\pi^0$ [$\sigma = (200 \pm 30)\mu\text{b}$]
 - $pp \rightarrow pp\pi^+\pi^-\pi^0$ [$\sigma = (4.6 \pm 1.5)\mu\text{b}$]
- After proton identification and requesting an event topology reconstruct η -Meson via Missing mass: $|\mathbf{P}_{beam} + \mathbf{P}_{target} - (\mathbf{P}_{proton1} + \mathbf{P}_{proton2})|$



$pp \rightarrow pp\eta$ data taken in	2008	2010	2012
Duration of beam time	2 weeks	7 weeks	8 weeks
η mesons produced	$\sim 1 \cdot 10^8$	$\sim 4 \cdot 10^8$	$\sim 5 \cdot 10^8$

$\eta \rightarrow \pi^+ \pi^- \gamma$ Analysis and Results



- Use kinematic fit and rejection of low energetic satellite clusters in calorimeter to suppress background
- $\approx 209 \text{ k } \eta \rightarrow \pi^+ \pi^- \gamma$ events reconstructed
- E_γ -distribution after background correction from direct pion production
- To do:
 - i) Include efficiency corrections
 - ii) Calculate $\frac{\Gamma(\eta \rightarrow \pi^+ \pi^- \gamma)}{\Gamma(\eta \rightarrow \pi^+ \pi^- \pi^0)}$ and α

Summary and Outlook

- η -Decays with CLAS and WASA-at-COSY
(part of the Light Meson Decay group)
- **Dalitz Plot analysis of $\gamma p \rightarrow p\eta[\eta \rightarrow \pi^+\pi^-\pi^0]$:**

- ≈ 770 k events reconstructed in CLAS
- Background corrected Dalitz Plot
- Include simulations into analysis
- Calculate Dalitz Plot parameters and Q

- **Box Anomaly and $\pi^+\pi^-$ FSI in $p\bar{p} \rightarrow p\bar{p}\eta[\eta \rightarrow \pi^+\pi^-\gamma]$:**

- ≈ 209 k events reconstructed in WASA
- Background corrected E_γ -distribution
- Efficiency correction
- Extract α and $\frac{\Gamma(\eta \rightarrow \pi^+\pi^-\gamma)}{\Gamma(\eta \rightarrow \pi^+\pi^-\pi^0)}$