


# Synergies with EHM at JLab

## Talk outline:

- Insight into EHM from combined exploration of meson and baryon structure
- Mapping dressed quark mass function from the data on the ground nucleon structure
- EHM from combined studies of the nucleon elastic form factors and  $\gamma_v p N^*$  electrocouplings
- New opportunities for gaining insight into EHM from inclusive electron scattering data in the resonance region



V.I. Mokeev, Jefferson  
Laboratory, for the CLAS  
Collaboration



Perceiving the Emergence of Hadron Mass through AMBER@CERN  
April 27-29, 2021, Geneva Switzerland

# Studies of the Nucleon and N\* Structure as a Window into EHM

## Composition of the Nucleon Mass:


$M_p$ , MeV (PDG20)

938.2720813  
 $\pm 0.0000058$

Sum of bare quark  
masses, MeV

$2.16 + 2.16 + 4.67$   
 $= 8.99$  or  $< 1.0\%$

proton




$M_n$ , MeV (PDG20)

939.5654133  
 $\pm 0.0000058$

Sum of bare quark  
masses, MeV


$4.67 + 4.67 + 2.16$   
 $= 11.50$  or  $< 1.1\%$

neutron






Dominant part of nucleon mass emerges from strong interaction in the regime when QCD's process-independent running-coupling becomes comparable with unity

- Elastic/resonance electroexcitation amplitudes are sensitive to dressed quark propagator allowing us to map-out momentum dependence of dressed quark mass
- Consistent results on momentum dependence of dressed quark mass from independent studies of elastic and transition  $N \rightarrow N^*$  FFs validate credible insight into the dynamics of hadron mass generation



# Basics for Insight into EHM with Continuum QCD



## Emergence of Dressed Quarks and Gluons D. Binosi et al., Phys. Rev. D 95, 031501 (2017)



- Dressed quark/gluon masses converge at the complete QCD mass scale of 0.43(1) GeV elucidating the trace anomaly  $O(M_{\text{proton}})$
- Momentum-dependent quark/gluon masses shape the ground/excited hadron structure and assure that  $\alpha(k)$  is free of divergences, making QCD a well-defined theory at all distance scales

## QCD Running Coupling $\alpha(k)$ & Dressed Quark/Gluon Masses


C.D. Roberts, Symmetry 12, 1468 (2020)




Inferred from QCD Lagrangian with only the  $\Lambda_{\text{QCD}}$  parameter

# EHM from Global Hadron Structure Analysis


Nucleon Elastic FF




$\gamma_v p N^*$  Electrocouplings




(e,e'X) Inclusive Scattering




Pion Elastic FF



Pion PDF




Dressed Quark/Gluon Running Masses




Continuum QCD approach has demonstrated the capability of gaining insight into the dressed quark/gluon running masses from all of the experimental results above!



# Dressed Quark Mass Function from the Nucleon Elastic Form Factor Data





## Description of N/N\* electroexcitation in DSEQCD




- Elastic form factors (bottom plot) are sensitive to the rate for the transition between a fully dressed (constituent) quark in the infrared to an almost bare QCD quark in ultraviolet seen in the momentum dependence of quark mass function (top plot)

# Charting the Contributions from the Di-Quarks




# EHM from the Ground Nucleon Structure Exploration in 12 GeV Era

- A unique combination of high luminosity ( $10^{38} \text{ cm}^{-2}\text{s}^{-1}$ ), duty cycle, and polarization capabilities make the **SBS facility at JLab** the most suitable in the world for studies of the nucleon elastic form factor at high  $Q^2$  up to  $15 \text{ GeV}^2$
- The **BONUS installation in the CLAS12** detector extends the capabilities in the studies of the  $F_2$  DIS structure function off neutrons at large  $x_B$  and  $Q^2$  up to  $14 \text{ GeV}^2$



Shed light on the presence of di-quark correlations of spin-parity  $0^+$  and  $1^+$

- Provide strong constraints on the rate of the transition from fully dressed to pQCD quarks
- Further explore the relevance of di-quark correlations through the search for zero crossing in  $Q^2$ -evolution of d-quark contribution into Dirac nucleon elastic form factor





# Nucleon Resonance Electrocouplings from Data On Exclusive Meson Electroproduction with CLAS

Exclusive meson electroproduction channels	Excited proton states	$Q^2$ -ranges for extracted $\gamma_v p N^*$ electrocouplings, $\text{GeV}^2$
$\pi^0 p, \pi^+ n$	$\Delta(1232)3/2^+$	0.16-6.0
	$N(1440)1/2^+, N(1520)3/2^-, N(1535)1/2^-$	0.30-4.16
$\pi^+ n$	$N(1675)5/2^-, N(1680)5/2^+, N(1710)1/2^+$	1.6-4.5
$\eta p$	$N(1535)1/2^-$	0.2-2.9
$\pi^+ \pi^- p$	$N(1440)1/2^+, N(1520)3/2^-$	0.25-1.50
	$\Delta(1620)1/2^-, N(1650)1/2^-, N(1680)5/2^+, \Delta(1700)3/2^-, N(1720)3/2^+, N'(1720)3/2^+$	2.0-5.0 (preliminary) 0.5-1.5

- The  $N^*$  electroexcitation amplitudes ( $\gamma_v p N^*$  electrocouplings) in a broad range of  $Q^2$  offer a unique opportunity to explore universality on environmental sensitivity of dressed quark mass function
- Consistent results on dressed quark mass function from  $\gamma_v p N^*$  electrocouplings of different resonances validate insight into EHM in a nearly model-independent way





# Electrocouplings of $N(1440)1/2^+$ from $\pi N$ and $\pi^+\pi^-p$ Electroproduction off Proton Data



Consistent results on  $N(1440)1/2^+$  electrocouplings from independent studies of two major  $\pi N$  and  $\pi^+\pi^-p$  electroproduction channels with different non-resonant contributions allow us to evaluate the systematic uncertainties of these quantities in a nearly model-independent way

## Dyson-Schwinger Equations (DSE):

- J. Segovia et al., Phys. Rev. Lett. 115, 171801 (2015)
- J. Segovia et al., Few Body Syst. 55, 1185 (2014)




*DSE analyses of CLAS data on  $\Delta(1232)3/2^+$  electroexcitation demonstrate that dressed quark mass runs with momentum*

Good data description at  $Q^2 > 2.0 \text{ GeV}^2$  achieved with the same dressed quark mass function for the ground and two excited nucleon states of distinctively different structure **validates the DSE results** on momentum dependence of dressed quark mass.  $\gamma_V p N^*$  electrocoupling data offer access to the strong QCD dynamics underlying hadron mass generation.


**One of the most important achievements in hadron physics of the last decade in synergistic efforts between experimentalists, phenomenologists, and theorists**

# Predictions for Electrocouplings of the First Radial $\Delta(1600)3/2^+$ from Approaches with Momentum Dependent Dressed Quark Mass

$\Delta(1600)3/2^+$



Parameter free continuum QCD (DSE) predictions for  $\Delta(1600)3/2^+$  electrocouplings. Ya Lu et al., Phys. Rev. D100, 034001 (2019)




LFRQM accounting for 3-quark configuration mixing : I.G. Aznauryan and V.D. Burkert arXiv: 1603.06692 [nuc-ph]

# Description of the $\pi^+\pi^-p$ CLAS Data with Electrocouplings of $\Delta(1600)3/2^+$ from Continuum QCD Approach

$\chi^2/\text{d.p.} = 2.71$


$\chi^2/\text{d.p.} = 2.59$




- Reasonable data description and pronounced differences in the resonant/background contributions offer a good prospect for extraction of  $\Delta(1600)3/2^+$  electrocouplings from the  $\pi^+\pi^-p$  electroproduction data
- Confirmation of the continuum QCD expectations on  $\Delta(1600)3/2^+$  electrocouplings will provide strong evidence for credible access to the mass functions of u- and d-quarks at quark momenta  $< 0.5 \text{ GeV}$

# Quality of the $\pi^+\pi^-p$ Data Description with/without $\Delta(1600)3/2^+$


2.0  $\text{GeV}^2 < Q^2 < 2.4 \text{ GeV}^2$




2.4  $\text{GeV}^2 < Q^2 < 3.0 \text{ GeV}^2$



3.0  $\text{GeV}^2 < Q^2 < 3.5 \text{ GeV}^2$



4.2  $\text{GeV}^2 < Q^2 < 5.0 \text{ GeV}^2$



Thick dashed line stands for  $\Delta(1600)3/2^+$  mass, the interval between thin dashed lines corresponds to the resonance width

—  $\Delta(1600)3/2^+$  contribution is replaced by the non-resonant processes

—  $\Delta(1600)3/2^+$  resonance is included with electrocouplings predicted within continuum QCD (slide #11)


Implementation of  $\Delta(1600)3/2^+$  resonance with electrocouplings from the continuum QCD approach improves description of  $\pi^+\pi^-p$  electroproduction data at  $1.45 \text{ GeV} < W < 1.68 \text{ GeV}$  and  $2.0 < Q^2 < 5.0 \text{ GeV}^2$



# N\* Electroexcitation to High Q<sup>2</sup> with CLAS12

**Expected outcome:** The first results on the  $\gamma_p N^*$  electrocouplings of most N\* states from data in the range  $W < 3.0$  GeV and  $Q^2 > 5.0$  GeV<sup>2</sup> for exclusive reaction channels:  $\pi N$ ,  $\pi\pi N$ ,  $KY$ ,  $K^*Y$ ,  $KY^*$

kinematic coverage for RG-A data @ 10.6 GeV



Expected events per Q<sup>2</sup>/W bin for full RG-A dataset

$\pi^+n$			$K^+\Lambda$ & $K^+\Sigma^0$					$\pi^+\pi^+p$		
Q <sup>2</sup> [GeV <sup>2</sup> ]	W [GeV] 1.5-1.55	W [GeV] 1.7-1.75	Q <sup>2</sup> [GeV <sup>2</sup> ]	W $_{\Lambda}$ [GeV] 1.7-1.75	W $_{\Sigma}$ [GeV] 1.7-1.75	W $_{\Lambda}$ [GeV] 1.9-1.95	W $_{\Sigma}$ [GeV] 1.9-1.95	Q <sup>2</sup> [GeV <sup>2</sup> ]	W [GeV] 1.7-1.75	W [GeV] 1.9-1.95
			1.4-2.2	63417	6012	66564	33170			
			2.2-3.0	72144	5364	77443	28720			
5.2-5.8	15272	4175	3.0-4.0	52358	3945	51991	18936	5.2-5.8	2813	2808
5.8-6.5	10737	2637	4.0-5.0	24833	3103	26690	5925	5.8-6.5	1822	1969
6.5-7.2	7367	1684	5.0-6.0	11203	1598	11160	2642	6.5-7.2	1159	1294
7.2-8.1	4567	1290	6.0-7.0	5566	648	6300	943	7.2-8.1	661	924
8.1-9.1	2742	540	7.0-8.0	2606	338	3276	633	8.1-9.1	364	414
9.1-10.5	1453	194	8.0-9.0	1440	244	936	86	9.1-10.5	118	179

Collecting the remainder of the approved RG-A beam time will give a factor of two more statistics

This will extend the Q<sup>2</sup> range of the  $\gamma_p N^*$  electrocouplings to **8-10 GeV<sup>2</sup>** for each of these channels – *the data collected so far will limit us to 6-8 GeV<sup>2</sup>*

# Emergence of Hadron Mass and Quark-Gluon Confinement

**N\* electroexcitation studies at JLab will address the critical open questions:**


*How is >98% of visible mass generated?*

*How does confinement emerge from QCD and how is it related to Dynamical Chiral Symmetry Breaking?*

*What is the behavior of QCD's running coupling at infrared momenta?*


*(S.J, Brodsky et al., Int. J. Mod. Phys. Rev. E29, 2030006 (2020))*

**Mapping-out quark mass function from the CLAS12 results on  $\gamma_v p N^*$  electrocouplings of spin-isospin flip, radial, and orbital excited nucleon resonances at  $5 < Q^2 < 12 \text{ GeV}^2$  will allow us to explore the transition from strong QCD to pQCD regimes**



CLAS results vs. theory expectations with running quark mass

**Access to the dressed quark/hadron mass generation**




# Resonant Contributions into Inclusive $F_2(W, Q^2)$ Structure Functions

Data points are from interpolation of the CLAS results re-evaluated with the  $\sigma_L/\sigma_T$  ratio from Hall C data

CLAS data:  
M. Osipenko et al., PRD 67, 092001 (2003)

Hall C data:  
Y. Liang, PhD thesis of American University (2003)

$N^*$  contributions computed with  $\gamma_p N^*$  electrocouplings from the CLAS data:  
A.N. Hiller Blin et al, Phys. Rev. C100, 035201 (2019)



**The non-resonant parts of  $F_2$  structure function can be computed with the dressed quark mass function supported by the results on pion and nucleon elastic FFs, and on  $\gamma_p N^*$  electrocouplings. From the resonant/non-resonant contributions, full  $F_2$  structure function can be computed and confronted with the data offering a complementary way to validate insight into EHM**




# Evolution of the Resonant Contributions with Photon Virtuality

Resonant contributions into the  $F_2$ ,  $F_L$  structure functions are in the range of 40-60%, suggesting good prospects for the extraction of the  $\gamma_v p N^*$  electrocouplings at  $Q^2 > 4.0 \text{ GeV}^2$ , allowing to map out the dressed quark mass towards higher quark momenta

**Intriguing feature:** the same rate in  $Q^2$ -evolution of the resonant and non-resonant contributions into The  $F_2$  structure function within the second resonance region at  $Q^2 > 2.0 \text{ GeV}^2$

Complementary information from  $F_2$  and  $F_L$



## Conclusions and Outlook

- Mapping dressed quark mass function from the data on the ground and excited nucleon state structure is of particular importance for validation of the insight into EHM.
- The Hall A/C data on nucleon elastic form factors **provided constraints on the transition rate from fully dressed constituent quark in infrared ( $p < 0.5$  GeV) towards almost bare QCD quark in ultraviolet ( $p > 2.0$  GeV).**
- A good description of CLAS results on  $\Delta(1232)3/2^+$  and  $N(1440)1/2^+$  electroexcitation amplitudes **achieved with the same dressed quark mass function as used previously in successful evaluations of the elastic ground nucleon and pion form factors, validate insight to the dynamics that underlie the emergence of hadron mass.** Studies of the  $\Delta(1600)3/2^+$  electrocouplings are in progress.
- The expected results from JLab in the 12 GeV era on the pion, kaon, ground and excited nucleon structure will allow us to map out the dressed quark mass function at the distances where the transition from quark-gluon confinement to pQCD regime is expected, **addressing the most challenging problem of the Standard Model on the nature hadron mass.**
- **Synergy between experiment, phenomenology, and theory is of particular importance in order to achieve these challenging objectives.**





# Back Up



## N\* Structure in Experiments with CLAS/CLAS12

The experimental program on the studies of N\* structure in exclusive meson photo-/electroproduction with CLAS/CLAS12 seeks to determine:

- $\gamma_v p N^*$  electrocouplings at photon virtualities  $Q^2$  up to  $5.0 \text{ GeV}^2$  for most excited proton states through analyzing major meson electroproduction channels from CLAS data
  - extend accessible  $Q^2$  range within  $5.0 \text{ GeV}^2 < Q^2 < 12 \text{ GeV}^2$  and down to  $0.05 \text{ GeV}^2$  from CLAS12 data
  - explore hadron mass emergence by mapping out running quark mass in the transition from almost massless pQCD quarks to fully dressed constituent quarks
- A unique source of information on many facets of strong QCD in generating N\* states with different structural features
  - Allow evaluation of the resonant contributions to inclusive  $F_1$ ,  $F_2$ , and  $F_L$  structure functions from experimental results on  $\gamma_v p N^*$  electrocouplings

### References:

1. I.G. Aznauryan and V.D. Burkert, Prog. Part. Nucl. Phys. 67, 1 (2012)
2. V.D. Burkert and C.D. Roberts, Rev. Mod. Phys. 91, 011003 (2019)
3. D.S. Carman, K. Joo, and V.I. Mokeev, Few Body Syst. 61, 29 (2020)
4. A.N. Hiller Blin et al., Phys. Rev. C100, 035201 (2019)





# Resonant Contributions into Inclusive $F_1(W, Q^2)$ Structure Functions & the Contributions from the PDF in the Ground State of the Nucleon Evaluated from the Data in DIS Region

Resonant contributions:

A.N. Hiller Blin et al.,  
PRC 100, 035201 (2019)



$N(1440) 1/2^+$   
 $N(1520) 3/2^-$   
 $N(1535) 1/2^-$   
 $N(1650) 1/2^-$   
 $N(1675) 5/2^-$   
 $N(1680) 5/2^+$   
 $N(1710) 1/2^+$   
 $N(1720) 3/2^+$   
 $\Delta(1232) 3/2^+$   
 $\Delta(1620) 1/2^-$   
 $\Delta(1700) 3/2^-$   
 $N^*(1720) 3/2^+$   
 Total  
 Total coherent  
 CLAS Data  
 JAM  
 JAM TMC Moffat  
 JAM TMC Brady approx.  
 JAM TMC Brady



Data points are from  
interpolation of the  
CLAS results re-evaluated  
with the  $\sigma_L/\sigma_T$  ratio  
from Hall C data

CLAS data:  
M. Osipenko et al., PRD  
67, 092001 (2003)

Hall C data:  
Y. Liang, PhD thesis of  
American University  
(2003)



Green dot-dashed lines:  $F_1$  from JAM PDF

Other smooth curves:  $F_1$  from JAM PDF after target mass corrections within different prescriptions



## N\* studies at $0.05 \text{ GeV}^2 < Q^2 < 7.0 \text{ GeV}^2$ with CLAS12

<b>Hybrid Baryons</b> E12-16-010	Search for hybrid baryons (qqqq) focusing on $0.05 \text{ GeV}^2 < Q^2 < 2.0 \text{ GeV}^2$ in mass range from 1.8 to 3 GeV in $K\Lambda$ , $N\pi\pi$ , $N\pi$ ( <i>A. D'Angelo, et al.</i> )
<b>KY</b> <b>Electroproduction</b> E12-16-010A	Study N* structure for states that couple to KY through measurements of cross sections and polarization observables that will yield $Q^2$ evolution of electrocoupling amplitudes at $Q^2 < 7.0 \text{ GeV}^2$ ( <i>D. Carman, et al.</i> )

**Approved by PAC44**

Run Group conditions:

$E_b = 6.6 \text{ GeV}$ , 50 days


$E_b = 8.8 \text{ GeV}$ , 50 days

- Polarized electrons, unpolarized  $\text{LH}_2$  target
- $L = 1 \times 10^{35} \text{ cm}^{-2}\text{s}^{-1}$

# Hunting for Glue in Excited Baryons with CLAS12

Can glue be a structural component to generate hybrid  $q^3g$  baryon states?

Predictions of the  $N^*$  spectrum from QCD show both regular  $q^3$  and hybrid  $q^3g$  states



JLab LQCD group results

Search for hybrid baryons with CLAS12 in exclusive  $KY$  and  $\pi^+\pi^-p$  electroproduction

LQCD and/or QM predictions on  $Q^2$  evolution of the hybrid-baryon electroexcitation amplitudes are critical in order to establish the nature of a baryon state

