

J/ ψ near threshold photoproduction at CLAS12

RICHARD TYSON




University
of Glasgow

J/ψ Near Threshold Photoproduction

- ▶ CLAS12 operates close to the 8.2 GeV J/ψ photoproduction threshold.
- ▶ Near threshold, all the valence quarks of the nucleon are predicted to participate in J/ψ photoproduction while at higher energies it is predicted that one or two hard gluons can be involved [2]. This is studied by measuring the total cross section as a function of beam energy.
- ▶ [3] predicts that the t dependency of the differential cross section is defined by the nucleon gluonic form-factor, for which a dipole form is assumed with $m_g^2 \approx 1 \text{ GeV}^2$ as:

$$F(t) \propto (1 - t/m_g^2)^{-2}$$
- ▶ CLAS12 will also make a first measurement of J/ψ photoproduction on the neutron.




Measurements of the J/ψ total cross section as a function of the photon beam energy and theoretical predictions scaled to GlueX data [1].

- [1] A. Ali, et. al. (GlueX Collaboration), *Phys. Rev. Lett.* **123**, 072001 (2019).
 [2] S. Brodsky, E. Chudakov, P. Hoyer, J. Laget, *Phys. Lett. B.* **498**, 23 (2001).
 [3] L. Frankfurt, M. Strikman, *Phys. Rev. D.* **66**, 031502 (2002)

P_c^+ resonances with CLAS12

- ▶ Different theoretical models for the structure of the P_c^+ pentaquarks favor different decay mechanisms.
- ▶ CLAS12 should be able to place upper limits on the branching fraction $B(P_c^+ \rightarrow J/\psi p)$.
- ▶ J/ψ photoproduction on the neutron further offers the possibility of looking for the isospin partners of the P_c^+ Pentaquarks.



The $J/\psi p$ invariant mass distribution measured at the LHCb. Taken from:

R. Aaij, et. al. (LHCb Collaboration), *Phys. Rev. Lett.* **122**, 22 (2019).



The CLAS12 Detector

- ▶ The CLAS12 Detector is located in Jefferson Lab's Hall B, in Newport News, Virginia.
- ▶ The GlueX detector is located in Hall D.




The CLAS12 Detector

- ▶ The CLAS12 Detector is located in Jefferson Lab's Hall B, in Newport News, Virginia.
- ▶ The recently upgraded CEBAF accelerator facility produces a 12 GeV electron beam, with beam energies up to 11 GeV delivered to Hall B.
- ▶ The Forward Detector has polar angle coverage of 5 to 35 degrees.
- ▶ The Central Detector has polar angle coverage of 35 to 125 degrees.



CLAS12 Forward Detector

- ▶ All final state particles are detected with the Forward Detector.
- ▶ The High Threshold Cherenkov Counter (HTCC) was built to identify electrons as other particle types generally won't fire the HTCC.
- ▶ The tracking system and Drift Chambers (DC) measure the charge and momentum of particles.
- ▶ The Forward Time Of Flight (FTOF) counters were designed to resolve pions, kaons, protons and deuterons.
- ▶ The Electromagnetic Calorimeters (PCAL and EC) are used to detect neutrals and identify electrons as they should deposit more energy than other particle types.



Experiment Overview


- ▶ J/ψ decays to a lepton pair, with l^+l^- denoting either e^+e^- or $\mu^+\mu^-$.
- ▶ CLAS12 took data with both a proton and a deuterium target offering several potential final states:

$$ep \rightarrow (e')l^+l^-p$$


$$ed \rightarrow (e')l^+l^-d$$

$$e p_{\text{bound}} \rightarrow (e')l^+l^-p$$

$$e n_{\text{bound}} \rightarrow (e')l^+l^-n$$





J/ψ quasi-real photoproduction on a proton target



Feynmann diagram of P_c^+ pentaquark photoproduction with a proton target.



Initial Event Selection

- ▶ To select only quasi-real photoproduction events regime we can minimize:
 - ▶ The difference between the initial and scattered electron momentum, Q^2
 - ▶ The scattered electron transverse momentum fractions in the x and y components, $|\frac{p_x}{p}|$ and $|\frac{p_y}{p}|$.
- ▶ Similarly, we want the missing mass close to the mass of the scattered electron (which is effectively 0).
- ▶ The widths of these distributions can be parametrised as a function of the photon energy.





Initial Particle Identification

- ▶ Electrons and positrons are required to produce a signal in the HTCC and high energy deposition in the calorimeter. Their main source of background is due to high momentum pions firing the HTCC.
- ▶ Muon candidates are minimum ionizing particles, and susceptible to a significant charged pion contamination.
- ▶ For protons (and charged hadrons) a cut is made on the Beta versus Momentum parametrization.
- ▶ For neutrons, the initial requirement is simply $\text{Beta} < 0.9$. Their main source of background comes from photons reconstructed with low Beta.





Particle ID Refinement

- ▶ We trained two classifiers to distinguish between pions and either muons or positrons. Simulated CLAS12 data was used to create the training samples.
- ▶ The classifier output is given as a probability of being a signal event, called the response, and effectively reduces the PID process down to a cut on the response.
- ▶ This cut can be varied to study the systematic effect introduced by the classifiers.
- ▶ This was implemented using the ROOT TMVA package: A. Hoecker, *et. al.*, user guide available online at: [arXiv:physics/0703039v5](https://arxiv.org/abs/physics/0703039v5)



Di-lepton Invariant Mass

- ▶ Plotted here are the invariant mass distributions of:
 - ▶ $\mu^+\mu^-$ produced on a proton target
 - ▶ e^+e^- produced on a bound proton in the deuteron target.
- ▶ Radiative effects shift the e^+e^- J/ ψ mass peak away from the J/ ψ mass (3.097 GeV) as the reconstructed momentum is post-radiation.
- ▶ These are preliminary and produced with only a subset of all available data.



Conclusion




- ▶ The total and differential J/ψ photoproduction cross section provide unique insight about the nucleon gluonic form factor and the J/ψ production mechanism.
- ▶ CLAS12 will make a first measurement of the J/ψ photoproduction cross section ratio on proton and neutron, and a possible comment on the isospin partners of the P_c^+ Pentaquarks.
- ▶ Several CLAS12 analyses aiming for these measurements are ongoing:
 - ▶ J/ψ near threshold photoproduction in $ep \rightarrow (e')e^+e^-p$, Joseph Newton
 - ▶ Timelike Compton Scattering in $ep \rightarrow (e')e^+e^-p$, Pierre Chatagnon
 - ▶ J/ψ near threshold photoproduction in $ep \rightarrow (e')\mu^+\mu^-p$
 - ▶ J/ψ near threshold photoproduction in $ep_{bound} \rightarrow (e')e^+e^-p$
 - ▶ J/ψ near threshold photoproduction in $en_{bound} \rightarrow (e')e^+e^-n$
- ▶ Next: total and differential cross sections for the proton and deuterium targets.



Backup Slides


P_c^+ Models

- ▶ Hadronic molecules: Weakly coupled charmed baryon and charmed meson.
- ▶ Hadro-charmonium states: compact bound $c\bar{c}$ state and light quarks.
- ▶ Quarks in a bag: Two tightly correlated di-quarks and an anti-quark.



ρ , ω and ϕ mesons

- ▶ Plotted here is the invariant mass of e^+e^- produced on a bound proton in the deuteron target.
- ▶ ϕ mesons are clearly resolved.
- ▶ ρ and ω mesons are unresolvable but clearly present.



Fiducial Cuts and Momentum Corrections

- ▶ If an electron or positron hits close to the edges of the PCAL, the shower may not be fully contained within the calorimeter volume.
- ▶ This can lead to a wrong sampling fraction and reduced identification power for electrons and positrons.
- ▶ Electrons and positrons radiate photons before reaching the forward detector. The reconstructed electron momentum is therefore the post-radiation momentum.
- ▶ This is corrected by adding the momentum of the radiated photon, identified by a small angular difference with the electron.

