Insight into Emergence of Hadron Mass in the Exploration of N* Structure at JLab after Energy and Luminosity Increase





FFA concept for CEBAF energy upgrade:

- 5 additional recirculations in existing tunnel
- Use existing LINACs 5 more times each





Tellerson Pab

- Concept on emergence of hadron mass (EHM) from QCD within continuum Schwinger function method (CSM)
- Gaining insight into EHM from <u>combined</u> studies of π/K and N/N* structure. Why is this needed?
- Evidence and limitations for gaining insight into EHM from experiments at JLab with 6/12 GeV beam
- Understanding EHM from the results on N* electrocouplings determined within Q²-range up to 35 GeV²





Hadron Physics Opportunities with JLab Energy and Luminosity Upgrade



How do the Ground/Excited State Nucleon Masses Emerge?





- Higgs mechanism generates the masses of bare quarks
- Dominant part of nucleon mass is generated in processes other than the Higgs mechanism

The Continuum Schwinger method (CSM) has conclusively demonstrated that the dominant part of hadron mass is generated by the strong interaction in the regime where the QCD running coupling becomes comparable with unity the so-called strong QCD regime



Basics for Insight into EHM: CSM and Lattice QCD Synergy



In the regime of the QCD running coupling comparable with unity, dressed quarks and gluons with distance (momentum) dependent masses emerge from QCD, as follows from the equations of the motion for the QCD fields depicted above.



Basics for Insight into EHM: Continuum and Lattice QCD Synergy

- Express the fundamental feature: emergence of the quark and gluon masses even in the case of massless quarks in the chiral limit and massless QCD gluons
- Continuum QCD results are confirmed by LQCD
- Insight into EHM from data on hadron structure represents a challenge for experimental hadron physics



Dressed Quark/Gluon Masses (Continuum QCD) C.D. Roberts, Symmetry 12, 1468 (2020)



Inferred from QCD Lagrangian with only Λ_{QCD} parameter



V.I. Mokeev, APCTP Workshop on Nuclear Physics 2022, July 11-16, 2022, Jeju, Korea



Mass Budgets

- Studies of π/K structure elucidate the interference between emergent and Higgs mechanisms in EHM
- Studies of ground/excited state nucleon structure allow us to explore the dressed quark mass function in a different environment where the sum of dressed quark masses is the dominant contribution to the physical masses of these states, offering insight into emergent mechanisms

• Successful description of the π/K elastic FF and PDF, nucleon elastic/axial FFs, and the $\gamma_v pN^*$ electrocouplings of prominent nucleon resonances of different structure achieved with the *same* dressed quark mass function is of particular importance for the validation of insight into EHM.



N* Photo-/Electroexcitation Amplitudes and their Extraction from Exclusive Data



• Consistent results on the $\gamma_{r,v}$ pN* photo-/electrocouplings from different meson photo-/electroproduction channels allow us to validate reliable extraction of these quantities.



Nucleon Resonance Electrocouplings from Data On Exclusive Meson Electroproduction with CLAS

Exclusive meson electroproduction channels	Excited proton states	Q ² -ranges for extracted γ _v pN* electrocouplings, GeV ²
π ⁰ p, π ⁺ n	∆(1232)3/2⁺	0.16-6.0
	N(1440)1/2⁺,N(1520)3/2⁻, N(1535)1/2⁻	0.30-4.16
π ⁺ n	N(1675)5/2 ⁻ , N(1680)5/2 ⁺ N(1710)1/2 ⁺	1.6-4.5
ηρ	N(1535)1/2 ⁻	0.2-2.9
π ⁺ π ⁻ p	N(1440)1/2 ⁺ , N(1520)3/2 ⁻ ∆(1620)1/2 ⁻ , N(1650)1/2 ⁻ , N(1680)5/2 ⁺ , ∆(1700)3/2 ⁻ , N(1720)3/2 ⁺ , N'(1720)3/2 ⁺	0.25-1.50 2.0-5.0 (preliminary) 0.5-1.5

- The N* electroexcitation amplitudes (γ_vpN* electrocouplings) have become available in a broad range of Q² < 5 GeV²
- In the mass range W < 1.6 GeV, the $\gamma_v pN^*$ electrocoupling were obtained from independent studies of πN , ηp , and $\pi^* \pi^- p$ electroproduction

Most recent results can be found in: A.N. Hiller Blin et al, PRC100, 035201 (2019)



Electrocouplings of N(1440)1/2⁺ and N(1520)3/2⁻ Resonances from π N and $\pi^+\pi^-p$ Electroproduction off Proton Data



Consistent results on the N(1440)1/2⁺ and N(1520)3/2⁻ electrocouplings from independent studies of the two major π N and $\pi^+\pi^-p$ electroproduction channels with different non-resonant contributions allow us to evaluate the systematic uncertainties of these quantities in a nearly model-independent way.



Emergence of Hadron Mass: Concept from CSM vs. Available Experimental Results

Successful description of the pion and nucleon elastic FFs, and the electrocouplings of the $\Delta(1232)3/2^+$ and N(1440)1/2⁺ resonances has been achieved <u>with the same dressed</u> <u>quark/gluon mass functions</u>



Dressed Quark/Gluon Masses from CSM C.D. Roberts, Symmetry 12, 1468 (2020)



- Dressed quarks with dynamically generated masses represent active degrees of freedom in the structure of the pion, nucleon, and the Δ(1232)3/2⁺, N(1440)1/2⁺ resonances
- Strong evidence for insight into EHM



Electrocouplings of the ∆(1600)3/2⁺: CSM Prediction vs. Data Determination



Parameter-free CSM predictions for ∆(1600)3/2⁺ electrocouplings Ya Lu et al., Phys. Rev. D 100, 034001 (2019)

efferson V



V.I. Mokeev, APCTP Workshop on Nuclear Physics 2022, July 11-16, 2022, Jeju, Korea

Electrocouplings of the ∆(1600)3/2⁺: CSM prediction vs. Data Determination



--- CSM predictions, Ya Lu et al., Phys. Rev. D 100, 034001 (2019)

Electrocouplings from independent analyses of $\pi^+\pi^-p$ differential cross sections within three W-intervals, 1.46<W<1.56 GeV, 1.51<W<1.61 GeV, and 1.56<W<1.66 GeV for 2.0<Q²<5.0 GeV²

CLAS results on $\Delta(1600)3/2^+$ electrocouplings confirmed the CSM prediction, solidifying evidence for gaining insight into dressed quark mass function and, consequently, into EHM from the studies of $\gamma_v pN^*$ electrocouplings



N* Electroexcitation to High Q² with CLAS12

Expected outcome: The first results on the $\gamma_v pN^*$ electrocouplings of most N* states from data in the range W < 2.5 GeV and Q² > 5.0 GeV² for exclusive reaction channels: πN , $\pi \pi N$, KY, KYY, KY*



Expected events per Q²/W bin for full RG-A dataset

	π*n		K+Λ & K+Σ ⁰				<i>π</i> ⁺ <i>π</i> ⁻p			
Q² [GeV²]	W [GeV] 1.5-1.55	W [GeV] 1.7-1.75	Q² [GeV²]	₩ _^ [GeV] 1.7-1.75	W _Σ [GeV] 1.7-1.75	₩ _^ [GeV] 1.9-1.95	W _Σ [GeV] 1.9-1.95	Q² [GeV²]	W [GeV] 1.7-1.75	W [GeV] 1.9-1.95
			1.4-2.2	63417	6012	66564	33170			
			2.2-3.0	72144	5364	77443	28720			
5.2 - 5.8	15272	4175	3.0-4.0	52358	3945	51991	18936	5.2-5.8	2813	2808
5.8 - 6.5	10737	2637	4.0-5.0	24833	3103	26690	5925	5.8-6.5	1822	1969
6.5 - 7.2	7367	1684	5.0-6.0	11203	1598	11160	2642	6.5-7.2	1159	1294
7.2-8.1	4567	1290	6.0-7.0	5566	648	6300	943	7.2-8.1	661	924
8.1-9.1	2742	540	7.0-8.0	2606	338	3276	633	8.1-9.1	364	414
9.1-10.5	1453	194	8.0-9.0	1440	244	936	86	9.1-10.5	118	179

Collecting the remainder of the approved RG-A beam time will give a factor of two more statistics

This will extend the Q² range of the $\gamma_v pN^*$ electrocouplings to 8-10 GeV² for each of these channels – the data collected so far will limit us to 6-8 GeV²



Emergence of Hadron Mass from N* Studies with CLAS/CLAS12 and Expected with CLAS24

N* electroexcitation studies at JLab during 12 GeV era will address the critical questions:

How is >98% of visible mass generated?

How EHM is related to Dynamical Chiral Symmetry Breaking?

(S.J, Brodsky et al., Int. J. Mod. Phys. Rev. E29, 2030006 (2020))

Mapping-out the dressed quark mass function from $\gamma_v pN^*$ electrocouplings of different spin-isospin flip, radial, and orbital excited nucleon states at 5<Q²<12 GeV² will increase knowledge on EHM and motivate efforts to determine $\gamma_v pN^*$ electrocouplings for Q² up to 35 GeV² to explore the full range of distances (quark momenta) where the dominant part of hadron mass is expected to be generated

				0.4		Continuum Met	Schwing hod	er
	Q^2 -coverage for $\gamma_v p N^*$ electrocoupling results	Accessible range of quark momenta p	Fraction of fully dressed quark mass generated at p <p<sub>max</p<sub>			essed auarks	$p = \sqrt{Q^2}/$	3
CLAS	<5.0 GeV ²	<0.8 GeV	15-20 %		C. C. S. C. S. S. C. S.		Dressed	gluons
CLAS12	<9.0 GeV ²	<1.0 GeV	40-50 %		CLAS LAS12	524		
CLAS24	<35.0 GeV ²	<2.0 GeV	>80 %				***********	
				0	1	2	3	4
	Momentum, p (GeV)							

Energy and luminosity increase are needed in order to obtain information on the $\gamma_v pN^*$ electrocouplings at Q²>10 GeV², allowing us to map out the momentum dependence of the dressed quark mass within the entire range of distances where the dominant part of hadron mass is generated



Both EicC and EIC would need much higher, unlikely feasible luminosity

The exclusive electroproduction measurements foreseen at JLab after completion of the 12 GeV program:

- Beam energy at fixed target: 24 GeV
- Nearly 4π coverage
- High luminosity

Offer maximal achievable luminosity for extraction of $\gamma_v p N^*$ electrocouplings at Q²>10 GeV²



Studies of $\gamma_v p N^*$ Electrocouplings at $Q^2 > 10 \text{ GeV}^2$

• Kinematic coverage in the resonance region for W<2.0 GeV with electron beam energy $E_b=24$ GeV, $\theta_{e max}=35^{\circ}$.

$\Omega^2 = \frac{2M_N E_b - W^2 + M_N^2}{M_N^2}$	W, GeV	1.1	2.0
$Q_{max} = \frac{M_N}{(1 + \frac{M_N}{2E - \sin^2(\theta_{e})})}$	Q ² _{max} [,] GeV ²	36.7	34.4
$2E_b sin^2(\frac{1}{2})$	P quark, GeV	2.02	1.96

- The coverage over Q² within the resonance region allows us to map out the momentum dependence of the dressed quark mass function within the range of p_{quark} < 2.0 GeV and observe how the dominant part of hadron mass emerges.
- Luminosity ~10³⁶ cm⁻²s⁻¹ may be sufficient for the extraction of the γ_vpN* electrocouplings if the rate of exclusive cross section fall-off with Q² at Q²> 10 GeV² is the same as at Q²<10 GeV² and the (resonant)/(background) ratio does not deteriorate.



Hadron Structure Studies with CLAS24



Hadron Structure Group in Hall B developing physics case to support CLAS24 upgrade

List of Participating Institutions:

- Jefferson Lab (Hall B and Theory Division)
- University of Connecticut
- Genova University and INFN of Genova
- Lamar University
- Ohio University
- Skobeltsyn Nuclear Physics Institute and Physics Department at Lomonosov Moscow State University
- University of South Carolina
- INFN Sez di Roma Tor Vergata and Universita di Roma Tor Vergata
- Nanjing University and affiliated institutes
- Tubingen University
- Tomsk State University and Tomsk Polytechnic University
- James Madison University
- George Washington University

https://userweb.jlab.org/~carman/clas24



Luminosity to Determine $\gamma_v pN^*$ Electrocouplings at Q²>10 GeV² from N π Electroproduction



 Luminosities needed for extraction of γ_vpN* electrocouplings from Nπ electroproduction at Q²>10 GeV² were evaluated in each bin of (W,Q²) as:

$$L(W,Q^2)$$
, = 10³⁴ cm⁻²sec⁻¹ Y(W,Q² current)/Y(W,Q² = 5.0 GeV²) (1),

assuming that statistics comparable with those achieved in the measurements with CLAS in the bin of $(W,Q^2 = 5 \text{ GeV}^2)$ at luminosity $10^{34} \text{ cm}^{-2} \text{sec}^{-1}$ will be sufficient

• The ratio $Y(W,Q^2_{current})/Y(W,Q^2=5.0 \text{ GeV}^2)$ was obtained in MC simulation for $E_{beam}=24$ GeV with N π cross sections computed from the MAID07 multipoles at Q²=5.0 GeV², extrapolated into the range of Q²>10 GeV² as the accepted event ratio computed for CLAS12.

 $\gamma_v pN^*$ electrocouplings can be determined up to Q^2_{max} in the range from 17 GeV² to 22 GeV² where the required luminosity remains below ~10³⁶ cm⁻²sec⁻¹

$\gamma_v pN^*$ Electrocouplings at Q²>10 GeV² from K Λ Channel

D.S. Carman, Jefferson Lab



Yields of K Λ events in the bins of (W,Q²,cos θ_{K}^{cm}) were evaluated in MC simulation by employing 2-fold differential cross sections from the CLAS measurements at Q²<4 GeV² and extrapolated into the range of Q²>10 GeV² as 1/Q⁴ (left panel), 1/Q² (right panel)

 $\gamma_v pN^*$ electrocouplings can be determined up to Q^2_{max} in the range from 17 GeV² < Q^2 < 22 GeV², where the projected yields remains comparable with those achieved in the CLAS measurements



Beam Time Needs for Exclusive $p\pi^+\pi^-$

K. Neupane, R.W. Gothe - USC

Based on RG-A fall 2018 Luminosity of 5.96 10³⁴ cm⁻² s⁻¹ at 45 nA



Implementing all analysis cuts (3/2), Golden Run Selection (3), PAC Days (2) 6 (12) years at 5.96 10³⁴ cm⁻² s⁻¹ or 4 (8) month at 10³⁶ cm⁻² s⁻¹

Conclusions and Outlook

- CSM paradigm for EHM makes a broad array of predictions. Those for N* structure in terms of the Q²-evolution of the γ_vpN* electrocoupling are worth testing by confronting the CSM predictions obtained with the same dressed quark mass function against the experimental results determined from exclusive meson electroproduction data.
- A good description of the ∆(1232)3/2⁺ and N(1440)1/2⁺ electroexcitation amplitudes <u>achieved within</u> <u>CSM approach starting from the QCD Lagrangian with the same dressed quark mass function</u> as used in the successful evaluations of the ground state nucleon electromagnetic/axial and pion form factors, and the pion PDF, demonstrated the capability for gaining insight into EHM.
- The CSM parameter-free predictions on the Δ(1600)3/2⁺ electrocouplings have been confirmed by the first and still preliminary results on the electrocouplings of this state obtained from π⁺π⁻p electroproduction data solidifying evidence for gaining insight into the dressed quark mass function and, consequently, into EHM from the studies of γ_vpN* electrocouplings.
- Increase of the CEBAF beam energy up to 24 GeV and the upgrade of the detector capabilities to measure electroproduction events within ~4π acceptance at ~10³⁶cm⁻²sec⁻¹ will make it possible to determine the electrocouplings of prominent N* states at Q² from 10 GeV² to 20 GeV². Analyses of the results on Q²-evolution of γ_vpN* electrocouplings at Q²<20 GeV² will allow us to explore nearly entire range of distances where the dominant part of hadron mass is expected to be generated in the transition from the sQCD to pQCD regimes.
- Confirmation of the CSM predictions on the Q²-evolution of the γ_vpN* electrocouplings of nucleon resonances of different structure obtained with the same dressed quark mass function by the experimental results within the range of Q² up to 20 GeV² will provide the sound evidence for <u>understanding how the dominant part of hadron mass and the N* structructure emerge from QCD</u> and will make JLab@24 GeV the unique and ultimate QCD-facility at the luminosity frontier.







Basics for Insight into EHM: Continuum and Lattice QCD Synergy

- Dressed quark/gluon masses converge at the complete QCD mass scale of 0.43(1) GeV - value impacted by Higgs mechanism
- Express the fundamental feature: emergence of the quark and gluon masses even in the case of massless quarks in the chiral limit and massless QCD gluons
- Continuum QCD results get support from LQCD
- Insight into dressed quark mass function from data on hadron structure represents a challenge for experimental hadron physics

Dressed Quark/Gluon Masses (continuum QCD) C.D. Roberts, Symmetry 12, 1468 (2020)



Inferred from QCD Lagrangian with only Λ_{QCD} parameter





V.I. Mokeev, APCTP Workshop on Nuclear Physics 2022, July 11-16, 2022, Jeju, Korea



Mass Budgets

 Studies of the ground and excited state nucleon structure allow us to explore the dressed quark mass function in a different environment where the sum of dressed quark masses is the dominant contribution into the physical masses of the ground and excited states of the nucleon

 Consistent results on the momentum dependence of the dressed quark mass function from independent studies of the pseudo-scalar mesons and the ground and excited state nucleon structure are of particular importance for the validation of insight into EHM.



Insight to EHM From Resonance Electrocouplings



Good data description at Q²>2.0 GeV² achieved with <u>the same dressed quark mass function</u> for the ground pion/nucleon and two excited nucleon states of distinctively different structure validates the continuum QCD results on the momentum dependence of the dressed quark mass. $\gamma_v pN^*$ electrocoupling data shed light on the strong QCD dynamics underlying hadron mass generation.

One of the most important achievements in hadron physics of the last decade in synergistic efforts between experimentalists, phenomenologists, and theorists.



Studies of $\gamma_v p N^*$ Electrocouplings at $Q^2 > 10 \text{ GeV}^2$

Energy and luminosity increase up to >10³⁶ cm⁻²s⁻¹ are needed in order to obtain information on the $\gamma_v pN^*$ electrocouplings at Q²>10 GeV², allowing us to map out the momentum dependence of the dressed quark mass within the entire range of distances where the dominant part of hadron mass is generated



Both EicC and EIC would need much higher, unlikely feasible luminosity

The exclusive electroproduction measurements foreseen at JLab after completion of the 12 GeV program:

- Beam energy at fixed target: 24 GeV
- Nearly 4π coverage
- High luminosity

Offer maximal achievable luminosity for extraction of $\gamma_v pN^*$ electrocouplings at Q²>10 GeV²



Conclusions and Outlook

 Resolving the most significant open problem in the Standard model on the EHM will make JLab@24 GeV the unique and ultimate QCD-facility at the luminosity frontier with significant contribution into achieving this objective from the studies of the structure of the excited state of the nucleon from the data of experiments with electromagnetic probes (see details in Contribution of the Hadron Structure Group to the Physics Motivation to Increase the Energy and Luminosity of JLab, <u>https://userweb.ilab.org/~carman/clas24/CLAS24-NstarStructure.pdf</u>)

