

J/ ψ Near-Threshold Photoproduction off the Proton and Neutron with CLAS12


RICHARD TYSON



University
of Glasgow

J/ψ Near Threshold Photoproduction

- ▶ CLAS12 operates close to the 8.2 GeV J/ψ ($c\bar{c}$ meson) photoproduction threshold.
- ▶ Near threshold, the 3 gluon exchange's contribution to the cross section is expected to dominate that of the 2 gluon exchange [2].
- ▶ [3] relates the nucleon gluonic form-factor to the t dependency of the differential cross section.
- ▶ CLAS12 will make a first measurement of J/ψ near-threshold photoproduction on the neutron.




Measurements of the J/ψ total cross section as a function of the photon beam energy and theoretical predictions scaled to GlueX data [1].

- [1] A. Ali, et. al. (GlueX Collaboration), *Phys. Rev. Lett.* **123**, 072001 (2019).
[2] S. Brodsky, E. Chudakov, P. Hoyer, J. Laget, *Phys. Lett. B.* **498**, 23 (2001).
[3] L. Frankfurt, M. Strikman, *Phys. Rev. D.* **66**, 031502 (2002).

Mass Radius of the Nucleon

- ▶ The J/ψ differential cross section as a function of t can be related to the mass radius of the nucleon [4].
- ▶ Measurements on the proton at Jlab's Hall-C are in good agreement with those from GlueX [5].
- ▶ The proton mass radius is significantly different from the charge radius ($R_c = 0.8409 \pm 4 \cdot 10^{-4}$ fm), suggesting that its gluon radius is significantly smaller than its quark radius [4].



J/ψ differential cross section as a function of $-t$.
Data from the GlueX Collaboration [1], plot taken from [4].

[4] D.E. Kharzeev, *Phys. Rev. D* **104**, 054015 (2021).

[5] B. Duran, et. al. (J/ψ - 007 Collaboration), preprint available at [arXiv:2207.05212](https://arxiv.org/abs/2207.05212) (2022).



CEBAF

- ▶ The Continuous Electron Beam Accelerator Facility (CEBAF) is located in Newport News, Virginia.
- ▶ The CEBAF Large Acceptance Spectrometer (CLAS12) is located in Hall B.
- ▶ The GlueX detector is located in Hall D, with the $J/\psi - 007$ Collaboration located in Hall C.




The CLAS12 Detector

- ▶ The recently upgraded CEBAF accelerator facility produces a 12 GeV electron beam, with beam energies up to 11 GeV delivered to Hall B.
- ▶ The Forward Detector has polar angle coverage of 5 to 35 degrees.
- ▶ The Central Detector has polar angle coverage of 35 to 125 degrees.



CLAS12 Forward Detector

- ▶ The High Threshold Cherenkov Counter (HTCC) was built to identify electrons.
- ▶ The Drift Chambers (DC) measure the charge and momentum of particles.
- ▶ The Forward Time Of Flight (FTOF) counters were designed to identify charged hadrons.
- ▶ The Electromagnetic Calorimeters (PCAL and EC) are used to detect neutrals and identify electrons and muons.



Experiment Overview


- ▶ J/ψ decays to a lepton pair, with l^+l^- denoting either e^+e^- or $\mu^+\mu^-$.
- ▶ CLAS12 took data with both a proton and a deuterium target offering several potential final states:

$$ep \rightarrow e' J/\psi p \rightarrow (e') l^+ l^- p$$

$$e p_{\text{bound}} \rightarrow e' J/\psi p \rightarrow (e') l^+ l^- p$$

$$e n_{\text{bound}} \rightarrow e' J/\psi n \rightarrow (e') l^+ l^- n$$


$$ed \rightarrow e' J/\psi d \rightarrow (e') l^+ l^- d$$




J/ψ quasi-real photoproduction
on a proton target

Lepton Identification

- ▶ Electrons and positrons are required to produce a signal in the HTCC and have a ratio of the energy deposition to momentum around 0.25.
- ▶ Muons are minimum ionising particles which we select with cuts on their energy deposition in the calorimeters.
- ▶ We refine leptons PID by training a machine learning classifier on variables from several CLAS12 detector subsystems such as:
 - ▶ Energy deposition and cluster information in the calorimeters.
 - ▶ Number of photoelectrons produced in the HTCC.
- ▶ The PID process is then reduced to a cut on the response of the classifier.




J/ ψ and Background Yields vs Response Cut Value




Hadron Identification

- ▶ For protons (and charged hadrons in general) a cut is made on the Beta versus Momentum parametrization.
- ▶ For neutrons we require a neutral charge. No further cuts were applied as there isn't any strong evidence of photon contamination.




Particle Corrections

- ▶ Radiative corrections for **electrons/positrons** add the momentum of **radiated photons**.
- ▶ **Neutrons** also produce **secondary clusters**. These are removed by taking the earliest neutral in a given sector.
- ▶ The reconstructed path length for neutrons is corrected for a more accurate calculation of the momentum.
- ▶ We apply fiducial cuts to remove e^+/e^- hits close to the edges of the PCAL where the shower is not fully contained within the calorimeter.
- ▶ Fiducial cuts in the drift chambers are applied to electrons, positrons, protons and muons by removing hits at the edge of the layers.





e- Sampling Fraction vs LV



Event Selection

- ▶ We remove high Q^2 events to select only quasi-real photoproduction events.
- ▶ We also want the missing mass close to the mass of the scattered electron.
- ▶ Alternatively, we can train a neural network to distinguish between:
 - ▶ J/ψ simulation.
 - ▶ Mismatched particles from different events of the above.
- ▶ The neural network is trained on the four momentum of the final state particles and quantities like the missing mass.




J/ψ Total Cross Section

- Shown here is the total cross section measured in:

- $en_{bound} \rightarrow (e')e^+e^-n$
- $ep_{bound} \rightarrow (e')e^+e^-p$
- $ep \rightarrow (e')\mu^+\mu^-p$

- We are still working on the absolute normalization.
- The differential cross section measurements are also consistent between channels.



Conclusion

- ▶ The near-threshold J/ψ photoproduction cross sections are related to its production mechanism and the nucleon gluonic form factor.
- ▶ Near-threshold J/ψ photoproduction can be directly connected to the nucleon mass radius.
- ▶ We're aiming for a first measurement directly comparing the J/ψ photoproduction cross section on proton and neutron.
- ▶ The analysis of data at CLAS12 is ongoing and well advanced.

Outlook



- ▶ AI based improvements in the tracking reconstruction at CLAS12 show an average 30% increase in the efficiency for 3 charged particles. The data already taken at CLAS12 is about to be reprocessed with the new tracking improvements.
- ▶ The experiment aiming for the measurements of J/ψ photoproduction on deuterium still has roughly ~40% left to run.
- ▶ Future luminosity upgrades at Jlab and CLAS12 will enable high statistics measurements of J/ψ photoproduction. An energy upgrade would also provide complementary studies of higher mass charmonium states.



Backup Slides

P_c^+ resonances with CLAS12

- CLAS12 should be able to place upper limits on the branching fraction $B(P_c^+ \rightarrow J/\psi p)$ and $B(P_c^+ \rightarrow J/\psi n)$.






The $J/\psi p$ invariant mass distribution measured at the LHCb. Taken from:

R. Aaij, et. al. (LHCb Collaboration), *Phys. Rev. Lett.* **122**, 22 (2019).


P_c^+ Models

- ▶ Hadronic molecules: Weakly coupled charmed baryon and charmed meson.
- ▶ Hadro-charmonium states: compact bound $c\bar{c}$ state and light quarks.
- ▶ Quarks in a bag: Two tightly correlated di-quarks and an anti-quark.




Neutron Identification

- Qualitative arguments suggest there isn't strong evidence of photon contamination.
- For example, photons are more likely than neutrons to interact with multiple layers in the CLAS12 calorimeters.



ρ , ω and ϕ mesons

- ▶ Plotted here is the invariant mass of e^+e^- produced on a bound proton in the deuteron target.
- ▶ ϕ mesons are clearly resolved.
- ▶ ρ and ω mesons are unresolvable but clearly present.



Cross Section Calculation

- We can calculate the total cross section as:

$$\sigma_0(E_\gamma) = \frac{N_{J/\psi}}{N_\gamma \cdot \eta_T \cdot \omega_c \cdot Br \cdot \epsilon(E_\gamma)}$$

- Where:

- $N_{J/\psi}$ is the J/ψ yield in each E_γ bin
- N_γ is the sum of real and virtual photon flux
- η_T is the integrated luminosity
- ω_c is the Bethe Heitler normalisation factor
- Br is the branching ratio ($\sim 6\%$)
- $\epsilon(E_\gamma)$ is the acceptance in each E_γ bin

