



Hall C at 12 GeV
January 10, 2003

Moments and Lattice QCD

Wally Melnitchouk

Jefferson Lab

Outline

- What can Hall C @ 12 GeV do?
 - Structure functions over large kinematic range
 - L-T separation for $F_{1,2}(x, Q^2)$ up to $Q^2 = 12 \text{ GeV}^2$
 - Polarization asymmetries $A_{1,2}$ up to $Q^2 = 10 \text{ GeV}^2$
 - *Moments* of $F_{1,2}$ and $g_{1,2}$ structure functions
- What can lattice QCD actually calculate?
 - Lattice moments
 - Chiral extrapolation (π cloud)
 - Plans of LHPC (Lattice Hadron Physics Collaboration)

INELASTIC SCATTERING

$$\frac{d^2\sigma}{d\Omega dE'} = \frac{4\alpha^2 E'^2}{Q^4} \left[2 \sin^2 \frac{\theta}{2} W_1 + \cos^2 \frac{\theta}{2} W_2 \right]$$

$$\left. \begin{aligned} Q^2 &= 4EE' \sin^2 \frac{\theta}{2} \\ \nu &= E - E' \end{aligned} \right\} x = \frac{Q^2}{2M\nu}$$

BJORKEN LIMIT

$$Q^2, \nu \rightarrow \infty$$

x FIXED




$$M W_1(x, Q^2) \rightarrow F_1(x)$$


$$\nu W_2(x, Q^2) \rightarrow F_2(x)$$

ALSO $F_2(x) = 2x F_1(x)$

STRUCTURE FUNCTIONS IN PARTON MODEL



$$F_2(x) = x \sum_q e_q^2 q(x)$$
$$= \frac{4}{9} x u(x) + \frac{1}{9} x d(x) + \dots$$



Parton distributions in the proton

POLARIZED SCATTERING

$$\frac{d^2\sigma}{d\Omega dE'} \uparrow\uparrow \leftarrow \leftarrow = \frac{4\alpha^2 E'}{M\nu Q^2 E} \left[(E + E' \cos\theta) g_1 - 2Mx g_2 \right]$$

$$\frac{d^2\sigma}{d\Omega dE'} \uparrow \rightarrow \leftarrow \leftarrow = \frac{4\alpha^2 E'}{M\nu Q^2 E} \sin\theta \left[g_1 + \frac{2E}{\nu} g_2 \right]$$

QUARK-PARTON MODEL

$$g_1(x) = \frac{1}{2} \sum_q e_q^2 \Delta q(x)$$

\uparrow
 $q^\uparrow - q^\downarrow + \bar{q}^\uparrow - \bar{q}^\downarrow$

$$g_2(x) = g_2^{WW}(x) + g_2^{\text{twist-3}}(x)$$

$$\int_x^1 \frac{dy}{y} g_1(y) - g_1(x)$$


Wandzura - Wilczek

$$\langle \psi^\dagger \vec{B} \psi \rangle$$


quark-gluon correlations
(color \vec{B} & \vec{E} polarizabilities)

[Faint handwritten notes]

World data on F_1^p




World data on g_1^p



December 1998

ArXiv:9812101




GAUGE INVARIANCE

$\sigma^{\sigma p}$ FINITE AT $Q^2 = 0$


$$\sigma^{\sigma^* p} \propto \frac{F_2^p}{Q^2}$$

$\Rightarrow F_2^p (Q^2 \rightarrow 0) \rightarrow 0$



DUALITY DIAGRAMS FOR $ep \rightarrow eX$




≡




=



≡



=



COLLINS 1977

For γ^*N scattering at large s ,

$$F_2 \sim x^{1-\alpha(0)}$$


(since $x \sim 1/s$)


→ Valence (non-diffractive, Reggeon)

$$F_2^{\text{val}} \sim x^{1/2}$$

→ Sea (diffractive, Pomeron)

$$F_2^{\text{sea}} \sim x^0$$





Kinematic regions covered by global data on F_2^p

TRANSITION REGION $Q^2 < 1 \text{ GeV}^2$
 FOR $0.005 \lesssim x \lesssim 0.2$ UNEXPLORED

Longitudinal/Transverse Separation

Measured cross section

$$\frac{d\sigma}{d\Omega dE'} = \Gamma (\sigma_T(x, Q^2) + \epsilon \sigma_L(x, Q^2))$$

→ virtual photon flux Γ

→ virtual photon polarization

$$\epsilon = \left(1 + 2(1 + \nu^2/Q^2) \tan^2(\theta/2)\right)^{-1}$$

$\sigma_T(\sigma_L)$ cross section for photoabsorption of transverse (longitudinal) photon


$$\sigma_T = \frac{4\pi^2\alpha}{2x(W^2 - M^2)} 2xF_T$$
$$\sigma_L = \frac{4\pi^2\alpha}{2x(W^2 - M^2)} F_L$$

where $F_T = F_1$, $F_L = (1 + Q^2/\nu^2)F_2 - 2xF_1$

Rosenbluth separation


→ vary ϵ for fixed x and Q^2

→ fit intercept σ_T , slope σ_L



*Range of $\Delta\epsilon$ accessible at $x = 0.8$
in Hall C at 12 GeV*

$$R = \sigma_L / \sigma_T$$




World data on R for $x > 0.75$. Arrow indicates $W = 2 \text{ GeV}$ boundary for $x = 0.8$.


Spin Dependence of Transition

- Transition in spin dependent structure functions even more fascinating
 - structure functions need not be positive!
 - dramatic transition from (negative) GDH integral at $Q^2 = 0$ to (positive) Bjorken integral at $Q^2 \geq 2 \text{ GeV}^2$
- Are higher twist effects more important for g_1 than F_2 ?
- How does g_1 at low Q^2 approach the scaling curve?
- Transition for (twist-3) g_2 structure function?
- Complete absence of data in resonance region at high x


Testing duality in g_1

g_1 vs. Nachtmann Variable





Expected uncertainties for $A_{||}$ at $Q^2 = 8 \text{ GeV}^2$.



Lowest moments of F_2^p , compared with lattice calculations.

Structure Functions in Lattice QCD

Cannot calculate x -distribution, only moments

$$\langle x^n \rangle = \int_0^1 dx x^n (q(x) + (-1)^{n+1} \bar{q}(x))$$

where $\langle x^n \rangle$ are matrix elements of twist-2 operators (twist = dimension - spin):

$$\langle x^n \rangle_{p\mu_1 \cdots p\mu_{n+1}} = \langle N | \mathcal{O}_{\{\mu_1 \cdots \mu_{n+1}\}} | N \rangle$$

where


$$\mathcal{O}_{\{\mu_1 \cdots \mu_{n+1}\}} = \bar{\psi} \gamma_{\{\mu_1} \overleftrightarrow{D}_{\mu_2} \cdots \overleftrightarrow{D}_{\mu_{n+1}\}} \psi$$

On the lattice, operators transform according to hypercubic $H(4)$ subgroup of $O(4)$

→ because of operator mixing, only $n \leq 3$ moments can be calculated

Lattice Calculation of SF Moments

Matrix elements of twist-2 operators have contributions from **connected** and **disconnected** diagrams:



Disconnected diagrams are very noisy
→ at present only connected diagrams
can be reliably calculated on the lattice

However, disconnected contributions **cancel**
in (non-singlet) $u - d$ difference!


Reference	Q/U	Quark Action	Lattice	a [fm]	m_π [GeV]	Moments	Symbol
QCDSF [30]	Q	Wilson	$16^3 \times 32$	0.1	0.6 - 1.0	All	▲
QCDSF [31]	Q	Wilson	$24^3 \times 32$	0.1	0.35 - 0.6	All	■
QCDSF [32]	Q	NPIC	$16^3 \times 32$	0.1	0.6 - 1.0	All	×
QCDSF [33]	Q	NPIC	$16^3 \times 32$	0.1	0.65 - 1.2	$\langle 1 \rangle_{\Delta q}, \langle x^2 \rangle_{\Delta q}$	◆
	Q	NPIC	$24^3 \times 48$	0.075	0.7 - 1.2	$\langle 1 \rangle_{\Delta q}, \langle x^2 \rangle_{\Delta q}$	●
	Q	NPIC	$32^3 \times 48$	0.05	0.6 - 1.25	$\langle 1 \rangle_{\Delta q}, \langle x^2 \rangle_{\Delta q}$	*
MIT [20]	Q	Wilson	$16^3 \times 32$	0.1	0.58 - 0.82	All	★
MIT-SESAM [20]	U	Wilson	$16^3 \times 32$	0.1	0.63 - 1.0	All	◇
MIT-SCRI [20]	U	Wilson	$16^3 \times 32$	0.1	0.48 - 0.67	All	□
KEK [27, 28]	Q	Wilson	$16^3 \times 20$	0.14	0.52 - 0.97	$\langle 1 \rangle_{\Delta q}, \langle 1 \rangle_{\delta q}$	▼

TABLE I: Simulation parameters for lattice calculations of the moments of PDFs included in our analysis. Q/U corresponds to quenched/unquenched simulations, and NPIC denotes the nonperturbatively improved clover quark action. "All" moments correspond to $\langle x^i \rangle_q$ for $i = 1, 2, 3$, $\langle x^i \rangle_{\Delta q}$ for $i = 0, 1, 2$, and $\langle x^i \rangle_{\delta q}$ for $i = 0, 1, 1$. The symbols shown in the final column correspond to those plotted in Figs. 8, 9 and 10.

VOLUME LARGE ENOUGH TO INCLUDE π CLOUD


$$L \gtrsim 4/m_\pi$$

Linear Extrapolation





Chiral Symmetry and QCD

- In limit $m_q \rightarrow 0$, QCD has exact *chiral* $SU(N_f)_L \times SU(N_f)_R$ symmetry
- Spontaneous breaking of chiral symmetry
→ appearance of pseudoscalar Goldstone bosons (π, K)
- Pions play important role at small $m_\pi^2 \sim m_q$
→ expand observables in powers of m_π
(e.g. χPT)
- Pion loops in χPT generate non-polynomial terms which are *non-analytic* in m_q
(odd powers of m_π or $\log m_\pi$)
- Coefficients of non-analytic terms are **model-independent**
→ given in terms of g_A and f_π



WAVEFUNCTION
RENORMALIZATION

VERTEX
CORRECTIONS



WEINBERG-TOMOZAWA CONTACT
(TADPOLE)

Chiral Extrapolation of SF Moments

Extrapolation formula for n -th moment of $u-d$:

$$\langle x^n \rangle_{u-d} = a_n + b_n \frac{m_\pi^2}{m_\pi^2 + m_b^2} + c_n m_\pi^2 \log \left(\frac{m_\pi^2}{m_\pi^2 + \mu^2} \right)$$

Detmold, WM, Negele, Renner, Thomas,
Phys.Rev.Lett. **87** (2001) 172001


Thomas, WM, Steffens
Phys.Rev.Lett. **85** (2000) 2892

Coefficient of non-analytic term:

$$c_n = -a_n \frac{3g_A^2 + 1}{4\pi^2 f_\pi^2}$$

calculated in chiral perturbation theory

Arndt, Savage, nucl-th/0105045
Chen, Ji, hep-ph/0105197


Chiral Extrapolation ($\mu = 550$ MeV)

x dependence and Regge Trajectories

- What can lattice moments tell us about x dependence of parton distributions?
- Typical parameterization of (non-singlet) distributions (CTEQ, MRS, GRV)


$$x q(x) = a x^b (1-x)^c (1 + \epsilon\sqrt{x} + \gamma x)$$

- Fit parameters b and c to lattice data using extrapolation formula
→ quark distribution as function of m_q !
- Exponent b related to intercept of ρ Regge trajectory
→ relate m_q dependence of masses of $1^{--}, 2^{++}, 3^{--}, \dots$ mesons to $x \rightarrow 0$ behavior of structure function moments!



Extracted $x(u_v - d_v)$ distribution
at the physical quark mass

Detmold, WM, Thomas
Eur.Phys.J. C (2001)



Moments of $u_v - d_v$ distribution
 extracted from lattice data (scaled by 3^n)

Detmold, WM, Thomas
Eur.Phys.J. C (2001)

Spin Dependent Distributions

Moments of spin dependent distributions:

$$\langle \Delta x^n \rangle = \int_0^1 dx x^n (\Delta q(x) + (-1)^n \Delta \bar{q}(x))$$

helicity

$$\langle \delta x^n \rangle = \int_0^1 dx x^n (\delta q(x) + (-1)^{n+1} \delta \bar{q}(x))$$

transversity




obtained from matrix elements of twist-2 operators:

$$\langle N | \bar{\psi} \gamma_{\{\mu_1} \gamma_5 D_{\mu_2} \cdots D_{\mu_{n+1}} \} \psi | N \rangle \longrightarrow \langle \Delta x^n \rangle$$

$$\langle N | \bar{\psi} \sigma_{[\alpha \{\mu_1} \gamma_5 D_{\mu_2} \cdots D_{\mu_{n+1}} \} \psi | N \rangle \longrightarrow \langle \delta x^n \rangle$$



Leading nonanalytic chiral coefficients
recently calculated for both $\langle \Delta x^n \rangle$ and $\langle \delta x^n \rangle$

(Chen & Ji, hep-ph/0105197, hep-ph/0105296)




Riken-BNL \Rightarrow FINITE \checkmark

\rightarrow JURY SHALL OUT ON g_A^{lat}




The Promise of Lattice

- Lattice Hadron Physics Collaboration has ambitious long-term plan to calculate structure function moments from first principles
 - currently 128 nodes, 0.1 Tflops
 - another 256 nodes in FY03, 0.5 Tflops
 - expect 1 Tflops by FY04
 - **8 Tflops by FY06**
(c.f. largest cluster today 5 Tflops, Earth simulator \sim 30 Tflops)
- Simulations usually in two stages
 - computation of quark propagators **(expensive)**
 - tying propagators together into matrix elements **(easy)**
- Push closer to physical quark masses, $m_q \rightarrow m_q^{phys}$ ($m_q \propto m_\pi^2$)
 - simulations with $m_\pi \approx 300$ MeV currently running
- Larger volumes to include pion cloud



Momentum carried by $u-d$ quarks in proton,
with projected lattice errors on chiral extrapolation.



- Scale Q^2 determined by (inverse) lattice spacing, $Q^2 \sim 1/a^2$
 - currently $a \approx 0.1$ fm $\Leftrightarrow Q^2 \approx 4$ GeV²
 - extend to $a \approx 0.05$ fm to access $Q^2 \approx 16$ GeV²
BUT $\sim 2^4$ times more expensive
 - larger spacings, e.g. $a = 0.2$ fm $\Leftrightarrow Q^2 \approx 1$ GeV², but must match onto perturbative renormalization (e.g. $\overline{\text{MS}}$)
- Singlet quantities (e.g. $F_2^p + F_2^n$) require computationally difficult *disconnected insertions*
 - ~ 20 harder for currently accessible m_q
 - worse for smaller m_q
- Similar expectations for spin-dependent moments
 - possibly larger finite V effects for g_A
- Unquenching — inclusion of quark loops
 - **1000** times more expensive than quenched for quark propagators
 - more important at small m_q where quark loops not suppressed
 - expect to match today's accuracy of quenched simulations by \sim **2006**

Summary

- Reconstructing complete quark structure of nucleon requires measurement of structure functions over wide range of kinematics
- Hall C @ 12 GeV, with HMS-SHMS
→ (high \mathcal{L} , small θ , high \vec{p})
will provide unprecedented access to unpolarized ($F_{2,L}$) and polarized ($g_{1,2}$) structure functions and their *moments*
- Measurements will confront QCD directly through moments calculated in lattice QCD
→ complement long-term plans by LHPC