

GPD's in Lattice QCD

George T. Fleming

Yale University

[Lattice Hadron Physics Collaboration]

- GPD's and generalized form factors (GFF's).
- Summary of LHPC hadron structure program.
- Some preliminary lattice results.

LHPC Hadron Structure project on USQCD resources

Dru Renner

University of Arizona

Richard Brower, James Osborn

Boston University

Rebecca Irwin, Michael Ramsey-Musolf

California Institute of Technology

Robert Edwards, David Richards

Thomas Jefferson National Accelerator Facility

Bojan Bistrovic, Jonathan Bratt, Patrick Dreher, Oliver Jahn,

John Negele, Kostas Orginos, Andrew Pochinsky, Dmitry Sigaev

Massachusetts Institute of Technology

Matthias Burkardt, Michael Engelhardt

New Mexico State University

George Fleming

Yale University

Constantia Alexandrou, Antonios Tsapalis

University of Cyprus

Wolfram Schroers

DESY, Zeuthen

Philippe de Forcrand

ETH-Zürich and CERN

Philipp Hägler

Vrije Universiteit, Amsterdam

GPD's and Generalized Form Factors (GFF's)

- Experimentalists measure matrix elements of light cone operators

$$\langle P' S' | \mathcal{O}_\Gamma^q | PS \rangle = \left\langle P' S' \left| \bar{q} \left(-\frac{x^-}{2} \right) \Gamma \mathcal{P} \exp \left[-ig \int_{x^-/2}^{-x^-/2} A^+(y) dy \right] q \left(\frac{x}{2} \right) \right| PS \right\rangle$$

- They can be written in terms of generalized parton distributions (GPD's)

$$\begin{aligned} & \int \frac{dx^-}{2\pi} e^{ix^- \bar{p}^+ x} \langle P' S' | \mathcal{O}_\Gamma^q | PS \rangle \\ &= \overline{U}(P', S') [H^q(x, \xi, \Delta^2) \Gamma + E^q(x, \xi, \Delta^2) \frac{i\sigma \cdot \Delta}{2M}] U(P, S) \end{aligned}$$

- Eight GPD's in all: $H, E, \tilde{H}, \tilde{E}, H_T, E_T, \tilde{H}_T, \tilde{E}_T$
- Using OPE, light cone operators replaced by tower of local twist two operators

$$\langle P' S' | \mathcal{O}_\Gamma^{\mu_1 \dots \mu_n} | PS \rangle = \left\langle P' S' \left| \bar{q}(x) i D^{(\mu_1} \dots i D^{\mu_{n-1}} \Gamma^{\mu_n)} q(x) \right| PS \right\rangle$$

- They can be parameterized by generalized form factors (GFF's), *i. e.*

$$\begin{aligned} & \langle P' S' | \mathcal{O}_q^{\mu_1 \mu_2} | PS \rangle = \\ & \overline{U}(P', S') \left[A_{20}^q(Q^2) \gamma^{(\mu_1} \Delta^{\mu_2)} + B_{20}^q(Q^2) \frac{i\sigma^{(\mu_1 \alpha} \Delta^\alpha}{2M} \Delta^{\mu_2)} + C_2^q(Q^2) \frac{\Delta^{(\mu_1} \Delta^{\mu_2)}}{2M} \right] U(P, S) \end{aligned}$$

- Nine GFF's in all: $A_{ni}, B_{ni}, C_{ni}, \tilde{A}_{ni}, \tilde{B}_{ni}, A_{Tni}, B_{Tni}, \tilde{A}_{Tni}, \tilde{B}_{Tni}$

Equivalence of GPD's and GFF's

- GPD's and GFF's are formally equivalent by Mellin transformation *e. g.*

$$\int_{-1}^1 dx \ x^{n-1} H^q(x, \xi, Q^2) = \sum_{i=0, \text{ even}}^{n-1} A_{ni}^q(Q^2) (-2\xi)^i + \delta_{n, \text{ even}} C_n^q(Q^2) (-2\xi)^n$$

$$\int_{-1}^1 dx \ x^{n-1} E^q(x, \xi, Q^2) = \sum_{i=0, \text{ even}}^{n-1} B_{ni}^q(Q^2) (-2\xi)^i - \delta_{n, \text{ even}} C_n^q(Q^2) (-2\xi)^n$$

- Choice of GPD's *vs.* GFF's depends on physics.

GPD: PDF's and transverse PDF's

GFF: elastic form factors and nucleon spin

- In Euclidean lattice QCD, only GFF's can be computed directly.
- Many GFF's are familiar experimental quantities:

- $A_{10}^q(Q^2) = F_1^q(Q^2)$, $B_{10}^q(Q^2) = F_2^2(Q^2)$
- $\tilde{A}_{10}^q(Q^2) = G_A^q(Q^2)$, $\tilde{B}_{10}^q(Q^2) = G_P^q(Q^2)$,
- $J^q = \frac{1}{2} (A_{20}^q(0) + B_{20}^q(0))$, $\frac{1}{2}\Delta\Sigma^q = \tilde{A}_{10}^q(0)$
- $L^q = J^q - \frac{1}{2}\Delta\Sigma^q$
- $\langle x^{n-1} \rangle_q = A_{n0}^q(0)$, $\langle x^{n-1} \rangle_{\Delta q} = \tilde{A}_{n0}^q(0)$, $\langle x^{n-1} \rangle_{\delta q} = A_{Tn0}^q(0)$

Summary of LHPC hadron structure program

- Long term program to compute all $n \leq 4$ GFF's in dynamical lattice QCD.
- Current pion masses $m_\pi \approx 350 - 750$ MeV and lattice spacing $a \approx \frac{1}{8}$ fm.
- Status of the calculation

Operators	Matrix	Operator	GFF	
	elements	renorm.	extraction	Analysis
$\bar{q}\Gamma_\mu q$	Done!	Done!	Almost done	Starting
$\bar{q}\Gamma_{(\mu}D_{\nu)}q$	Done!	Done!	Almost done	Starting
$\bar{q}\Gamma_{(\mu}D_{\nu}D_{\rho)}q$	Done!	Done!	Almost done	Starting
$\bar{q}\Gamma_{(\mu}D_{\nu}D_{\rho}D_{\sigma)}q$	Not yet	Done!	Not yet	Not yet

- Only isovector flavor combinations for GFF's in this round.
- Finite perturbative renormalization needed to quote results in $\overline{\text{MS}}$ scheme.

$$\langle P' S' | \mathcal{O}_\Gamma^{\mu_1 \dots \mu_n} | PS \rangle_{\overline{\text{MS}}} = Z \langle P' S' | \mathcal{O}_\Gamma^{\mu_1 \dots \mu_n} | PS \rangle_{\text{latt}}$$

- Lighter pion masses $m_\pi \approx 250 - 350$ MeV finished by next year.

Perturbative renormalization of twist two matrix elements

Tree level: $Z = 1$, One loop HYP corrections: $< 10\%$.

operator	$H(4)$	NOS	HYP	APE
$\bar{q}[\gamma_5]q$	1^{\pm}_1	0.68	0.971	1.07
$\bar{q}[\gamma_5]\gamma_\mu q$	4^{\mp}_4	0.765	0.964	0.99
$\bar{q}[\gamma_5]\sigma_{\mu\nu}q$	6^{\mp}_1	0.821	0.987	0.989
$\bar{q}[\gamma_5]\gamma_{\{\mu}D_{\nu\}}q$	6^{\pm}_3	0.986	0.968	0.929
$\bar{q}[\gamma_5]\gamma_{\{\mu}D_{\nu\}}q$	3^{\pm}_1	0.972	0.962	0.925
$\bar{q}[\gamma_5]\gamma_{\{\mu}D_{\nu}D_{\alpha\}}q$	8^{\mp}_1	1.206	0.982	0.898
$\bar{q}[\gamma_5]\gamma_{\{\mu}D_{\nu}D_{\alpha\}}q$	mixing	8.78×10^{-3}	2.88×10^{-3}	1.26×10^{-3}
$\bar{q}[\gamma_5]\gamma_{\{\mu}D_{\nu}D_{\alpha\}}q$	4^{\mp}_2	1.191	0.98	0.898
$\bar{q}[\gamma_5]\gamma_{\{\mu}D_{\nu}D_{\alpha}D_{\beta\}}q$	2^{\pm}_1	1.375	0.989	0.876
$\bar{q}[\gamma_5]\sigma_{\mu\{\nu}D_{\alpha\}}q$	8^{\pm}_1	1.018	0.991	0.945
$\bar{q}[\gamma_5]\gamma_{[\mu}D_{\nu]}q$	6^{\mp}_1	0.967	0.973	0.983
$\bar{q}[\gamma_5]\gamma_{[\mu}D_{\{\nu\}}D_{\alpha\}}q$	8^{\pm}_1	0.931	0.937	0.947

Table 11.17: Full \overline{MS} to lattice renormalization coefficients for $M = 1.7$ and 1-loop expression for g . By chiral symmetry matrix elements are the same (except for parity) with and without γ_5 , and this is indicated by the $[\gamma_5]$ notation where the upper parity arises in the absence of γ_5 .


B. Bistrovic, Ph. D. Thesis, MIT, 2005

The Nucleon Electromagnetic Form Factors

- Brodsky-Lepage predicted $F_1 \sim Q^{-4}$ and $F_2 \sim Q^{-6}$ as $Q^2 \rightarrow \infty$.
- Older L/T separation experiments observed predicted behavior.
- New polarization transfer experiments do not see expected scaling.
- *One Photon Exchange* approximation may not be justified.
- Softer scaling also possible
(hep-ph/0212351)

$$\frac{Q^2}{\log^2(Q^2/\Lambda^2)} \frac{F_2(Q^2)}{F_1(Q^2)} \sim \text{const}$$


Phys. Rev. Lett. **88**, 092301 (2002)



Nucleon F_2/F_1 on the Lattice (I)

PRELIMINARY


- Only $I = 1$ form factors computed so far to avoid disconnected diagrams. $F_1^{I=1} = F_{1p} - F_{1n}$ but F_{1n}, F_{2n} not known accurately for $Q^2 \gtrsim 1 \text{ GeV}^2$.
- Our normalization is $F_2(Q^2) \rightarrow \kappa$ as $Q^2 \rightarrow 0$.



Nucleon F_2/F_1 on the Lattice (II)


PRELIMINARY

- $F_2^{I=1}/F_1^{I=1} \rightarrow \kappa_p - \kappa_n$ as $Q^2 \rightarrow 0$.
- PDG: $\kappa_p = 1.792847351(28)$
- PDG: $\kappa_n = -1.91304273(45)$
- So, comparison of $I = 1$ with $p - n$ could be OK with proper chiral extrapolation.




Nucleon axial charge g_A in a finite volume

- g_A is strongly suppressed by finite volume when $m_\pi L < 4$.
- Graph from hep-lat/0409161 (QCDSF). $m_\pi = 717$ MeV, curve is LO χ PT.




NLO χ PT (with Δ) extrapolation of g_A

- Fitting formula proposed by Hemmert *et al.* hep-lat/0303002.
- Two free parameters: g_{NN}^A and $g_{\Delta\Delta}^A$.
- Other parameters fixed by phenomenology: f_π , $g_{N\Delta}^A$, $m_\Delta - m_N$, $B_9 - g_A B_{20}$.



Transverse quark distributions




$$A_{n0}^q(-\Delta_\perp^2) = \int d^2 b_\perp e^{i \Delta_\perp \cdot b_\perp} \int_{-1}^1 x^{n-1} q(x, b_\perp)$$

$$\langle b_\perp^2 \rangle_{(n)}^q = -4 \frac{A_{n0}^{q'}(0)}{A_{n0}^q(0)}$$

$$\lim_{x \rightarrow 1} q(x, b_\perp) \propto \delta(b_\perp^2)$$

M. Burkardt hep-ph/0207047

- Higher moments A_{n0} weight $x \sim 1$.
- Slope of A_{n0}^q decreases as n increases.
- Slope of $A_{10}^{u-d}(0) = -0.93(4) \text{ (GeV)}^2$.
- Slope of $A_{30}^{u-d}(0) = -0.13(3) \text{ (GeV)}^2$.
- Will this continue at light pion masses?



D. Renner (LHPC/SESAM)

Summary and outlook

- Large scale computation of isovector matrix elements ($n \leq 3$) is done. Data analysis is proceeding rapidly. Expect published results soon.
- Isoscalar and strange matrix elements are $\mathcal{O}(10) - \mathcal{O}(100)$ times harder to compute due to statistical noise. We'll make our first serious attempt this year.
- Perturbative renormalization complete. Bojan Bistrovic (MIT)
- Reaching higher Q^2 is high priority for nucleon form factors.