

## Vertical angle determination

We try to determine the vertical angle of the magnetic field's direction by fitting the following formula to the data:

$$\phi = \arctan \left( \frac{\sqrt{(I_s + A)^2 + R^2(I_l + B)^2}}{K(I_v + E)} \right)$$

Here  $I_s$  is the current in the small coil,  $I_l$  the current in the large coil and  $I_v$  the current in the vertical coil.

A, B, E, R, K are free parameters. A, B, E represent corrections in the all three directions of the components of the magnetic field; R and K are scaling factors.

The angle is counted as zero when the field is vertical.

We can get an estimate of the parameters, which we will use in fitting of the whole formula, by examining the data with only one of the planar coils switched on. Here, assuming the correction B (or A) is much smaller than the field generated by  $I_s$  or  $I_l$ , the above formula simplifies to:

$$\Phi = \arctan \left( \frac{(I_s + A)}{(K(I_v + E))} \right)$$

and

$$\Phi = \arctan \left( \kappa \frac{(I_l + B)}{(I_v + E)} \right)$$

where  $\kappa$  is R/K.

### Measured data:

#### Small coil + vertical coil:

| $I_s$ | $I_v$ | Compass | Compass - reverse |
|-------|-------|---------|-------------------|
| 7     | 0     | 1990    | 11                |
| 5     | 10    | 1462.5  | 537               |
| 0     | 14    | 997.5   | 1002              |
| -5    | 10    | 534     | 1466              |
| -7    | 0     | 7       | 1994              |

The first column is  $I_s$ , second  $I_v$ ; the third and fourth columns are reading taken from the compass, the second one after rotating it round the vertical axis by  $180^\circ$ .

We first use both readings to find the zero of the compass. Looking at our formula, we see that we must limit ourselves to the interval  $(0, \pi/2)$ . The angle is then given by the deviation of the measurement from the compass' zero.

The (average) zero of the compass in this measurement is **1000.1**.

Data from the first measurement, with the angle in radians:

| $I_s$ | $I_l$ | $I_v$ | Angle (rad) |
|-------|-------|-------|-------------|
| 7     | 0     | 0     | 1.5543      |
| 5     | 0     | 10    | 0.7269      |
| 0     | 0     | 14    | 0.0035      |
| -5    | 0     | 10    | 0.7320      |
| -7    | 0     | 0     | 1.5606      |

(Here I wrote the values of all three currents:  $I_s$ ,  $I_l$ , and  $I_v$ ).

The error for the angle is  **$\pm 0.0007854$**  (roughly  $0.05^\circ$ ) for all values.

Large coil + vertical coil:

| $I_l$ | $I_v$ | Compass | Compass - reverse |
|-------|-------|---------|-------------------|
| -7    | 0     | 3972    | 1959              |
| -5    | 10    | 518     | 1411.5            |
| 0     | 14    | 983     | 947               |
| 5     | 10    | 1438    | 491               |
| 7     | 0     | 1955    | 3976              |

To avoid problems using mod(4000), we re-brand the measurements over 2000 as negative:

| $I_l$ | $I_v$ | Compass | Compass - reverse  |
|-------|-------|---------|--|
| -7    | 0     | -28     | 1959   |
| -5    | 10    | 518     | $\Phi = \arctan \left( \kappa \frac{(I_l + I_v)}{(I_v + I_l)} \right)$<br>1411.5 |
| 0     | 14    | 983     | 947  |
| 5     | 10    | 1438    | 491  |
| 7     | 0     | 1955    | -24  |

The (average) zero of the compass in this measurement is **965.5** .

Data from the second measurement:

| $I_s$ | $I_l$ | $I_v$ | Angle (rad) |
|-------|-------|-------|-------------|
| 0     | -7    | 0     | 1.5610      |
| 0     | -5    | 10    | 0.7018      |
| 0     | 0     | 14    | 0.0283      |
| 0     | 5     | 10    | 0.7438      |
| 0     | 7     | 0     | 1.5543      |

First, I tried fitting both the “partial” formulas to get an idea for the parameters' values.

Small coil fit:

**A = -0.026297**      +/- 0.03113      (118.4%)  
**K = 0.549914**      +/- 0.004105      (0.7465%)  
**E = 0.175074**      +/- 0.0401      (22.9%) Arial

Large coil fit:

**$\kappa$  = 1.79487**      +/- 0.01537      (0.8565%)  
**B = 0.210944**      +/- 0.03583      (16.98%)  
**E = 0.171157**      +/- 0.04543      (26.54%)

Using these as starting points for the fit of the whole function (with  $R = \kappa * K$ ), we get these values for the parameters:



**A = -0.0195053**      +/- 0.06669      (341.9%)  
**K = 0.550365**      +/- 0.008133      (1.478%)  
**E = 0.168082**      +/- 0.06048      (35.98%)  
**B = 0.157533**      +/- 0.0419      (26.6%)  
**R = 0.987262**      +/- 0.01873      (1.898%)


The total error of our fit is a function of the three currents with the values of parameters and their error as above. Here is the measured angle, calculated angle, their difference, the absolute error of the fit at that value, and the relative error of the fit, for every measurement. All angles are in **degrees**.

| $I_s$ | $I_l$ | $I_v$ | measured<br>[°] | calculated<br>[°] | $\Delta(\text{meas.-calc.})$ [°] | $d\phi$<br>[°] | $\frac{d\phi}{\phi}$<br>[%] |
|-------|-------|-------|-----------------|-------------------|----------------------------------|----------------|-----------------------------|
| 7     | 0     | 0     | 89.2409         | 89.0325           | 0.2085                           | 0.2917         | 0.3276                      |
| 5     | 0     | 10    | 41.6825         | 41.6371           | 0.0454                           | 0.9783         | 2.3497                      |
| 0     | 0     | 14    | 1.1522          | 0.2024            | 0.9498                           | 0.4060         | 200.5440                    |
| -5    | 0     | 10    | 41.9044         | 41.9295           | 0.0251                           | 0.9761         | 2.3280                      |
| -7    | 0     | 0     | 89.2452         | 89.3929           | 0.1477                           | 0.2900         | 0.3244                      |
| 0     | -7    | 0     | 89.2159         | 89.3929           | 0.1770                           | 0.3134         | 0.3506                      |
| 0     | -5    | 10    | 40.5229         | 40.1975           | 0.3254                           | 1.3695         | 3.4070                      |
| 0     | 0     | 14    | 1.1522          | 1.6190            | 0.4674                           | 0.4060         | 25.0680                     |
| 0     | 5     | 10    | 42.3142         | 42.6043           | 0.2901                           | 1.3651         | 3.2041                      |
| 0     | 7     | 0     | 89.2504         | 89.0325           | 0.2179                           | 0.2994         | 0.3363                      |


**Graphs:**



Small coil:





Large coil:





## Planar angle determination

Here, the formula is:

$$\phi = \arctan \left( \frac{R (I_l + B)}{(I_s + A)} \right)$$

Parameters R, B, and A are the same as in the vertical case. (R is the scaling factor between the two coils, A and B are corrections in both directions).

We get the angle from the measurements, which are in the *Hall system* (i.e. Cartesian):

$$\tan(\Phi) = \frac{x}{z}$$

For least-squares fitting, the following expression is often more robust:

$$\frac{x}{z} = \frac{R (I_l + B)}{(I_s + A)}$$

I used both formulas, and will show the results from both.

Here, a mechanical compass (a needle) was used; (z, x) are the coordinates of the compass' endpoints, effectively defining two angles for every measurement.

The *coil system* is rotated from the *Hall system* by **143°**. (-z<sub>coil</sub> axis is rotated 37° degrees from the z axis). All data was transformed into the coil system prior to analysis.

Because I was using *atan2*, the angles were transformed to [-π, π] .

**Measured data:**With BigBite ON:

| $I_s$ | $I_l$ | Z       | X       |
|-------|-------|---------|---------|
| 4.2   | -5.6  | -5.08   | 231.36  |
|       |       | 8.53    | -225.24 |
| -1.0  | -6.9  | 163.58  | 164.09  |
|       |       | -160.76 | -157.66 |
| -5.6  | -4.2  | 230.12  | -1.63   |
|       |       | -226.93 | 10.15   |
| -6.9  | 1.0   | 124.34  | -194.29 |
|       |       | -136.74 | 180.47  |
| -4.2  | 5.6   | -8.92   | -235.87 |
|       |       | -3.86   | 221.07  |
| 1.0   | 6.9   | -169.16 | -168.16 |
|       |       | 155.83  | 153.44  |
| 5.6   | 4.2   | -234.87 | -11.25  |
|       |       | 222.11  | -4.57   |
| 6.9   | -1.0  | -149.41 | 170.82  |
|       |       | 135.21  | -186.50 |

(The two sets of coordinates corresponding to each set of currents represent the coordinates of the endpoints of the needle. Each pair of coordinates in a set is always the same endpoint. So far, the currents are in the *Coil System* and the coordinates in the *Hall System*.)

With BigBite OFF:

| $I_s$ | $I_l$ | Z       | X       |
|-------|-------|---------|---------|
| 4.2   | -5.6  | -8.52   | 229.46  |
|       |       | 1.88    | -227.36 |
| -5.6  | -4.2  | 225.13  | 0.12    |
|       |       | -231.79 | 4.20    |
| -4.2  | 5.6   | -4.84   | -238.88 |
|       |       | -3.00   | 217.99  |
| 5.6   | 4.2   | -235.88 | -3.15   |
|       |       | 220.98  | -3.36   |

To determine the angle, we must do the following:

- Find the zero of the compass – subtract the average value of a coordinate in a pair from each endpoint. Thus we get symmetric pairs, differing only in their sign.
- Use atan2 to get the angle, ranging from  $-\pi$  to  $\pi$ .
- Rotate the angle by  $143^\circ$ , taking care of the discontinuity at  $180^\circ$

The algorithm for deriving the data, used for fitting the simplified formula (“*the quotient*”), is a bit different:

- Find the zero of the compass – subtract the average value of a coordinate in a pair from each endpoint. Again we get symmetric pairs, differing only in their sign. (We only get one angle per pair of currents)
- Rotate the whole dataset of coordinates by  $143^\circ$ .
- Compute the quotient  $x'/z'$ .

### The final set of data:

#### With BigBite ON:

| $I_s$ | $I_l$ | $\Phi$ (rad)<br>= $\text{atan2}(x', z')$ | $d\Phi$<br>(rad) | $x'/z'$ (the<br>“quotient”) | $d(x'/z')$ |
|-------|-------|--|------------------|-----------------------------|------------|
| 4.2   | -5.6  | -0.89523                                 | 0.00006          | -1.24788                    | 0.00016    |
| -1.0  | -6.9  | -1.71443                                 | 0.00005          | 6.91414                     | 0.00242    |
| -5.6  | -4.2  | -2.52159                                 | 0.00006          | 0.71391                     | 0.00009    |
| -6.9  | 1.0   | 2.82505                                  | 0.00006          | -0.32756                    | 0.00006    |
| -4.2  | 5.6   | 2.20549                                  | 0.00006          | -1.35808                    | 0.00017    |
| 1.0   | 6.9   | 1.42593                                  | 0.00005          | 6.85442                     | 0.00238    |
| 5.6   | 4.2   | 0.66039                                  | 0.00006          | 0.77673                     | 0.00010    |
| 6.9   | -1.0  | -0.25240                                 | 0.00005          | -0.25790                    | 0.00006    |

#### With BigBite OFF:

| $I_s$ | $I_l$ | $\Phi$ (rad)<br>= $\text{atan2}(x', z')$ | $d\Phi$<br>(rad) | $x'/z'$ (the<br>“quotient”) | $d(x'/z')$ |
|-------|-------|--|------------------|-----------------------------|------------|
| 4.2   | -5.6  | -0.90226                                 | 0.00006          | -1.26603                    | 0.00016    |
| -5.6  | -4.2  | -2.50475                                 | 0.00006          | 0.73965                     | 0.00009    |
| -4.2  | 5.6   | 2.21254                                  | 0.00006          | -1.33822                    | 0.00017    |
| 5.6   | 4.2   | 0.64531                                  | 0.00006          | 0.75283                     | 0.00010    |

The measurement error in coordinates is **0.01 mm**. The errors are computed using this value, and used as weights in least-square fitting.

### Data analysis:

I used two approaches, the first one (called “*atan2*” from here on), was to calculate the measured angle and then fit the arctan formula. The second one (called “*the quotient*” from here on) fitted the ratio of both currents (plus corrections etc) to the quotient of the coordinates,

For fitting, I used two tools: *MATLAB* and *gnuplot*. The regression statistics they provide are a bit different. *MATLAB* also has the option of using robust non-linear regression. All the fits are displayed in the following table:

BigBite ON:

| Approach                             | Tool           | A              | dA      | B              | dB      | R              | dR      | STATS   |
|--------------------------------------|----------------|----------------|---------|----------------|---------|----------------|---------|---|
| <i>atan2</i>                         | <i>MATLAB</i>  | <b>0.04421</b> | 0.40520 | <b>0.21440</b> | 0.38940 | <b>1.02100</b> | 0.11220 | SSE: 2.13e-06<br>R-square: 0.9985<br>Adjusted R-square: 0.9979<br>RMSE: 0.0006527     |
| <i>atan2</i> -<br><i>LAR(robust)</i> | <i>MATLAB</i>  | <b>0.03747</b> | 0.2912  | <b>0.205</b>   | 0.2871  | <b>0.994</b>   | 0.0796  | SSE: 1.128e-06<br>R-square: 0.9992<br>Adjusted R-square: 0.9989<br>RMSE: 0.0004749    |
| <i>atan2</i>                         | <i>gnuplot</i> | <b>0.04464</b> | 0.31530 | <b>0.19184</b> | 0.31760 | <b>1.03845</b> | 0.10140 | rms of residuals : 1618.39<br>variance of residuals (reduced chisquare) : 2.61917e+06 |
| <i>quotient</i>                      | <i>MATLAB</i>  | <b>0.03236</b> | 0.02470 | <b>0.19340</b> | 0.16980 | <b>0.99750</b> | 0.00230 | SSE: 2.853e-06<br>R-square: 0.9999<br>Adjusted R-square: 0.9999<br>RMSE: 0.0007554    |
| <i>quotient</i>                      | <i>gnuplot</i> | <b>0.04008</b> | 0.28730 | <b>0.18507</b> | 0.31360 | <b>1.02990</b> | 0.09350 | rms of residuals : 1570.67<br>variance of residuals (reduced chisquare) : 2.46701e+06 |

BigBite OFF:

| Approach                                | Tool           | A                   | dA      | B                | dB      | R               | dR       | STATS  |
|---|----------------|---------------------|---------|------------------|---------|-----------------|----------|--|
| <i>atan2</i>                            | <i>MATLAB</i>  | -<br><b>0.01952</b> | 0.7635  | <b>0.1575</b>    | 0.7646  | <b>0.9873</b>   | 0.159    | SSE: 3.474e-08<br>R-square: 1<br>Adjusted R-square: 0.9999<br>RMSE: 0.0001864      |
| <i>atan2</i><br>-<br><i>LAR(robust)</i> | <i>MATLAB</i>  | <b>0.04412</b>      | 0.6622  | <b>0.08593</b>   | 0.6638  | <b>0.981</b>    | 0.1367   | SSE: 2.616e-08<br>R-square: 1<br>Adjusted R-square: 0.9999<br>RMSE: 0.0001618      |
| <i>atan2</i>                            | <i>gnuplot</i> | <b>0.0567881</b>    | 0.04545 | <b>0.0797418</b> | 0.04552 | <b>0.985554</b> | 0.009442 | rms of residuals: 149.841<br>variance of residuals (reduced chisquare) : 22452.2   |
| <i>quotient</i>                         | <i>MATLAB</i>  | <b>0.05389</b>      | 0.2662  | <b>0.08174</b>   | 0.3240  | <b>0.9793</b>   | 0.0442   | SSE: 3.224e-08<br>R-square: 0.9999<br>Adjusted R-square: 0.9998<br>RMSE: 0.0001796 |
| <i>quotient</i>                         | <i>gnuplot</i> | <b>0.05678</b>      | 0.04528 | <b>0.07973</b>   | 0.04559 | <b>0.98542</b>  | 0.00942  | rms of residuals: 150.017<br>variance of residuals: 22505.2                        |

Finally, to get a bigger sample size, we try fitting both datasets together:

BigBite ON + BigBite OFF:

| Approach                                | Tool           | A                | dA     | B               | dB      | R              | dR      | STATS  |
|---|----------------|------------------|--------|-----------------|---------|----------------|---------|--|
| <i>atan2</i>                            | <i>MATLAB</i>  | <b>0.04917</b>   | 0.2155 | <b>0.1683</b>   | 0.2120  | <b>1.004</b>   | 0.0531  | SSE: 2.206e-06<br>R-square: 0.999<br>Adjusted R-square: 0.9988<br>RMSE: 0.0004951    |
| <i>atan2</i><br>-<br><i>LAR(robust)</i> | <i>MATLAB</i>  | <b>0.02925</b>   | 0.1401 | <b>0.1575</b>   | 0.1398  | <b>0.9838</b>  | 0.0338  | SSE: 9.453e-07<br>R-square: 0.9996<br>Adjusted R-square: 0.9995<br>RMSE: 0.0003241   |
| <i>atan2</i>                            | <i>gnuplot</i> | <b>0.0464188</b> | 0.1992 | <b>0.15999</b>  | 0.2041  | <b>1.01274</b> | 0.05453 | rms of residuals : 1227.24<br>variance of residuals (reduced chisquare): 1.50613e+06 |
| <i>quotient</i>                         | <i>MATLAB</i>  | <b>0.02638</b>   | 0.0125 | <b>0.152</b>    | 0.0859  | <b>0.9975</b>  | 0.0013  | SSE: 3.258e-06<br>R-square: 0.9999<br>Adjusted R-square: 0.9999<br>RMSE: 0.0006017   |
| <i>quotient</i>                         | <i>gnuplot</i> | <b>0.04307</b>   | 0.1872 | <b>0.151802</b> | 0.19960 | <b>1.00878</b> | 0.05132 | rms of residuals: 1187.95<br>variance of residuals (reduced chisquare): 1.41122e+06  |

To get a better illustration, here's a table of measured angles, calculated angles, their difference, the absolute and the relative error in that point. (The error is a function of both currents).

I used both sets of measurements, and for the parameters' values I chose the “*quotient with MATLAB*” fit, which is second-to-last in the previous table. All angles are in **degrees**.

| $I_s$ | $I_l$ | measured<br>[°] | calculated[°<br>] | $\Delta(\text{meas.}-\text{calc.})$ [°] | $d\phi$<br>[°] | $\frac{d\phi}{\phi}$<br>[%] |
|-------|-------|-----------------|-------------------|---|----------------|-----------------------------|
| 4.2   | -5.6  | -51.6958        | -52.1274          | 0.4316                                  | 0.5561         | 1.0668                      |
| 4.2   | -5.6  | -51.2927        | -52.1274          | 0.8347                                  | 0.5561         | 1.0668                      |
| -1.0  | -6.9  | -98.2297        | -98.2304          | 0.0007                                  | 0.2181         | 0.2221                      |
| -5.6  | -4.2  | -144.4764       | -144.0781         | 0.3983                                  | 0.6742         | 0.4679                      |
| -5.6  | -4.2  | -143.5116       | -144.0781         | 0.5665                                  | 0.6742         | 0.4679                      |
| -6.9  | 1.0   | 161.8634        | 170.5092          | 8.6458                                  | 0.7239         | 0.4246                      |
| -4.2  | 5.6   | 126.3656        | 126.0326          | 0.3329                                  | 0.5242         | 0.4159                      |
| -4.2  | 5.6   | 126.7692        | 126.0326          | 0.7366                                  | 0.5242         | 0.4159                      |
| 1.0   | 6.9   | 81.6996         | 81.6986           | 0.0010                                  | 0.2101         | 0.2571                      |
| 5.6   | 4.2   | 37.8375         | 37.6525           | 0.1849                                  | 0.6446         | 1.7121                      |
| 5.6   | 4.2   | 36.9737         | 37.6525           | 0.6789                                  | 0.6446         | 1.7121                      |
| 6.9   | -1.0  | -14.4612        | -6.9627           | 7.4985                                  | 0.7198         | 10.3380                     |

### Graphs:

Both currents go through these values: (-6.9, -5.6, -4.2, -1, 1, 4.2, 5.6, 6.9). I decided to plot the angle as a function of the current in the small coil, keeping the current in the large coil fixed, resulting in 8 graphs total.

