

DVCS WITH LONGITUDINALLY POLARIZED TARGET USING CLAS AT 6 GEV



Angela Biselli
Fairfield University
For the CLAS collaboration

Outline



- ▶ Background
- ▶ Previous measurements
- ▶ Experimental apparatus
- ▶ Analysis procedure
- ▶ Expected results

Generalized Parton Distributions (GPD)



3-dimensional quark structure of nucleon



Elastic Scattering
transverse quark
distribution in
coordinate space



Deep inelastic scattering
Longitudinal quark
Distribution in
momentum space




Deep exclusive scattering
Fully-correlated
quark distribution in
both coordinate and
momentum space – GPDs

Deeply Virtual Compton Scattering and GPDs

► DVCS is the cleanest process to access GPDs

Handbag mechanism: the exclusive process can be factorized into a hard scattering part and a nucleon structure part parameterized via GPDs



Link between GPDs, DIS and form factors

- ▶ DIS at $t=\xi=0$: ordinary parton distributions

$$H^q(x, \xi = 0, t = 0) = q(x)$$

unpolarized quark distributions

$$\tilde{H}^q(x, \xi = 0, t = 0) = \Delta q(x)$$

polarized quark distributions

$$E^q, \tilde{E}^q$$

new information

- ▶ First moments: form factors

Dirac

$$\int_{-1}^1 dx H^q(x, \xi, t) = F_1^q(t)$$

axial

$$\int_{-1}^1 dx \tilde{H}^q(x, \xi, t) = G_A^q(t)$$

Pauli


$$\int_{-1}^1 dx E^q(x, \xi, t) = F_2^q(t)$$

pseudo-scalar

$$\int_{-1}^1 dx \tilde{E}^q(x, \xi, t) = G_P^q(t)$$

Accessing GPDs via DVCS

DVCS
BH



$$\frac{d^4\sigma}{dQ^2 dx_B dt d\phi} \approx |T^{DVCS} + T^{BH}|^2$$


GPDs
Elastic form factors

$$\frac{\sigma^+ - \sigma^-}{\sigma^+ + \sigma^-} \propto \frac{I(DVCS \bullet BH)}{|DVCS|^2 + |BH|^2 + I}$$

BH and DVCS interference is helicity dependent


DVCS can be measured via beam and target asymmetries

Cross section of $ep \rightarrow ep\gamma$ at $Q^2=2 \text{ GeV}/c^2$ and $X_B=0.35$



Polarization observables

$$A = \frac{\sigma^+ - \sigma^-}{\sigma^+ + \sigma^-} \quad \begin{array}{l} \xi = x_B/(2-x_B) \\ k = -t/4M^2 \end{array}$$



Polarized beam, unpolarized target

$$\Delta\sigma_{LU} \sim \sin\phi \operatorname{Im}\{F_1 H + \xi(F_1 + F_2)\tilde{H} + kF_2 E\}d\phi \quad \longrightarrow \quad H, \tilde{H}, E$$

Unpolarized beam, longitudinal target

$$\Delta\sigma_{UL} \sim \sin\phi \operatorname{Im}\{F_1 \tilde{H} + \xi(F_1 + F_2)(H + ..)\}d\phi \quad \longrightarrow \quad H, \tilde{H}$$

Unpolarized beam, transverse target


$$\Delta\sigma_{UT} \sim \sin\phi \operatorname{Im}\{k(F_2 H - F_1 E) + ...\}d\phi \quad \longrightarrow \quad H, E$$

Beam charge asymmetry


$$\Delta\sigma_C \sim \cos\phi \operatorname{Re}\{F_1 H + \xi(F_1 + F_2)\tilde{H} - kF_2 E\}d\phi \quad \longrightarrow \quad H, \tilde{H}, E$$

Cross section and BSA measurements

CLAS
Beam-spin asymmetry




Hall A
Cross section




HERMES

BSA A. Airapetian et al, Phys. Rev. D 75 (2007)
BCA A. Airapetian et al, Phys. Rev. Lett 96 (2001)



TSA measurements

S. Chen et al., PRL 97 (2006)



CLAS

Data were collected as a by-product during the **Eg1 2000** run:
5.7 GeV
with NH₃ longitudinally polarized target, Q^2 up to 4.5 GeV²

Non dedicated experiment


Target-spin (L) asymmetry (**H**, **\tilde{H}**)

HERMES

TSA with transverse polarized target

Also longitudinal polarized target asymmetry

A. Airapetian et al., JHEP 0806 (2008)



Target-spin (T) asymmetry (**H**, **E**)

DVCS at 6 GeV with polarized target and polarized beam using the CLAS detector

H. Avakian, S. Boyarinov, V.D. Burkert, **L. Elouadrhiri**¹, M. Ito, Y. Sharabian, E. Smith, S. Stepanyan
Jefferson Lab

J. Ball, P. Bertin, A. Camsonne, R. De Masi, F. Descamps, M. Garçon, F-X. Girod, M. Guidal, H-S. Jo,
M. Mazouz, M. Mac Cormick, B. Michel, C. Munoz Camacho, **S. Niccolai**¹, F. Sabatie, E. Voutier
LPC (Clermont) / LPSC (Grenoble) / IPNO (Orsay) / SPhN (Saclay)
CEA / DSM/DAPNIA & CNRS/IN2P3

K. Joo¹, N. Markov, M. Ungaro, B. Aaho
University of Connecticut

A. Biselli¹
Fairfield University

S. Kuleshov, O. Pogorelko, P. Shiskov
Institute of Theoretical and Experimental Physics, Moscow

H. Aznauryan, N. Dashyan, K. Egiyan
Yerevan Physics Institute

V. Koubarovsky, P. Stoler
Rensselaer Polytechnic Institute

M. Amarian, G. Dodge, H. Juengst, S.E. Kuhn, A. Radyushkin, L. Weinstein
Old Dominion University

M. Vanderhaeghen
College of William and Mary

S. Chen
Florida State University

and the CLAS collaboration

¹Co-spokesperson

DVCS at 6 GeV with polarized target and polarized beam using the CLAS detector

- ▶ First dedicated DVCS experiment with the polarized target
- ▶ Approved for 60 PAC days
- ▶ According to the current accelerator schedule: Feb 4th – Mar 11th @ 5.9 GeV
Apr 9th – May 31st @ 6.1 GeV
Aug 21st – Sept 20th @ 5.9 GeV


Jefferson Lab



Continuous
Electron
Beam
Accelerator
Facility


$I_{\max} = 200 \mu\text{A}$
 $E_{\max} = 6 \text{ GeV}$
 $\sigma_E/E \sim 2.5 \cdot 10^{-5}$
Beam Pol $\sim 80\%$

The CLAS detector




- Six individually instrumented sectors
- Toroidal magnetic field
- Multi-particle final state
- Large acceptance

CEBAF
Large
Acceptance
Spectrometer




The polarized target




Targets:

- $^{14}\text{NH}_3$ $P \sim 70 - 80\%$
- ^{12}C & He-4 background studies





- Polarized by DNP along the beam
- 5 Tesla magnetic holding field
- He-4 cooling bath $T=1\text{ K}$
- Field uniformity $\Delta B/B \sim 10^{-4}$

The Inner Calorimeter (IC)





- ▶ Designed for the e1-DVCS experiment
- ▶ Increase γ acceptance
 - EC: $17^\circ < \theta < 43^\circ$
 - IC: $4^\circ < \theta < 15^\circ$
- ▶ Better resolution



- 424 PbWO_4 crystals
- 16 cm x 1.3 cm x 1.3 cm
- Pointing geometry
- ~ 1.2 degree/crystal
- 18 radiation lengths
- APD readout




Moller background

- ▶ Moller background “naturally” shielded by the target field





Analysis

- ▶ Detection of all final state particles e p γ
- ▶ DVCS: $Q^2 > 1 \text{ GeV}^2$, $W > 2 \text{ GeV}$, $-t < 0.6 \text{ GeV}^2$
- ▶ Background from unpolarized target nucleons and π^0 highly reduced by geometry cut




Kinematics and acceptance




- Improved acceptance as compared to eg1
- Much higher statistics \rightarrow bins in t and x_B

Target spin asymmetry: ϕ dependence




Target spin asymmetry: Q^2 dependence

$$x_B=0.3, t = 0.325 \text{ GeV}^2, \phi = 90^\circ$$




Target spin asymmetry: t dependence

$$x_B = .3, Q^2 = 2.3 \text{ GeV}^2, \phi = 90^\circ$$



Semi-Inclusive Pion Production with a Longitudinally Polarized Target at 6 GeV

- ▶ Concurrent to DVCS experiment
- ▶ Single- and double spin azimuthal asymmetries in semi-inclusive electroproduction of pions using the 6 GeV
- ▶ Study of transverse momentum dependent parton distributions at $Q^2 > 1 \text{ GeV}^2$



Conclusions


- ▶ First dedicated DVCS experiment with polarized target
- ▶ Improvement on statistical and systematic errors by a factor 4 and 2.
- ▶ The large acceptance will allow to study x_B , Q^2 and t dependence of the DVCS amplitude
- ▶ The target asymmetry will increase the sensitivity to H , adding new information to existing polarized beam data.
- ▶ This experiment together with polarized beam experiment with the CLAS detector and the Hall-A DVCS program, will greatly contribute to our knowledge of GPDs and will provide a crucial step for the 12 GeV program.

Target polarization

- ▶ Target polarization extracted by measuring elastic asymmetry


$$A_{meas} = \frac{N^+ - N^-}{N^+ + N^-} = P_e P_t A_{theo}$$

- ▶ IC–target distance is important for electron detection




Target spin asymmetry: x_B dependence

$$t=0.325 \text{ GeV}^2, Q^2 = 2.3 \text{ GeV}^2, \phi = 90^\circ$$



Target spin asymmetry: Q^2 dependence

$$x_B=0.3, t = 0.325 \text{ GeV}^2, \phi = 90^\circ$$



Target spin asymmetry: t dependence

$$x_B = .3, Q^2 = 2.3 \text{ GeV}^2, \phi = 90^\circ$$

