



Learning New Chemistry and Physics with THz Light



Charles Schmuttenmaer
Chemistry Department,
Yale University

September 20, 2004

Overview



Characteristics of THz Pulse Spectroscopy

- measures electric field, $E(t)$, directly $\Rightarrow \hat{n}(\omega) = n(\omega) - ik(\omega)$
- minimum detectable fractional absorption better than 1 in 10^4
- **transient FIR absorption with sub-picosecond temporal resolution !**

$$1 \text{ THz} = 33.333 \text{ cm}^{-1} = 4 \text{ meV} = 300 \text{ micron } \lambda$$

Transient Photoconductivity

- semiconductors (ultrafast switches)
- **Semiconductor nanocrystals**
- **nano-electronics, molecular electronics**
- **thermalization and relaxation processes**
- **frequency dependent conductivity**

Binary Liquid Mixtures


- H_2O & CH_3OH , CH_3CN , CH_3COCH_3
- CH_3OH & CH_3CN , CH_3COCH_3
- OH librational dynamics
- reverse micelles

Intramolecular Electron Transfer

- accelerating charge generates pulse

THz Imaging

- plastic and paper are transparent




Relation to JLab THz Source ...

Much of this could be done with accelerator-based sources

- 100 fs pulse duration, or shorter.
- μJ to mJ per pulse, or more.
- kHz rep rates, or higher.
- At least 100 ns, or longer, between pulses for some studies.
- Synchronize intense NIR/visible/UV/x-ray pulses with far-IR.


- Nonlinear spectroscopy (2D THz, photon echoes, etc).
- Rapid, large-area THz imaging.
- (Ultra)high magnetic fields.

How do we stack up?




Two-Dimensional THz Spectroscopy

- Far infrared analog of two-dimensional NMR spectroscopy
- Spectra will show coupling between modes
 - Diagonal peaks=Same as 1D spectrum
 - Off-diagonal peaks=Shows which modes are coupled to one another
 - Can arise from anharmonicity in the potential or nonlinear dependence of the transition dipole on the mode coordinates
 - Probe the coupling between the 42 cm⁻¹ and 49 cm⁻¹ libron modes in crystalline L-alanine.
 - Probe the coupling between low-frequency modes in proteins
- 3rd order process
 - Complementary to the 5th order Raman process



2-Dimensional (Nonlinear) THz Spectroscopy



Anharmonicity in intermolecular potential ...


$$V(Q) = \sum_{i=1}^s \frac{1}{2} k_i Q_i^2 + \sum_{i,j,k} \frac{g_{i,j,k}}{3!} Q_i Q_j Q_k + \dots$$

... or nonlinear dependence of transition dipole ...


$$\mu(Q) = \mu_0 + \sum_{i=1}^s \mu_i Q_i + \sum_{i,j} \frac{\mu_{i,j}}{2!} Q_i Q_j + \dots$$

Model System: Tryptophan Crystals


Intermolecular modes depend on enantiomers



Nonlinear two-dimensional THz spectroscopy




Previous work: 2D IR studies




1,3-cyclohexane

Hochstrasser



acetylproline- NH_2


Hochstrasser



acetyl-acetonato dicarbonyl rhodium(I)
(RDC)

Tokmakoff

L-alanine crystals




Anharmonicity in potential



$$V(Q) = \sum_{i=1}^s \frac{1}{2} k_i Q_i^2 + \sum_{i,j,k} \frac{g_{i,j,k}}{3!} Q_i Q_j Q_k + \dots$$

Nonlinearity in dipole

$$\mu(Q) = \mu_0 + \sum_{i=1}^s \mu_i Q_i + \sum_{i,j} \frac{\mu_{i,j}}{2!} Q_i Q_j + \dots$$



Simulated Results




Summary: 2D THz Spectroscopy

- Separate potential anharmonicity from dipole nonlinearity.
- Observe couplings between phonon modes.
- Observe FIR photon echo in liquids.

Shameless Self-Promotion

Lots of Recent Activity and Interest in THz Radiation




©Martin Koch 2002




Additional THz Information

Reasonably comprehensive review:

C. A. Schmuttenmaer, "Exploring Dynamics in the Far-Infrared with Terahertz Spectroscopy"
Chemical Reviews, **104**, 1759 (2004).



Report is available on DOE BES website.



Eric
Beaurepaire
(CNRS)

Gordon Turner Jim Schleicher
Shayne Harrel Tim French
Matt Beard Alan True