

DIS2023: XXX International Workshop on Deep-Inelastic Scattering and Related Subjects

Inclusive electron scattering off the proton with CLAS12 at JLab

V. Klimenko*, University of Connecticut
For the CLAS Collaboration

03/30/2023



* contact person
valerii.klimenko@uconn.edu

Extending Knowledge of the Nucleon PDF in the Resonance Region

- Global QCD analyses have provided detailed information on the nucleon PDFs in a wide range of parton fractional longitudinal momentum, x , from 10^{-4} to 0.9.
- At large x , in the nucleon resonance region $W < 2.5$ GeV, the PDFs are significantly less explored.
- Extractions in this region require accounting for higher twist effects, target-mass corrections and evaluation from the nucleon resonance electroexcitations.

A. N. Hiller Blin et al., *Phys. Rev. C* 100 (2019) 3, 035201, [hep-ph 1904.08016]


A. Accardi et al., *Phys. Rev. D.* 11, 114017 (2016), [hep-ph 1602.03154]




CLAS Results

- CLAS measured the inclusive cross section up to $x = 0.9$ and Q^2 from 0.25 to 4.5 GeV^2 .
- Owing to large acceptance of CLAS, the information on inclusive structure function F_2 can be obtained within a wide range of W from pion threshold to maximal kinematically allowed W -values in any given bin of Q^2 covered in the measurements.

M. Osipenko et al., *Phys. Rev. D* 67 (2003), 092001, [hep-ph 0301204]



World data used for moment evaluations of F_2 . Shaded area corresponds to CLAS.



Resonant Contributions

- CLAS results on $\gamma_p N^*$ electrocouplings for most N^* in the mass range $W < 1.8$ GeV allowed us to evaluate the resonant contributions to F_2 structure function from the experimental results on resonance electroexcitation amplitudes.
- Resonant contributions demonstrate pronounced evolution with photon virtuality Q^2 different in the first, the second and the third resonance regions.
- Information on Q^2 evolution $\gamma_p N^*$ electrocouplings for all prominent N^* is needed for realistic evaluation of the resonant contribution into inclusive electron scattering observables.


$$\sigma_{T,L}^R(W, Q^2) = \frac{\pi}{q_\gamma^2} \sum_R (2J_R + 1) \frac{M_R^2 \Gamma_R(W) \Gamma_{\gamma,R}^{T,L}(M_R, Q^2)}{(M_R^2 - W^2)^2 + (M_R \Gamma_R(W))^2}$$

Decay widths of resonance R to $\gamma^* p$ related to electrocouplings from previous slide.



$$\Gamma_{\gamma,R}^T(W = M_R, Q^2) = \frac{q_{\gamma,R}^2(Q^2)}{\pi} \frac{2M}{(2J_R + 1)M_R} \times \left(|A_{1/2}^R(Q^2)|^2 + |A_{3/2}^R(Q^2)|^2 \right),$$

$$\Gamma_{\gamma,R}^L(W = M_R, Q^2) = \frac{2q_{\gamma,R}^2(Q^2)}{\pi} \frac{2M}{(2J_R + 1)M_R} \times |S_{1/2}^R(Q^2)|^2,$$

A. N. Hiller Blin et al., *Phys. Rev. C* 104 (2021) 2, 025201, [hep-ph 2105.05834]



Inclusive Measurement



V. Burkert et al., Nucl. Instrum. Meth. A 959 (2020) 163419

- Measurements of $(e,e'X)$ inclusive cross sections are important to understand electron detection efficiency needed for evaluation of the cross sections of semi-inclusive and exclusive processes foreseen in the exploration with the CLAS12 detector
- CLAS12: $10^{35} \text{ cm}^{-2}\text{sec}^{-1}$ luminosity, nearly 4π acceptance, $0.05 \text{ GeV}^2 < Q^2 < 10.0 \text{ GeV}^2$ coverage over photon virtuality.
- Began data taking in Spring 2018 – many “run periods” now available.
- Data from Fall 2018 - 10.6 GeV electron beam, longitudinally polarized beam, liquid H_2 target.

Cross Section Calculation




$$\frac{d\sigma}{dQ^2 dW} = \frac{1}{\Delta Q^2 \Delta W} \cdot \frac{N}{\eta \cdot R \cdot BC \cdot N_0} \cdot \frac{1}{N_A \rho t / A_\omega} \quad (1)$$

- Q^2 - four-momentum transfer squared
- W - invariant mass of the final hadron system
- R - radiative correction factor
- BC - bin centering correction
- N - bin event yield
- η - is the product of geometrical acceptance and electron detection efficiency
- N_0 - live-time corrected incident electron flux summed over all data runs
- N_A - Avogadro's number
- ρ - target density
- t - target length
- A_ω - atomic weight of the target


Electron PID

Event yield extraction N in (1)

- Limited to Forward Detector (5 - 35° coverage in polar angle)
- Negative track with a hit in Time-of-Flight, Electromagnetic Calorimeters and High Threshold Cherenkov Counter (HTCC)
- >2.0 photoelectrons in HTCC
- **DC and PCAL Fiducial cuts.**
- **-8 < Vertex Z < 2 cm**
- **3.5- σ cuts on a parameterized momentum-dependent sampling fraction.**
- >70 MeV PCAL
- Electron/pion separation (triangular cut)



CLAS12 measurement




CLAS Data

Forward Calorimeter sampling fraction for electrons: $3.5\sigma \pm 0.5\sigma$

Simulation

Acceptance η in (1)

- Inclusive EG: M. Sargsyan, CLAS-NOTE 90-007 (1990).
- Elastic tail + Inelastic radiated.
- Background merging 45 nA.
- Kinematic range:
 - Theta range 5 - 40°.
 - Scattered electron momentum 1.4 – 11 GeV
 - Full Q^2 coverage.
- Additional kinematic smearing to match the resolution of reconstructed data.



Acceptance corrections

Acceptance η in (1)

Acceptance corrections can be done with multiple methods. One is bin-by-bin method and there are plenty of matrix methods:

- **Bin-by-bin:**

$$\text{Acceptance} = \frac{\# \text{ Events Reconstructed}}{\# \text{ Events Generated}}$$

CERN RooUnfold package was used:
<https://gitlab.cern.ch/RooUnfold/RooUnfold>

It is enough in case of “realistic” EG and “good” MC. It does not include connection between Gen. and Rec. event.


- **Acceptance Matrix:** $A_{(i,j)}$ describes both Acceptance (geometrical acceptance and detector efficiency) and Bin Migration:

$$A_{(i,j)} = \frac{\# \text{ Events Generated in bin } j \text{ but Reconstructed in bin } i}{\text{Total number of Events Generated in the } j\text{th bin}}$$

Acceptance unfolding: $Y_i = A_{(i,j)} X_j \Rightarrow X_j = A^{-1}_{(i,j)} Y_i$ where Y_i number of measured events in i -th bin, X_j is number of acceptance corrected events in j -th bin

We used:


1. Bin-by-bin method.
2. SVD Matrix Unfolding method.
3. Bayesian Matrix 2D unfolding is 1D - 3D Matrix method



Acceptance corrected W-yield in all Q^2 bins

Acceptance η in (1)

Red - 2D Bayesian method
Black - Bin-by-bin method



Radiative Corrections


$$R_{in(1)}$$

Each (Q^2, W) bin was divided into 21x11 sub bins. Cross Sections with rad. effects on and off were calculated in every sub bin.


Radiative Correction factor:

$$\frac{\text{Mean Cross Section (Rad)}}{\text{Mean Cross Section (No Rad)}}$$


Inclusive with radiative effects




Elastic with radiative effects




$2.55 < Q^2 < 2.99 \text{ GeV}^2$



$2.99 < Q^2 < 3.49 \text{ GeV}^2$



$3.49 < Q^2 < 4.08 \text{ GeV}^2$




Bin Centering Corrections

BC in (1)

$$\frac{d\sigma}{dQ^2 dW} = \frac{1}{\Delta Q^2 \Delta W} \cdot \frac{N}{\eta \cdot R \cdot BC \cdot N_0} \cdot \frac{1}{N_A \rho t / A_\omega}$$



Each (Q^2, W) bin was divided into (the same) 21x11 sub bins.

$$BC \text{ Corrections (BCC)} = \frac{\text{Cross Section (No Rad) in the central point}}{\text{Mean Cross Section (No Rad)}}$$



Iterations

After applying all the corrections and normalization accordingly to faraday cup charge we obtained preliminary cross section. That cross sections can be used as a base for new event generator and as a new model for RC and BC estimation.




Cross Section with three models:

- Green** – Cross Sections with initial model
- Orange** - Cross Sections with iteration 1 model
- Purple** - Cross Sections with iteration 2 model

There is small peak shift. Peaks become narrower with iterations number.

Preliminary Cross Section

- Preliminary CLAS12 measurements.
- CLAS data (after interpolation into the grid of our experiment), Phys. Rev. D67, 092001 (2003).



Summary

- Preliminary results on inclusive electron scattering cross sections are available from CLAS12 in the kinematic range of $1.15 < W < 2.5$ GeV and $2.55 < Q^2 < 9.0$ GeV². Our new measurements show reasonable agreements with world data in overlapping Q^2 regions.
- First (e,e'X) data from CLAS12 have become available within a broad coverage over W from pion threshold to 2.5 GeV at any given bin of Q^2 within the range of photon virtuality from 2.55 GeV² to 9.0 GeV².
- Evaluation of the resonant contributions from exclusive meson electroproduction data will pave a way to extend knowledge on PDF at large x in the resonance region.
- The (e,e'X) data from CLAS12 offer an opportunity to explore evolution of inclusive structure function F_2 within the range of distances where the transition from strongly coupled to pQCD regimes is anticipated.