

## Charmonium properties in a renormalization scheme

J. Segovia, D. R. Entem, F. Fernández and E. Ruiz-Arriola  
[segonza@usal.es](mailto:segonza@usal.es)

12th International Conference on Meson-Nucleon Physics and the Structure  
of the Nucleon (MENU 2010)  
Williamsburg, May - June 2010



University of Granada



University of Salamanca

# Contents

- 1 Constituent quark model
- 2 Renormalization scheme with boundary conditions
- 3 Masses in charmonium from both schemes
- 4 Study of some physical observables in function of different model parameters
  - Masses vs  $\alpha_s$
  - Masses vs  $\sigma$
  - Leptonic decays vs  $\alpha_s$
- 5 Conclusions

# 1.- Constituent quark model

1.1.- Most important ingredients / J. Phys. G: Nucl. Part. Phys. **31**, 481 (2005)

- Spontaneous chiral symmetry breaking (Goldstone-Bosons exchange):

$$L = \bar{\psi} (i\gamma^\mu \partial_\mu - MU^{\gamma_5}) \psi \rightarrow U^{\gamma_5} = 1 + \frac{i}{f_\pi} \gamma^5 \lambda^a \pi^a - \frac{1}{2f_\pi^2} \pi^a \pi^a + \dots$$

$$M(q^2) = m_q F(q^2) = m_q \left[ \frac{\Lambda^2}{\Lambda^2 + q^2} \right]^{1/2}$$

- QCD perturbative effects (One gluon exchange):

$$L = i\sqrt{4\pi\alpha_s} \bar{\psi} \gamma_\mu G^\mu \lambda^c \psi$$

- Confinement (screened potential):

$$V_{CON}^C(\vec{r}_{ij}) = [-a_c(1 - e^{-\mu_c r_{ij}}) + \Delta] (\vec{\lambda}_i^c \cdot \vec{\lambda}_j^c)$$

$$\begin{cases} V_{CON}^C(\vec{r}_{ij}) = (-a_c \mu_c r_{ij} + \Delta) (\vec{\lambda}_i^c \cdot \vec{\lambda}_j^c) & r_{ij} \rightarrow 0 \\ V_{CON}^C(\vec{r}_{ij}) = (-a_c + \Delta) (\vec{\lambda}_i^c \cdot \vec{\lambda}_j^c) & r_{ij} \rightarrow \infty \end{cases}$$

*Remember:*  $\sigma = -a_c \mu_c (\vec{\lambda}_i^c \cdot \vec{\lambda}_j^c)$



## 1.2.- Non-relativistic reduction of our potential

- One-gluon exchange (OGE)

$$V_{OGE}^C(\vec{r}_{ij}) = \frac{1}{4} \alpha_s (\vec{\lambda}_i^c \cdot \vec{\lambda}_j^c) \left[ \frac{1}{r_{ij}} - \frac{1}{6m_i m_j} (\vec{\sigma}_i \cdot \vec{\sigma}_j) \frac{e^{-r_{ij}/r_0(\mu)}}{r_{ij} r_0^2(\mu)} \right]$$

$$V_{OGE}^T(\vec{r}_{ij}) = -\frac{1}{16} \frac{\alpha_s}{m_i m_j} (\vec{\lambda}_i^c \cdot \vec{\lambda}_j^c) \left[ \frac{1}{r_{ij}^3} - \frac{e^{-r_{ij}/r_g(\mu)}}{r_{ij}} \left( \frac{1}{r_{ij}^2} + \frac{1}{3r_g^2(\mu)} + \frac{1}{r_{ij} r_g(\mu)} \right) \right] S_{ij}$$

$$\begin{aligned} V_{OGE}^{SO}(\vec{r}_{ij}) = & -\frac{1}{16} \frac{\alpha_s}{m_i^2 m_j^2} (\vec{\lambda}_i^c \cdot \vec{\lambda}_j^c) \left[ \frac{1}{r_{ij}^3} - \frac{e^{-r_{ij}/r_g(\mu)}}{r_{ij}^3} \left( 1 + \frac{r_{ij}}{r_g(\mu)} \right) \right] \times \\ & \times \left[ ((m_i + m_j)^2 + 2m_i m_j)(\vec{S}_+ \cdot \vec{L}) + (m_j^2 - m_i^2)(\vec{S}_- \cdot \vec{L}) \right] \end{aligned}$$

- Confinement

$$V_{CON}^C(\vec{r}_{ij}) = [-a_c(1 - e^{-\mu_c r_{ij}}) + \Delta] (\vec{\lambda}_i^c \cdot \vec{\lambda}_j^c)$$

$$V_{CON}^{SO}(\vec{r}_{ij}) = -(\vec{\lambda}_i^c \cdot \vec{\lambda}_j^c) \frac{a_c \mu_c e^{-\mu_c r_{ij}}}{4m_i^2 m_j^2 r_{ij}} \left[ ((m_i^2 + m_j^2)(1 - 2a_s) \right.$$

$$\left. + 4m_i m_j (1 - a_s))(\vec{S}_+ \cdot \vec{L}) + (m_j^2 - m_i^2)(1 - 2a_s)(\vec{S}_- \cdot \vec{L}) \right]$$

## 1.2.- Non-relativistic reduction of our potential. Singular contributions

$$V_{OGE}^C(\vec{r}_{ij}) = \frac{1}{4}\alpha_s(\vec{\lambda}_i^c \cdot \vec{\lambda}_j^c) \left[ \frac{1}{r_{ij}} - \frac{1}{6m_i m_j} (\vec{\sigma}_i \cdot \vec{\sigma}_j) 4\pi \delta(r_{ij}) \right]$$

$$V_{OGE}^T(\vec{r}_{ij}) = -\frac{1}{16} \frac{\alpha_s}{m_i m_j} (\vec{\lambda}_i^c \cdot \vec{\lambda}_j^c) \frac{1}{r_{ij}^3} S_{ij}$$

$$\begin{aligned} V_{OGE}^{SO}(\vec{r}_{ij}) = & -\frac{1}{16} \frac{\alpha_s}{m_i^2 m_j^2} (\vec{\lambda}_i^c \cdot \vec{\lambda}_j^c) \frac{1}{r_{ij}^3} \times \\ & \times \left[ ((m_i + m_j)^2 + 2m_i m_j)(\vec{S}_+ \cdot \vec{L}) + (m_j^2 - m_i^2)(\vec{S}_- \cdot \vec{L}) \right] \end{aligned}$$

$$V_{OGE}^C(\vec{r}_{ij}) = \frac{1}{4}\alpha_s(\vec{\lambda}_i^c \cdot \vec{\lambda}_j^c) \left[ \frac{1}{r_{ij}} - \frac{1}{6m_i m_j} (\vec{\sigma}_i \cdot \vec{\sigma}_j) \frac{e^{-r_{ij}/r_0(\mu)}}{r_{ij} r_0^2(\mu)} \right]$$

$$V_{OGE}^T(\vec{r}_{ij}) = -\frac{1}{16} \frac{\alpha_s}{m_i m_j} (\vec{\lambda}_i^c \cdot \vec{\lambda}_j^c) \left[ \frac{1}{r_{ij}^3} - \frac{e^{-r_{ij}/r_g(\mu)}}{r_{ij}} \left( \frac{1}{r_{ij}^2} + \frac{1}{3r_g^2(\mu)} + \frac{1}{r_{ij} r_g(\mu)} \right) \right] S_{ij}$$

$$\begin{aligned} V_{OGE}^{SO}(\vec{r}_{ij}) = & -\frac{1}{16} \frac{\alpha_s}{m_i^2 m_j^2} (\vec{\lambda}_i^c \cdot \vec{\lambda}_j^c) \left[ \frac{1}{r_{ij}^3} - \frac{e^{-r_{ij}/r_g(\mu)}}{r_{ij}^3} \left( 1 + \frac{r_{ij}}{r_g(\mu)} \right) \right] \times \\ & \times \left[ ((m_i + m_j)^2 + 2m_i m_j)(\vec{S}_+ \cdot \vec{L}) + (m_j^2 - m_i^2)(\vec{S}_- \cdot \vec{L}) \right] \end{aligned}$$



## 1.3.- Some recent applications

### • N-N interaction

- D. R. Entem, F. Fernández and A. Valcarce, Phys. Rev. C **62**, 034002 (2000)
- B. Julia-Díaz, J. Haidenbauer, A. Valcarce and F. Fernández, Phys. Rev. C **65**, 034001 (2002)

### • Baryon spectrum

- H. Garcilazo, A. Valcarce and F. Fernández, Phys. Rev. C **63**, 035207 (2001)
- H. Garcilazo, A. Valcarce and F. Fernández, Phys. Rev. C **64**, 058201 (2001)

### • Meson spectrum

- J. Vijande, A. Valcarce and F. Fernández, J. Phys. G **31**, 481 (2005)
- J. Segovia, D. R. Entem and F. Fernández, Phys. Rev. D **78** 114033 (2008)
- J. Segovia, D. R. Entem and F. Fernández, accepted by J. Phys. G

### • Molecular states

- P. G. Ortega, J. Segovia, D. R. Entem and F. Fernández, Phys. Rev. D **81**, 054023 (2010)

## 2.- Renormalization scheme with boundary conditions

### 2.1.- Features

- We have two particles interacting through a central potential

$$-u''(r) + \mathcal{U}(r)u(r) = k^2 u(r)$$

*Second order differential equation  $\Rightarrow$  two linear independent solutions*

- One of them is the physical solution and so we have to choose the solution that is given by the regular condition at the origin

$$u(0) = 0$$

- BUT there are a great number of physical systems which we do not know the exact potential



*Interaction potential is known at large distances but not at short distances*

- Solution: We can apply boundary conditions to potentials whose short range is not known

*Means that we can fix one physical observable of the large distances and it is equivalent to impose a boundary condition at origin*



## 2.2.- Application to hydrogen Spin-Orbit



$$V(r) = -\frac{\alpha \hbar c}{r} + \frac{\alpha (\hbar c)^3}{2m^2} \frac{1}{r^3} \vec{L} \cdot \vec{S}$$

$n = 2 \ J = 1/2 \ L = 1 \ S = 1/2 \Rightarrow \text{ATTRACTIVE}$

## 2.3.- Some recent applications and our aim

- 'Renormalization of NN-Scattering with one pion exchange and boundary conditions'  
M. Pavon Valderrama and E. Ruiz Arriola, Phys. Rev. C **70** 044006 (2004)
- 'Renormalization of the deuteron with one pion exchange'  
M. Pavon Valderrama and E. Ruiz Arriola, Phys. Rev. C **72** 054002 (2005)
- 'Renormalization vs strong form factors for one boson exchange'  
A. Calle Cordon and E. Ruiz Arriola, Phys. Rev. C **81**, 044002 (2010)
- 'Low energy universality and scaling of Van der Waals forces'  
A. Calle Cordon and E. Ruiz Arriola, Phys. Rev. A **81**, 044701 (2010)

### OUR AIM

- *Renormalization with boundary conditions applied to our model  $\Rightarrow$  Non perturbative treatment of our singular potential parts without cut-offs (no  $\hat{r}_0$  and  $\hat{r}_g$ )*
- *The model in this framework has few parameters and allows us to study the correlations between physical observables and model parameters which have some physical meaning*



## 2.4.- Final expresions for the different model contributions

- One-gluon exchange (OGE)

$$V_{OGE}^C(\vec{r}_{ij}) = \frac{1}{4} \alpha_s (\vec{\lambda}_i^c \cdot \vec{\lambda}_j^c) \frac{1}{r_{ij}}$$

$$V_{OGE}^T(\vec{r}_{ij}) = -\frac{1}{16} \frac{\alpha_s}{m_i m_j} (\vec{\lambda}_i^c \cdot \vec{\lambda}_j^c) \frac{1}{r_{ij}^3} S_{ij}$$

$$\begin{aligned} V_{OGE}^{SO}(\vec{r}_{ij}) = & -\frac{1}{16} \frac{\alpha_s}{m_i^2 m_j^2} (\vec{\lambda}_i^c \cdot \vec{\lambda}_j^c) \frac{1}{r_{ij}^3} \times \\ & \times \left[ ((m_i + m_j)^2 + 2m_i m_j)(\vec{S}_+ \cdot \vec{L}) + (m_j^2 - m_i^2)(\vec{S}_- \cdot \vec{L}) \right] \end{aligned}$$

- Confinement

$$V_{CON}^C(\vec{r}_{ij}) = [-a_c(1 - e^{-\mu_c r_{ij}}) + \Delta] (\vec{\lambda}_i^c \cdot \vec{\lambda}_j^c)$$

$$\begin{aligned} V_{CON}^{SO}(\vec{r}_{ij}) = & - \left( \vec{\lambda}_i^c \cdot \vec{\lambda}_j^c \right) \frac{a_c \mu_c e^{-\mu_c r_{ij}}}{4m_i^2 m_j^2 r_{ij}} \left[ ((m_i^2 + m_j^2)(1 - 2a_s) \right. \\ & \left. + 4m_i m_j(1 - a_s))(\vec{S}_+ \cdot \vec{L}) + (m_j^2 - m_i^2)(1 - 2a_s)(\vec{S}_- \cdot \vec{L}) \right] \end{aligned}$$

### 3.- Masses in charmonium from both schemes

#### 3.1 Model parameters

Quark Mass	$m_c$ (MeV)	1763
OGE	$\alpha_0$	2.118
	$\Lambda_0$ (fm $^{-1}$ )	0.113
	$\mu_0$ (MeV)	36.976
	$\hat{r}_0$ (fm)	0.181
	$\hat{r}_g$ (fm)	0.259
Confinement	$a_c$ (MeV)	507.4
	$\mu_c$ (fm $^{-1}$ )	0.576
	$\Delta$ (MeV)	184.432
	$a_s$	0.81



*Remember: There are no cut-off in the renormalization scheme (no  $\hat{r}_0$  and  $\hat{r}_g$ )*

### 3.2.- Calculation

State	CQM (MeV)	$\mathcal{P}_{^3S_1}$	$\mathcal{P}_{^3D_1}$
$J/\psi$	3096	99.959	0.041
$\psi(2S)$	3703	99.958	0.042
$\psi(3770)$	3796	0.032	99.968
$\psi(4040)$	4097	99.935	0.065
$\psi(4160)$	4153	0.060	99.940
$\psi(4360)$	4389	99.908	0.092
$\psi(4415)$	4426	0.089	99.911
$\psi(4660)$	4614	99.884	0.116
$\psi(4660)$	4641	0.114	99.886

*The mixing between S-wave and D-wave states is negligible from our original model and so we can calculate  ${}^3S_1$  and  ${}^3D_1$  without coupling in a renormalization scheme*

### 3.3.- Solution stability in renormalization scheme







### 3.4.- Model results

State	n	$M_{RSC}$ (MeV)	$M_{CQM}$ (MeV)	$M_{exp}$ (MeV)
$^3S_1$	1	3096 <sup>†</sup>	3096	$3096.916 \pm 0.011$
	2	3703	3703	$3686.093 \pm 0.034$
	3	4097	4097	$4039.6 \pm 4.3$
	4	4389	4389	-
	5	4614	4614	-
$^3D_1$	1	3796 <sup>†</sup>	3796	$3772.92 \pm 0.35$
	2	4153	4153	$4153 \pm 3$
	3	4426	4426	$4421 \pm 4$
	4	4641	4641	-


*Both schemes are equivalent*

## 4.- Study of some physical observables in function of different parameters





### 4.1.- Masses vs $\alpha_s$







## 4.1.- Masses vs $\alpha_s$ . Continuation




## 4.1.- Masses vs $\alpha_s$ . Continuation







#### 4.3.- Masses vs $\sigma$




## 4.3.- Masses vs $\sigma$ . Continuation



## 4.3.- Masses vs $\sigma$ . Continuation







## 4.4.- Leptonic widths vs $\alpha_s$




$$\Gamma(n^3S_1 \rightarrow e^+ e^-) = \frac{4\alpha^2 e_c^2 |R_n(0)|^2}{E_n^2} \left(1 - \frac{16\alpha_s}{3\pi}\right)$$

$$\mathcal{R} = \frac{\Gamma(n^3S_1 \rightarrow e^+ e^-)}{\Gamma(1^3S_1 \rightarrow e^+ e^-)} = \frac{|R_n(0)|^2}{|R_1(0)|^2} \frac{E_1^2}{E_n^2}$$

## 4.4.- Leptonic widths vs $\alpha_s$ . Continuation



#### 4.4.- Leptonic widths vs $\alpha_s$ . Continuation



## 5.- Conclusions

- We re-analyze the calculation of the charmonium spectrum in constituent quark model using a renormalization boundary condition scheme
- We find a good agreement between both schemes which provides confidence on the way the original model take into account the unknown short distance dynamics
- The use of this scheme allows us to further study the dependence of the states on the model parameters in a cleaner way since the regulator dependence has been removed when a suitable renormalization condition is imposed
- We find:
  - The mass of the excited states strongly depend on the string tension parameter
  - There is a remarkable insensitivity to the strong coupling constant entering the one gluon exchange contribution to the potential. This avoids a great deal of unphysical fine tuning which suggested taking for this parameter unnaturally large values  $\alpha_s \sim 0.3 - 0.4$
  - The leptonic widths depend strongly on the strong coupling constant. As expected because is a short range observable