

The Physics Opportunities with a K^0_L Facility at



Moskov Amaryan



YSTAR2016, JLab, November 16, 2016

A Letter of Intent to Jefferson Lab PAC-43.

Physics Opportunities with a Secondary K_L^0 Beam at JLab.

Moskov J. Amaryan (spokesperson),^{1,*} Yakov I. Azimov,² William J. Briscoe,³ Eugene Chudakov,⁴
Pavel Degtyarenko,⁴ Gail Dodge,¹ Michael Döring,³ Helmut Haberzettl,³ Charles E. Hyde,¹ Benjamin C. Jackson,⁵
Christopher D. Keith,⁴ Ilya Larin,¹ Dave J. Mack,⁴ D. Mark Manley,⁶ Kanzo Nakayama,⁵ Yongseok Oh,⁷
Emilie Passemar,⁸ Diane Schott,³ Alexander Somov,⁴ Igor Strakovsky,³ and Ronald Workman³

¹*Old Dominion University, Norfolk, VA 23529*

²*Petersburg Nuclear Physics Institute, Gatchina, St. Petersburg 188300, Russia*

³*The George Washington University, Washington, DC 20052*

⁴*Thomas Jefferson National Accelerator Facility, Newport News, Virginia 23606*

⁵*The University of Georgia, Athens, GA 30602*

⁶*Kent State University, Kent, OH 44242*

⁷*Kyungpook National University, Daegu 702-701, Korea*

⁸*Indiana University, Bloomington, IN 47405*

(Dated: May 15, 2015)

Outline

- Introduction
- Baryon Multiplets
- Reactions with K_L^0 beam on proton target
- Experimental Arrangement
- K_L^0 Beam at GlueX
- Expected rates
- Summary

The nonexistent is whatever we
have not sufficiently desired.


Franz Kafka

(In some cases it may be true M.A.)


Constituent Quark Model

SU(3)

Octet



Decuplet





But there are many more states predicted, where are they?

Where are hybrids, glueballs, multiquark states ?



Well, some of them may already have been observed?

Lattice QCD calculations

Thick borders: Hybrid states




Low Lying states



Lattice QCD calculations

Thick borders: Hybrid states




Low Lying states


Edwards, Mathur, Richards and Wallace
Phys. Rev. D 87, 054506 (2013)

Status of $[I]^*$

well known




Status of Ω^{-*}



only one well known state?

- **Three light quarks** can be arranged in **6** baryonic families, \mathbf{N}^* , $\mathbf{\Delta}^*$, $\mathbf{\Lambda}^*$, $\mathbf{\Sigma}^*$, $\mathbf{\Xi}^*$, & $\mathbf{\Omega}^*$.
- **Number of members** in a family that can exist **is not arbitrary**.
- If $\mathbf{SU(3)}_F$ symmetry of **QCD** is controlling, then:

Octet: \mathbf{N}^* , $\mathbf{\Lambda}^*$, $\mathbf{\Sigma}^*$, $\mathbf{\Xi}^*$
 Decuplet: $\mathbf{\Delta}^*$, $\mathbf{\Sigma}^*$, $\mathbf{\Xi}^*$, & $\mathbf{\Omega}^*$

- Number of experimentally identified resonances of each baryon family in  summary tables is **17** \mathbf{N}^* , **24** $\mathbf{\Delta}^*$, **14** $\mathbf{\Lambda}^*$, **12** $\mathbf{\Sigma}^*$, **7** $\mathbf{\Xi}^*$, & **2** $\mathbf{\Omega}^*$.
- **Constituent Quark** models, for instance, predict existence of no less than **64** \mathbf{N}^* , **22** $\mathbf{\Delta}^*$ states with **mass** < **3** GeV.
- Seriousness of “**missing-states**” problem is obvious from these numbers.
- To complete $\mathbf{SU(3)}_F$ multiplets, one needs no less than **17** $\mathbf{\Lambda}^*$, **41** $\mathbf{\Sigma}^*$, **41** $\mathbf{\Xi}^*$, & **24** $\mathbf{\Omega}^*$.

Recourse to the Neutral Kaon System

Strangeness eigenstates with $J^{PC} = 0^{-+}$

$$|K^0\rangle = |d\bar{s}\rangle, \quad |\bar{K}^0\rangle = |\bar{d}s\rangle$$

$$S=+1$$

$$S=-1$$

Parity eigenstates with intrinsic $P = -1$

$$P|K^0\rangle = -|K^0\rangle, \quad P|\bar{K}^0\rangle = -|\bar{K}^0\rangle$$

Effect of C-Parity can be taken to be

$$C|K^0\rangle = |\bar{K}^0\rangle, \quad C|\bar{K}^0\rangle = |K^0\rangle$$



However not CP eigenstates

$$CP|K^0\rangle = -|\bar{K}^0\rangle, \quad CP|\bar{K}^0\rangle = -|K^0\rangle$$

CP eigenstates can be formed

$$|K_1\rangle \equiv \frac{1}{\sqrt{2}} (|K^0\rangle - |\bar{K}^0\rangle) ; \quad \underline{CP |K_1\rangle = + |K_1\rangle}$$

$$|K_2\rangle \equiv \frac{1}{\sqrt{2}} (|K^0\rangle + |\bar{K}^0\rangle) ; \quad \underline{CP |K_2\rangle = - |K_2\rangle}$$



$$K^0 \quad \text{and} \quad \bar{K}^0$$

are unstable particles decaying via W

$$K_S (K - \text{short}) \quad \text{and} \quad K_L (K - \text{long})$$

propagate as free particles and have distinct lifetimes

$$\underline{\tau_S = 0.9 \times 10^{-10} \text{ s}} \quad \text{and} \quad \underline{\tau_L = 0.5 \times 10^{-7} \text{ s}} \quad (c\tau = 15 \text{ m})$$

$$|K_S\rangle \equiv \frac{1}{\sqrt{1 + |\epsilon|^2}} (|K_1\rangle + \epsilon |K_2\rangle) \approx |K_1\rangle$$

$$|K_L\rangle \equiv \frac{1}{\sqrt{1 + |\epsilon|^2}} (|K_2\rangle + \epsilon |K_1\rangle) \approx |K_2\rangle$$

$$|\epsilon| \approx 2.3 \times 10^{-3}$$

defines the level of CP violation

CP conserving decays

$$\begin{array}{ll} K_S \rightarrow \pi^+ \pi^- & \text{BR} = 68.6\% \\ \rightarrow \pi^0 \pi^0 & \text{BR} = 31.4\% \end{array}$$

$$\begin{array}{ll} K_L \rightarrow \pi^+ \pi^- \pi^0 & \text{BR} = 12.6\% \\ \rightarrow \pi^0 \pi^0 \pi^0 & \text{BR} = 21.1\% \\ \rightarrow \pi^- e^+ \nu_e & \text{BR} = 19.4\% \\ \rightarrow \pi^+ e^- \bar{\nu}_e & \text{BR} = 19.4\% \\ \rightarrow \pi^- \mu^+ \nu_\mu & \text{BR} = 13.6\% \\ \rightarrow \pi^+ \mu^- \bar{\nu}_\mu & \text{BR} = 13.6\% \end{array}$$

CP violating decays observed in 1964

$$\begin{array}{ll} K_L \rightarrow \pi^+ \pi^- & \text{BR} = 2.1 \times 10^{-3} \\ \rightarrow \pi^0 \pi^0 & \text{BR} = 9.4 \times 10^{-4} \end{array}$$

What can be learned with a K_L^0 beam ?

List of reactions:

Elastic and charge-exchange

$$\begin{aligned} K_L^0 p &\rightarrow K_S^0 p \\ K_L^0 p &\rightarrow K^+ n \end{aligned}$$

Two-body with $S=-1$

$$\begin{aligned} K_L^0 p &\rightarrow \pi^+ \Lambda \\ K_L^0 p &\rightarrow \pi^+ \Sigma^0 \end{aligned}$$

Two-body with $S=-2$

$$\begin{aligned} K_L^0 p &\rightarrow K^+ \Xi^0 \\ K_L^0 p &\rightarrow K^+ \Xi^{0*} \end{aligned}$$

Three-body with $S=-2$

$$\begin{aligned} K_L^0 p &\rightarrow \pi^+ K^+ \Xi^- \\ K_L^0 p &\rightarrow \pi^+ K^+ \Xi^{*-} \end{aligned}$$

Three-body with $S=-3$





$$\begin{aligned} K_L^0 p &\rightarrow K^+ K^+ \Omega^- \\ K_L^0 p &\rightarrow K^+ K^+ \Omega^{*-} \end{aligned}$$

Very Limited World Data with K_L beam

(Mainly low stat. bubble chamber data. Compilation by I. Strakovsky)

blue points: $d\sigma/d\Omega$

red points: Polarization



we are not aware of any data on Neutron target


Expected Cross Sections vs Bubble Chamber Data

- **GlueX** measurements will span $\cos\theta$ from -0.95 to 0.95 in c.m. above $W = 1490$ MeV.


- K_L rate is 10^5 K_L /s.
- Uncertainties correspond to **100** days of running time.
- Cross section uncertainty estimates (statistics only) for

Courtesy of Simon Taylor, KL2016
Mark Manley, KL2016

$K_L p \rightarrow K_S p$





Expected
GlueX Data



BC Data

$K_L p \rightarrow \pi^+ \Lambda$



More details in KL2016 Workshop Proceedings


arXiv: 1604.02141

How to make a kaon beam?

Thomas Jefferson National Accelerator Facility




Aerial View




Hall D Beamline

Current setup




Hall D Tagger Area



- Design beam current limits: **5 μ A (60 kW) max**
- Design radiator thickness: **~ 0.0005 Radiation Lengths max**
- **Challenge:** Increase radiator thickness to **0.05-0.10 R.L.?!**

GEANT3 Model, 2000 electrons at 12 GeV

Compact γ Source




Compact Photon Source Concept

- **Strong magnet** after radiator deflects exiting electrons
- **Long-bore collimator** lets photon beam through
- **Electron beam dump** placed next to the collimator
- **Water-cooled Copper core** for better heat dissipation
- **Hermetic shielding** all around and close to the source
- **High Z and high density material** for bulk shielding
- **Borated Poly outer layer** for slowing, thermalizing, and absorbing fast neutrons still exiting the bulk shielding
- No need in tagging photons, so the design could be **compact**, as opposed to the Tagger Magnet concept


CPS: PR12-15-003 Proposal at JLab

Application example: CPS concept for new experiment in Hall A




MC simulation and direct calculations show acceptable background rates on SBS and NPS.


CPS at the Hall D Tagger Area




CPS, horizontal plane (1)



CPS, 50 electrons at 12 GeV



Dose Rate Evaluation and Comparison



- The dose rates in the Tagger vault for the **CPS** setup with 10% R.L. radiator are close to Standard XD ops
- The radiation spectral composition is different; most of the contribution in the **CPS** setup is from higher energy neutrons

Dose Rate Evaluation and Comparison

- The plots show comparison of dose rate estimates in the Tagger Area in two conditions: (1) **nominal Hall D operation** with the standard amorphous radiator at 0.0005 R.L., - with (2) radiator at 0.1 R.L., used as part of the **Compact Photon Source setup**.
- The comparison indicates that at equal beam currents, gamma radiation dose rates are much smaller for the CPS run (**~order of magnitude**), and neutron dose rates in the area are comparable.
- Design and shielding **optimization** may improve the comparison further in favor of the **CPS** solution

K^0_L beam (continued)

- Electron beam with $I_e = 5\mu A$
- Delivered with 60ns bunch spacing avoids overlap in the range of $P=0.35-10.0$ GeV/c
- Momentum measured with TOF
- K^0_L flux measured with pair spectrometer
- Side remark: Physics case with polarized targets is under study and feasible*

Detailed Design and Cost Estimate


- We do not see show-stoppers for implementation of the **CPS** concept in the experiment.
- 60 kW Copper-core dump will have characteristics close to the one installed already
- To make long and narrow photon beam collimation we propose to build the core using two symmetric flat plates, left and right, and make matching grooves in them for the beam entry cones, beam line, and the aperture collimator
- Cost would include detailed iterative modeling and simulation to optimize operation parameters, design, engineering and production, plus the choice and cost of bulk shielding material
- Crude cost expectation: within **\$0.5M**

Conclusions

- Compared to the alternative, the proposed **CPS** solution presents several advantages, including much less disturbance of the available infrastructure at the Tagger Area, and better flexibility in achieving high-intensity photon beam delivery to the Hall D
- The proposed **CPS** solution will satisfy proposed K^0_L beam production parameters
- We do not envision big technical or organizational difficulties in the implementation of the conceptual design


K_L -beam line

4



Rate of neutrons and K_L^0 on GlueX target

- JLAB



- PRL22.996 (1969) Brody et al.




FIG. 2. Comparison of the neutron and K_2^0 fluxes at the hydrogen bubble chamber for 2° production with 16-GeV electrons.

- With a proton beam ratio $n/K_L = 10^3-10^4$

K_L^0 beam

- **Electron beam** $E_e = 12GeV; I_e = 5\mu A$

- **Radiator (rad. length)**

10%

- **Be target (R=3cm)**

$L = 40cm$

- **LH2 target(L=30cm)**

$R = 3cm$

- **Distance Be-LH2**



16m


- **K_L Rate/sec**

$\sim 10^4$

Neutron Background

Neutron calculations for the KLF Project using MCMP6





Results:


Tally #1: 3200 n/(s cm²)

Tally #2: 40 n/(s cm²)

Tally #3: 140 n/(s cm²)

Tally #4: 3 n/(s cm²)

Neutron Flux $10e+10/4\pi/s$




- **Conclusion: Neutron Flux in Hall D is tolerable**

- Talk by Onishi at KL2016


J-PARC

Japan Proton Accelerator Research Complex



Two beam lines are under operation

K1.1 & High-p beam lines are under construction





- **ProjectX (Fermi Lab) arXiv:1306.5009**

Table III-2: Comparison of the K_L production yield. The BNL AGS kaon and neutron yields are taken from RSVP reviews in 2004 and 2005. The *Project X* yields are for a thick target, fully simulated with LAQGSM/MARS15 into the KOPIO beam solid angle and momentum acceptance.


	Beam energy	Target (λ_I)	$p(K)$ (MeV/ c)	K_L/s into 500 μ sr	$K_L : n$ ($E_n > 10$ MeV)
BNL AGS	24 GeV	1.1 Pt	300-1200	60×10^6	$\sim 1 : 1000$
<i>Project X</i>	3 GeV	1.0 C	300-1200	450×10^6	$\sim 1 : 2700$

*KL beam can be used to study rare decays
However it will be impossible to use it for hyperon spectroscopy
because of momentum range and n/K Ratio*



Momentum and W Resolution



W-Resolution




World Data on Ξ



Simulated with GlueX
 10^4 K_L /sec, one day of running




Jackson, Oh, Haberzettl, Nakayama
 Phys. Rev. C 91, 065208 (2015)



Status of $[I]^*$


Very poorly
measured at
AGS (BNL)
32 years ago

- C.M. Jenkins et al., Phys. Rev. Lett. 51, 951 (1983)




Cross Sections

$$K^- p \rightarrow \Xi^- X$$



$$K^- p \rightarrow \Omega^- X$$



J.K. Hassal et al., NPB 189 (1981)

Expected rates

<i>Production</i>	<i>J-PARC*</i>	<i>Jlab (this proposal)</i>
<i>flux/s</i>	$3 \times 10^4 K^-$	$10^4 K_L^0$
$\Xi^*/month$	3×10^5	2×10^5
$\Omega^{*-}/month$	600	4000

* *H.~Takahashi, NP A 914, 553 (2013)*


M.~Naruki and K.~Shirotori, LOI-2014-JPARC

Missing states and freezeout in heavy ion collisions

Close to T_c relaxation rates become small compared to the expansion rates and the system created in heavy ion collisions freezes out

The freeze-out is characterized by: (T^f, μ_B^f, μ_S^f) and hadron abundancies can be calculated from HRG

Lattice QCD Calculations



$$dU = TdS - PdV + \sum_{i=1}^n \mu_i dN_i$$

chemical potential

$$\mu_i = \frac{\partial U_i}{\partial N_i}$$

Bazavov et al., PRL 113(2014) 072001

12 GeV Approved Experiments by PAC Days

Topic	Hall A	Hall B	Hall C	Hall D	Other	Total
The Hadron spectra as probes of QCD		119		540		659
The transverse structure of the hadrons	145.5	85	102	25		357.5
The longitudinal structure of the hadrons	65	230	165			460
The 3D structure of the hadrons	409	872	212			1493
Hadrons and cold nuclear matter	180	175	201		14	570
Low-energy tests of the Standard Model and Fundamental Symmetries	547	180		79	60	866
Total Days	1346.5	1661	680	644	74	4405.5
Total Days – Without MIE Days	697.5	1661	680	644	28	3710.5
Total Approved Run Group Days (includes MIE)	1346.5	826	637	424	74	3307.5
Total Approved Run Group Days (without MIE)	528.5	826	637	424	28	2443.5
Total Days Completed	20	15	0	25	0	60
Total Days Remaining	508.5	811	637	399	28	2383.5

60 weeks

- **Bob McKeown's talk at 2016 UG meeting**

JLab Operations Budget ONP Briefing

- During FY01-FY12, CEBAF ops averaged 34.5 weeks/year (best year FY05 at 42 weeks)
- For 12 GeV era we estimate “optimal” operations at 37 weeks per year
- FY17 Pres. Budget includes JLab ops at \$104M
 - would fund 23 weeks (+ 3 weeks from 12 GeV project)
- FY18+ at cost of living implies 23 weeks/year running (62% of optimal)
- We propose FY18+ at 30 weeks/year (81%), will require ~\$6M increase in operations budget.

- Slide from Mont's talk at 2016 UG meeting
- Hall D Physics Program will be completed in 2-3 years

Summary

- KN scattering still remains very poorly studied
- lack of data on excited hyperon states requires significant experimental efforts to be completed
- Our preliminary studies show that few times $10^4 K^0_L/s$ at Jlab is feasible with GlueX setup in Hall D
- Proposed setup will have highest intensity K^0_L beam ever used for hadron spectroscopy
two orders of magnitude higher than
in LASS (SLAC) experiment
- Data obtained at Jlab will be unique and partially complementary to charged kaon data
- The possibility to run with polarized H and D targets is possible (see talk by C. Keith at KL2016 Workshop)

Thank You!