

# The **GLUE**X Meson Program


**Justin Stevens**




**WILLIAM & MARY**

CHARTERED 1693


# Confined states of quarks and gluons




mesons



baryons



tetraquark



pentaquark

Observed mesons and baryons well described by 1<sup>st</sup> principles QCD

But these aren't the only states permitted by QCD

A SCHEMATIC MODEL OF BARYONS AND MESONS \*


M. GELL-MANN

*California Institute of Technology, Pasadena, California*


... Baryons can now be constructed from quarks by using the combinations  $(qqq)$ ,  $(qqqq\bar{q})$ , etc., while mesons are made out of  $(q\bar{q})$ ,  $(qq\bar{q}\bar{q})$ , etc. ...

[Phys. Lett. 8 \(1964\) 214](#)

# Confined states of quarks and gluons




mesons




baryons

Observed mesons and baryons well described by 1<sup>st</sup> principles QCD


But these aren't the only states permitted by QCD




tetraquark



pentaquark



glueball




hybrid meson

Do gluonic degrees of freedom manifest themselves in the bound states we observe in nature?

# Hybrid mesons and gluonic excitations

- \* Excited gluonic field coupled to  $q\bar{q}$  pair
- \* Rich spectrum of hybrid mesons predicted by Lattice QCD
- \* Gluonic field with  $J^{PC} = 1^{+-}$  and mass = 1-1.5 GeV
- \* “Exotic”  $J^{PC}$  : not simple  $q\bar{q}$  from the non-rel. quark model

$$J^{PC} = 0^{+-}, 1^{-+}, 2^{+-} \dots$$




hybrid meson

$$\begin{aligned} \vec{J} &= \vec{L} + \vec{S} \\ P &= (-1)^{L+1} \\ C &= (-1)^{L+S} \end{aligned}$$

# Lattice QCD

Dudek et al. PRD 88 (2013) 094505



$$u\bar{u} + d\bar{d} \quad \blacksquare$$

$$s\bar{s} \quad \blacksquare$$

$$\phi = |s\bar{s}\rangle$$


$$\omega = |u\bar{u} + d\bar{d}\rangle$$

$$\pi^0 = |u\bar{u} - d\bar{d}\rangle$$

**Note:**  $m_\pi = 392 \text{ MeV}$


# Lattice QCD

Dudek et al. PRD 88 (2013) 094505





# Lattice QCD

Dudek et al. PRD 88 (2013) 094505





# Lattice QCD

Dudek et al. PRD 88 (2013) 094505




Most experimental searches for hybrids limited to the  $\pi_1$  state






# GLUEX in Hall D


- \* Linearly polarized photon beam from CEBAF 12 GeV
- \* Large acceptance detector for both charged and neutral particles
- \* ~45 billion events collected in 2017, expect ~2x more data this year




# Photon Beam and Tagger




## Measured Flux




## Measured Polarization



# Exotic $J^{PC}$ in photoproduction





**Meson X with particular  $J^{PC}$**



**Production through t-channel  
“quasi-particle” exchange**

# Non-exotic $J^{PC}$ in photoproduction



**Exchange  $J^{PC}$**

$1^{--} : \omega, \rho$

$1^{+-} : b, h$

- \* Begin by understanding non-exotic production mechanism
- \* Linear photon beam polarization critical to filter out “naturality” of the exchange particle

# Early **GLUEX** physics: $\gamma p \rightarrow \pi^0 p$

## High-Energy $\pi^0$ Photoproduction from Hydrogen with Unpolarized and Linearly Polarized Photons\*

R. L. Anderson, D. B. Gustavson, J. R. Johnson, I. D. Overman, D. M. Ritson, and B. H. Wiik

*Stanford Linear Accelerator Center, Stanford, California 94305*


and

D. Worcester†

*Harvard University, Cambridge, Massachusetts 02138*

(Received 25 June 1971)


1 OCTOBER 1971



# Early **GLUEX** physics: $\gamma p \rightarrow \pi^0 p$

## High-Energy $\pi^0$ Photoproduction from Hydrogen with Unpolarized and Linearly Polarized Photons\*

R. L. Anderson, D. B. Gustavson, J. R. Johnson, I. D. Overman, D. M. Ritson, and B. H. Wiik  
*Stanford Linear Accelerator Center, Stanford, California 94305*  
 and  
 D. Worcester†  
*Harvard University, Cambridge, Massachusetts 02138*  
 (Received 25 June 1971)




**Exchange  $J^{PC}$**


$1^{--} : \omega, \rho$

$1^{+-} : b, h$

1 OCTOBER 1971



$$\frac{d\sigma}{dt} = \sigma_{\perp} + \sigma_{\parallel} = |\rho + \omega|^2 + |b + h|^2$$



**JPAC: Mathieu et al. PRD 92, 074013**


# $\gamma p \rightarrow \pi^0 p$ beam asymmetry $\Sigma$

- \* Beam asymmetry  $\Sigma$  provides insight into dominant production mechanism

$$\Sigma = \frac{|\omega + \rho|^2 - |h + b|^2}{|\omega + \rho|^2 + |h + b|^2}$$

- \* From experimental standpoint it's easily extended to  $\gamma p \rightarrow \eta p$


\* **No previous measurements!**



**Exchange  $J^{PC}$**

$1^{--} : \omega, \rho$


$1^{+-} : b, h$



**JPAC: Mathieu et al. PRD 92, 074013**

# $\pi^0$ and $\eta$ beam asymmetries



$$\gamma p \rightarrow p \gamma \gamma$$



**Phys. Rev. C 95, 042201(R)**






# $\pi^0$ and $\eta$ beam asymmetries



- \* Dip in multiple theory predictions not observed
- \* Indication of vector exchange dominance at this energy
- \* Additional asymmetry measurements ongoing with this dataset



**First 12 GeV publication!**  
**Phys. Rev. C 95, 042201(R)**

# Pseudoscalar beam asymmetries






**Neutral pseudoscalars:  $\Sigma \sim 1$ , dominated by vector exchange**

# Pseudoscalar beam asymmetries




**Charged pseudoscalars: more complicated  $-t$  dependence**

# Early spectroscopy opportunities






- \* Enhancement consistent with earlier SLAC measurement, but  $\sim 100x$  more statistics with early GlueX data
- \* Polarization observables will provide further insight into the nature of this enhancement

# Early spectroscopy opportunities




- ✱ Previous photoproduction data very sparse for channels with multiple neutrals particles
- ✱ Preliminary studies are already showing interesting features




# Mapping the meson spectrum

PRD 88 (2013) 094505



- \* Already studying polarization observables for **“simple” final states**
- \* Beginning to identify **known mesons** in multi-particle final states

- \* Lattice predicts **strange** and **light** quark content for mesons
- \* Search for a **pattern** of hybrid states in many final states
- \* Requires clean identification of charged pions and kaons



	Approximate Mass (MeV)	$J^{PC}$	Final States
$\pi_1$	1900	$1^{-+}$	$\omega\pi\pi^\dagger, 3\pi^\dagger, 5\pi, \eta 3\pi^\dagger, \eta'\pi^\dagger$
$\eta_1$	2100	$1^{-+}$	$4\pi, \eta 4\pi, \eta\eta\pi\pi^\dagger$
$\eta'_1$	2300	$1^{-+}$	$KK\pi\pi^\dagger, KK\pi^\dagger, KK\omega^\dagger$

# Strangeness program: decay patterns


- \* Experimentally infer quark flavor composition through branching ratios to strange and non-strange decays

$$\frac{\mathcal{B}(f_2'(1525) \rightarrow \pi\pi)}{\mathcal{B}(f_2'(1525) \rightarrow KK)} \approx 0.009$$

$$\frac{\mathcal{B}(f_2(1270) \rightarrow \pi\pi)}{\mathcal{B}(f_2(1270) \rightarrow KK)} \approx 20$$


- \* Consistent with lattice QCD mixing angle for  $2^{++}$ , and predictions for hybrids

- \* Need capability to detect strange and non-strange to infer hybrid flavor content







# Strangeness program: $Y(2175)$



**Belle:**  $e^+e^- \rightarrow \phi\pi^+\pi^-(\gamma)$



**BES III:**  $J/\psi \rightarrow \eta\phi\pi^+\pi^-$






\*  $Y(2175)$   $J^{PC}=1^{--}$  state observed by 3 experiments

\* Decay pattern similar to  $Y(4260)$  in charmonium

$$Y(2175) \rightarrow \phi\pi^+\pi^- \quad Y(4260) \rightarrow J/\psi\pi^+\pi^-$$

\* Is there evidence for such strangeonium states in photoproduction?



# GLUEX DIRC upgrade



- \* The GlueX **DIRC** (**D**etection of **I**nternally **R**eflected **C**herenkov light) provides new  $K/\pi$  separation and will use components of the BaBar DIRC
- \* Partial installation and commissioning in **2018**

# Expected DIRC performance

**GLUEX** Simulation



- ✳ Significantly extends reach in search for exotic hadrons (hybrid, multi-quark, etc.) containing strange quarks

# Summary

- \* The **GLUEX** experiment is commissioned and the initial meson program is well underway
- \* First results aim at understanding the meson photoproduction mechanism through beam asymmetries and other polarization observables
- \* An upgrade is in progress to improve the identification of charged kaons to enhance the strange meson spectroscopy program

Supported by DE-SC0018224



U.S. DEPARTMENT OF  
**ENERGY**

Office of  
Science

# Backup

# Exotic $J^{PC}$ decays

C. A. Meyer and E. S. Swanson,  
Progress in Particle and Nuclear Physics B82, 21, (2015)

	Approximate Mass (MeV)	$J^{PC}$	Total Width MeV		Allowed Decay Modes
			PSS	IKP	
$\pi_1$	1900	$1^{-+}$	81 – 168	117	$b_1\pi, \pi\rho, \pi f_1, \pi\eta, \pi\eta', \eta a_1, \pi\eta(1295)$
$\eta_1$	2100	$1^{-+}$	59 – 158	107	$\pi a_1, \pi a_2, \eta f_1, \eta f_2, \pi\pi(1300), \eta\eta', KK_1^A, KK_1^B$
$\eta_1'$	2300	$1^{-+}$	95 – 216	172	$KK_1^B, KK_1^A, KK^*, \eta\eta'$
$b_0$	2400	$0^{+-}$	247 – 429	665	$\pi\pi(1300), \pi h_1, \rho f_1, \eta b_1$
$h_0$	2400	$0^{+-}$	59 – 262	94	$\pi b_1, \eta h_1, KK(1460)$
$h_0'$	2500	$0^{+-}$	259 – 490	426	$KK(1460), KK_1^A, \eta h_1$
$b_2$	2500	$2^{+-}$	5 – 11	248	$\pi a_1, \pi a_2, \pi h_1, \eta\rho, \eta b_1, \rho f_1$
$h_2$	2500	$2^{+-}$	4 – 12	166	$\pi\rho, \pi b_1, \eta\omega, \omega b_1$
$h_2'$	2600	$2^{+-}$	5 – 18	79	$KK_1^B, KK_1^A, KK_2^*, \eta h_1$

\* Predictions for the spectrum of hybrids from lattice, **but decay predictions are model dependent**

## 1<sup>-+</sup> channels observed


$$\begin{aligned}\pi\rho &\rightarrow \pi\pi\pi \\ \pi\eta' &\rightarrow \eta\pi\pi\pi \\ \pi b_1 &\rightarrow \omega\pi\pi\end{aligned}$$

## Some additional 1<sup>-+</sup> channels


$$\begin{aligned}\pi a_2 &\rightarrow \eta\pi\pi & \eta f_1 &\rightarrow \eta\eta\pi\pi \\ & & KK^* &\rightarrow KK\pi \\ & & KK_1(1270) &\rightarrow KK\pi\pi\end{aligned}$$

# “Typical” $\gamma p \rightarrow \pi^+ \pi^- p$ event


top view (looking down from above detector)






side view from beam right (south)



BCAL view from downstream looking upstream



# $\pi^0$ and $\eta$ beam asymmetries






$$\sigma = \sigma_0 \left( 1 - P_\gamma \Sigma \cos 2(\phi_p - \phi_\gamma^{\text{lin}}) \right)$$

**Phys. Rev. C 95, 042201(R)**



# $\pi^0$ and $\eta$ beam asymmetries




$$\frac{Y_{\perp} - F_R Y_{\parallel}}{Y_{\perp} + F_R Y_{\parallel}} = P_{\gamma} \Sigma \cos 2\phi_p$$

**Phys. Rev. C 95, 042201(R)**

# Early spectroscopy opportunities

$$\gamma p \rightarrow 5\gamma p$$




$$\gamma p \rightarrow b_1 p, b_1 \rightarrow \omega\pi^0, \omega \rightarrow \pi^0\gamma$$

- \* Successfully reconstructing  $5\gamma$  final state and observe  $b_1$  signal consistent with previous JLab photoproduction experiment (**RadPhi**)

# Amplitude Analysis

- \* **Goal:** Identify  $J^{PC}$  of  $X \rightarrow \pi^+\pi^-\pi^+$
- \* Model the intensity of events at the level of QM amplitudes (allow for interference)



$$I(\vec{x}) = \frac{dN}{d\vec{x}} = \left| \sum_{\alpha}^{N_{\text{amps}}} V_{\alpha} A_{\alpha}(\vec{x}) \right|^2$$




- \* 5-dimensional problem: two new angles at each decay step ( $X$  and  $I$ )

## Example Intensity:

$$X(1^{++}) \rightarrow \rho\pi^+ \text{ (S wave)}$$

$$X \rightarrow \rho\pi^+$$

$$\rho \rightarrow \pi^+\pi^-$$



# Amplitude Analysis

$$I(\vec{x}) = \frac{dN}{d\vec{x}} = \left| \sum_{\alpha}^{N_{\text{amps}}} V_{\alpha} A_{\alpha}(\vec{x}) \right|^2$$


- \* Expand set of possible amplitudes over many  $X$  and  $I$ , and determine  $V_{\alpha}$  via maximum likelihood fit
- \* Good angular acceptance critical for disentangling  $J^{PC}$

## Example Intensities:






$$X(1^{++}) \rightarrow \rho\pi^{+} \text{ (S wave)}$$

$$X(2^{++}) \rightarrow \rho\pi^{+} \text{ (D wave)}$$


$$X \rightarrow \rho\pi^{+}$$



$$\rho \rightarrow \pi^{+}\pi^{-}$$




# Amplitude Analysis



- \* **Simulate** production of known resonances and **exotic hybrid ( $1^{--}$ ) signal** with 1.6% relative strength
- \* Yields correspond to  **$\sim 3.5$  hours** of GlueX data taking (at full intensity)


## GLUEX Simulation




# Charm Quarks at JLab





**5-quark bound state**



**Hadronic molecule**



$$\Lambda_b \rightarrow J/\psi p K^-$$





**LHCb 2015**


# J/ $\psi$ photoproduction at **GLUEX**

$$\gamma p \rightarrow p e^+ e^-$$

MC normalized to  $\phi$  x-sec. kin.fit  $\chi^2 < 200, \theta_e > 2^\circ$




$\sigma(\gamma p \rightarrow J/\psi p)$  Arbitrary Units



# Strangeness program

	$J^{PC}$	Allowed Decay Modes
$\pi_1$	$1^{-+}$	$b_1\pi, \pi\rho, \pi f_1, \pi\eta, \pi\eta', \eta a_1, \pi\eta(1295)$
$\eta_1$	$1^{-+}$	$\pi a_1, \pi a_2, \eta f_1, \eta f_2, \pi\pi(1300), \eta\eta', KK_1^A, KK_1^B$
$\eta'_1$	$1^{-+}$	$KK_1^B, KK_1^A, KK^*, \eta\eta'$
$b_0$	$0^{+-}$	$\pi\pi(1300), \pi h_1, \rho f_1, \eta b_1$
$h_0$	$0^{+-}$	$\pi b_1, \eta h_1, KK(1460)$
$h'_0$	$0^{+-}$	$KK(1460), KK_1^A, \eta h_1$
$b_2$	$2^{+-}$	$\pi a_1, \pi a_2, \pi h_1, \eta\rho, \eta b_1, \rho f_1$
$h_2$	$2^{+-}$	$\pi\rho, \pi b_1, \eta\omega, \omega b_1$
$h'_2$	$2^{+-}$	$KK_1^B, KK_1^A, KK_2^*, \eta h_1$





- \* Mapping the hybrid spectrum requires: large statistics samples of many particle final states in **strange** and **non-strange** decay modes
- \* Experimentally access to strangeness content of the state by comparing strange vs non-strange decay modes







# Hyperon Spectroscopy: $\Xi^- (dss)$







6 GeV



# Hyperon Spectroscopy: $\Xi^- (dss)$



# Hyperon Spectroscopy: $\Xi^- (dss)$



✱ Longer term:  $K_L$  beam facility ([PAC proposal](#))

