

Photon impact factor and k_T -factorization in the next-to-leading order

I. Balitsky

JLAB & ODU

QCD Evolution Workshop 16 May 2012

- Regge limit in QCD.
- High-energy scattering and Wilson lines.
- Evolution equation for color dipoles.
- Leading order: BK equation.
- Conformal composite dipoles and NLO BK kernel in $\mathcal{N} = 4$.
- NLO amplitude in $\mathcal{N} = 4$ SYM
- Photon impact factor.
- NLO BK kernel in QCD.
- k_T -factorization and NLO BFKL.
- Conclusions
- Outlook: color dipoles, gluon light-ray operators and gluon TMDs

r



Strong interactions at asymptotic energies: Froissart bound

Regge limit: $E \gg$ everything else

$$\left. \begin{array}{c} \text{Causality} \\ \text{Unitarity} \end{array} \right\} \Rightarrow \sigma_{\text{tot}} \stackrel{E \rightarrow \infty}{\leq} \ln^2 E \quad \text{Froissart, 1962}$$


Long-standing problem - not explained in any quantum field theory (or string theory) in 50 years!

Experiment: $\sigma_{\text{tot}} \sim s^{0.08}$ ($s \equiv 4E_{\text{c.m.}}^2$). Numerically close to $\ln^2 E$.



In pQCD: Leading Log Approximation \Rightarrow BFKL pomeron

$$s = (p_A + p_B)^2 \simeq 4E^2$$




Leading Log Approximation (LLA):

$$\alpha_s \ll 1, \quad \alpha_s \ln s \sim 1$$

In pQCD: Leading Log Approximation \Rightarrow BFKL pomeron

$$s = (p_A + p_B)^2 \simeq 4E^2$$



Leading Log Approximation (LLA):

$$\alpha_s \ll 1, \quad \alpha_s \ln s \sim 1$$

The sum of gluon ladder diagrams gives

$$\sigma_{\text{tot}} \sim s^{12 \frac{\alpha_s}{\pi} \ln 2} \quad \text{BFKL pomeron}$$


Numerically: for DIS at HERA

$$\sigma \sim s^{0.3 \div 0.5} = x_B^{-0.3 \div 0.5}$$

- qualitatively OK

In pQCD: Leading Log Approximation \Rightarrow BFKL pomeron

$$s = (p_A + p_B)^2 \simeq 4E^2$$



Leading Log Approximation (LLA):

$$\alpha_s \ll 1, \quad \alpha_s \ln s \sim 1$$

The sum of gluon ladder diagrams gives


$$\sigma_{\text{tot}} \sim s^{12 \frac{\alpha_s}{\pi} \ln 2} \quad \text{BFKL pomeron}$$

Numerically: for DIS at HERA

$$\sigma \sim s^{0.3 \div 0.5} = x_B^{-0.3 \div 0.5}$$

- qualitatively OK

Towards the high-energy QCD




$\sigma_{\text{tot}} \sim s^{12 \frac{\alpha_s}{\pi} \ln 2}$ violates
Froissart bound $\sigma_{\text{tot}} \leq \ln^2 s$
 \Rightarrow pre-asymptotic behavior.

True asymptotics as $E \rightarrow \infty = ?$

Possible approaches:

- Sum all logs $\alpha_s^m \ln^n s$
- Reduce high-energy QCD to 2 + 1 effective theory

Towards the high-energy QCD



$\sigma_{\text{tot}} \sim s^{12 \frac{\alpha_s}{\pi} \ln 2}$ violates
Froissart bound $\sigma_{\text{tot}} \leq \ln^2 s$
 \Rightarrow pre-asymptotic behavior.


True asymptotics as $E \rightarrow \infty = ?$

Possible approaches:

- Sum all logs $\alpha_s^m \ln^n s$
- Reduce high-energy QCD to 2 + 1 effective theory

This talk: NLO corrections $\alpha_s^{n+1} \ln^n s$


High-energy scattering and “Wilson lines” in quantum mechanics



WKB approximation: $\Psi \sim e^{\frac{i}{\hbar}S}$

$$\begin{aligned} S &= \int (pdz - Edt) \\ &= -Et + \int^z dz' \sqrt{2m(E - V(z'))} \end{aligned}$$

High-energy scattering and “Wilson lines” in quantum mechanics




WKB approximation: $\Psi \sim e^{\frac{i}{\hbar}S}$

$$\begin{aligned} S &= \int (pdz - Edt) \\ &= -Et + \int_{-\infty}^z dz' \sqrt{2m(E - V(z'))} \end{aligned}$$

High energy: $E \gg V(x) \Rightarrow$

$$\Psi(\vec{r}, t) = e^{-\frac{i}{\hbar}(Et - kx)} e^{-\frac{i}{v\hbar} \int_{-\infty}^z dz' V(z')}$$

High-energy scattering and “Wilson lines” in quantum mechanics



WKB approximation: $\Psi \sim e^{\frac{i}{\hbar}S}$


$$\begin{aligned} S &= \int (pdz - Edt) \\ &= -Et + \int^z dz' \sqrt{2m(E - V(z'))} \end{aligned}$$

High energy: $E \gg V(x) \Rightarrow$

$$\Psi(\vec{r}, t) = e^{-\frac{i}{\hbar}(Et - kx)} e^{-\frac{i}{v\hbar} \int_{-\infty}^z dz' V(z')}$$

Ψ at high energy = free wave \times phase factor ordered along the line $\parallel \vec{v}$.

High-energy scattering and “Wilson lines” in quantum mechanics



WKB approximation: $\Psi \sim e^{\frac{i}{\hbar}S}$

$$\begin{aligned} S &= \int (pdz - Edt) \\ &= -Et + \int^z dz' \sqrt{2m(E - V(z'))} \end{aligned}$$

High energy: $E \gg V(x) \Rightarrow$

$$\Psi(\vec{r}, t) = e^{-\frac{i}{\hbar}(Et - kx)} e^{-\frac{i}{v\hbar} \int_{-\infty}^z dz' V(z')}$$


Ψ at high energy = free wave \times phase factor ordered along the line $\parallel \vec{v}$.

The scattering amplitude is proportional to $\Psi(t = \infty)$ defined by

$$U(x_\perp) = e^{-\frac{i}{v\hbar} \int_{-\infty}^{\infty} dz' V(z' + x_\perp)}$$

Glauber formula: $\sigma_{\text{tot}} = 2 \int d^2 x_\perp [1 - \Re U(x_\perp)]$

High-energy phase factor in QED and QCD




$$\begin{aligned} S_e &= \int dt \left\{ -mc^2 \sqrt{1 - \frac{v^2}{c^2}} - e\Phi + \frac{e}{c} \vec{v} \cdot \vec{A} \right\} \\ &= S_{\text{free}} + \int dt (-e\Phi + \frac{e}{c} \vec{v} \cdot \vec{A}) \end{aligned}$$

⇒ phase factor for the high-energy scattering is

$$\begin{aligned} U(x_\perp, v) &= e^{-\frac{ie}{\hbar c} \int_{-\infty}^{\infty} dt (-e\Phi + \frac{e}{c} \vec{v} \cdot \vec{A})} \\ &= e^{-\frac{ie}{\hbar c} \int_{-\infty}^{\infty} dt \dot{x}_\mu A^\mu(x(t))} \end{aligned}$$

High-energy phase factor in QED and QCD



$$\begin{aligned} S_e &= \int dt \left\{ -mc^2 \sqrt{1 - \frac{v^2}{c^2}} - e\Phi + \frac{e}{c} \vec{v} \cdot \vec{A} \right\} \\ &= S_{\text{free}} + \int dt (-e\Phi + \frac{e}{c} \vec{v} \cdot \vec{A}) \end{aligned}$$

⇒ phase factor for the high-energy scattering is

$$\begin{aligned} U(x_\perp, v) &= e^{-\frac{ie}{\hbar c} \int_{-\infty}^{\infty} dt (-e\Phi + \frac{e}{c} \vec{v} \cdot \vec{A})} \\ &= e^{-\frac{ie}{\hbar c} \int_{-\infty}^{\infty} dt \dot{x}_\mu A^\mu(x(t))} \end{aligned}$$

In QCD $e \rightarrow -g$, $A_\mu \rightarrow A_\mu \equiv A_\mu^a t^a$ t^a - color matrices


$$\Rightarrow U(x_\perp, v) = P \exp \left\{ \frac{ig}{\hbar c} \int_{-\infty}^{\infty} dt \dot{x}_\mu A^\mu(x(t)) \right\}$$

Wilson – line operator

(Later $\hbar = c = 1$)


DIS at high energy

- At high energies, particles move along straight lines \Rightarrow the amplitude of $\gamma^* A \rightarrow \gamma^* A$ scattering reduces to the matrix element of a two-Wilson-line operator (color dipole):



DIS at high energy


- At high energies, particles move along straight lines \Rightarrow the amplitude of $\gamma^* A \rightarrow \gamma^* A$ scattering reduces to the matrix element of a two-Wilson-line operator (color dipole):



$$A(s) = \int \frac{d^2 k_\perp}{4\pi^2} I^A(k_\perp) \langle B | \text{Tr}\{ \mathcal{U}(k_\perp) \mathcal{U}^\dagger(-k_\perp) \} | B \rangle$$

Formally, \rightarrow means the operator expansion in Wilson lines

Light-cone expansion and DGLAP evolution in the NLO




μ^2 - factorization scale (normalization point)

$k_\perp^2 > \mu^2$ - coefficient functions

$k_\perp^2 < \mu^2$ - matrix elements of light-ray operators (normalized at μ^2)

Light-cone expansion and DGLAP evolution in the NLO



μ^2 - factorization scale (normalization point)

$k_\perp^2 > \mu^2$ - coefficient functions

$k_\perp^2 < \mu^2$ - matrix elements of light-ray operators (normalized at μ^2)


OPE in light-ray operators

$$(x-y)^2 \rightarrow 0$$

$$T\{j_\mu(x)j_\nu(y)\} = \frac{x_\xi}{2\pi^2 x^4} \left[1 + \frac{\alpha_s}{\pi} (\ln x^2 \mu^2 + C) \right] \bar{\psi}(x) \gamma_\mu \gamma^\xi \gamma_\nu [x, y] \psi(y) + O(\frac{1}{x^2})$$

$$[x, y] \equiv P e^{ig \int_0^1 du (x-y)^\mu A_\mu(ux+(1-u)y)} - \text{gauge link}$$

Light-cone expansion and DGLAP evolution in the NLO



μ^2 - factorization scale (normalization point)

$k_\perp^2 > \mu^2$ - coefficient functions

$k_\perp^2 < \mu^2$ - matrix elements of light-ray operators (normalized at μ^2)

Renorm-group equation for light-ray operators \Rightarrow DGLAP evolution of parton densities
 $(x - y)^2 = 0$


$$\mu^2 \frac{d}{d\mu^2} \bar{\psi}(x)[x, y] \psi(y) = K_{\text{LO}} \bar{\psi}(x)[x, y] \psi(y) + \alpha_s K_{\text{NLO}} \bar{\psi}(x)[x, y] \psi(y)$$

Four steps of an OPE

- Factorize an amplitude into a product of coefficient functions and matrix elements of relevant operators.
- Find the evolution equations of the operators with respect to factorization scale.
- Solve these evolution equations.
- Convolute the solution with the initial conditions for the evolution and get the amplitude

DIS at high energy: relevant operators

- At high energies, particles move along straight lines \Rightarrow the amplitude of $\gamma^* A \rightarrow \gamma^* A$ scattering reduces to the matrix element of a two-Wilson-line operator (color dipole):




$$A(s) = \int \frac{d^2 k_\perp}{4\pi^2} I^A(k_\perp) \langle B | \text{Tr}\{U(k_\perp)U^\dagger(-k_\perp)\} | B \rangle$$

$$U(x_\perp) = \text{Pexp} \left[ig \int_{-\infty}^{\infty} du \, n^\mu A_\mu(un + x_\perp) \right]$$

Wilson line

DIS at high energy: relevant operators

- At high energies, particles move along straight lines \Rightarrow the amplitude of $\gamma^* A \rightarrow \gamma^* A$ scattering reduces to the matrix element of a two-Wilson-line operator (color dipole):




$$A(s) = \int \frac{d^2 k_\perp}{4\pi^2} I^A(k_\perp) \langle B | \text{Tr}\{U(k_\perp)U^\dagger(-k_\perp)\} | B \rangle$$

$$U(x_\perp) = P \exp \left[ig \int_{-\infty}^{\infty} du n^\mu A_\mu(un + x_\perp) \right] \quad \text{Wilson line}$$

Formally, \rightarrow means the operator expansion in Wilson lines

Rapidity factorization



η - rapidity factorization scale


Rapidity $Y > \eta$ - coefficient function (“impact factor”)

Rapidity $Y < \eta$ - matrix elements of (light-like) Wilson lines with rapidity divergence cut by η

$$U_x^\eta = \text{Pexp} \left[ig \int_{-\infty}^{\infty} dx^+ A_+^\eta(x_+, x_\perp) \right]$$

$$A_\mu^\eta(x) = \int \frac{d^4 k}{(2\pi)^4} \theta(e^\eta - |\alpha_k|) e^{-ik \cdot x} A_\mu(k)$$

High-energy expansion in color dipoles




The high-energy operator expansion is

$$T\{\hat{j}_\mu(x)\hat{j}_\nu(y)\} = \int d^2z_1 d^2z_2 I_{\mu\nu}^{\text{LO}}(z_1, z_2, x, y) \text{Tr}\{\hat{U}_{z_1}^\eta \hat{U}_{z_2}^{\dagger\eta}\}$$

+ NLO contribution

High-energy expansion in color dipoles



η - rapidity factorization scale

Evolution equation for color dipoles

$$\begin{aligned} \frac{d}{d\eta} \text{tr}\{U_x^\eta U_y^{\dagger\eta}\} &= \frac{\alpha_s}{2\pi^2} \int d^2 z \frac{(x-y)^2}{(x-z)^2(y-z)^2} [\text{tr}\{U_x^\eta U_y^{\dagger\eta}\} \text{tr}\{U_x^\eta U_y^{\dagger\eta}\} \\ &\quad - N_c \text{tr}\{U_x^\eta U_y^{\dagger\eta}\}] + \alpha_s K_{\text{NLO}} \text{tr}\{U_x^\eta U_y^{\dagger\eta}\} + O(\alpha_s^2) \end{aligned}$$

(Linear part of $K_{\text{NLO}} = K_{\text{NLO BFKL}}$)

Evolution equation for color dipoles

$$\hat{\mathcal{U}}(x, y) \equiv 1 - \frac{1}{N_c} \text{Tr}\{\hat{U}(x_\perp) \hat{U}^\dagger(y_\perp)\}$$

BK equation

$$\frac{d}{d\eta} \hat{\mathcal{U}}(x, y) = \frac{\alpha_s N_c}{2\pi^2} \int \frac{d^2 z}{(x-z)^2 (y-z)^2} \left\{ \hat{\mathcal{U}}(x, z) + \hat{\mathcal{U}}(z, y) - \hat{\mathcal{U}}(x, y) - \hat{\mathcal{U}}(x, z) \hat{\mathcal{U}}(z, y) \right\}$$

I. B. (1996), Yu. Kovchegov (1999)

Alternative approach: JIMWLK (1997-2000)

Non-linear evolution equation

$$\hat{\mathcal{U}}(x, y) \equiv 1 - \frac{1}{N_c} \text{Tr}\{\hat{U}(x_\perp) \hat{U}^\dagger(y_\perp)\}$$

BK equation

$$\frac{d}{d\eta} \hat{\mathcal{U}}(x, y) = \frac{\alpha_s N_c}{2\pi^2} \int \frac{d^2 z}{(x-z)^2 (y-z)^2} \left\{ \hat{\mathcal{U}}(x, z) + \hat{\mathcal{U}}(z, y) - \hat{\mathcal{U}}(x, y) - \hat{\mathcal{U}}(x, z) \hat{\mathcal{U}}(z, y) \right\}$$

I. B. (1996), Yu. Kovchegov (1999)

Alternative approach: JIMWLK (1997-2000)

LLA for DIS in pQCD \Rightarrow BFKL

(LLA: $\alpha_s \ll 1, \alpha_s \eta \sim 1$)

Non-linear evolution equation

$$\hat{\mathcal{U}}(x, y) \equiv 1 - \frac{1}{N_c} \text{Tr}\{\hat{U}(x_\perp) \hat{U}^\dagger(y_\perp)\}$$

BK equation

$$\frac{d}{d\eta} \hat{\mathcal{U}}(x, y) = \frac{\alpha_s N_c}{2\pi^2} \int \frac{d^2 z}{(x-z)^2 (y-z)^2} \left\{ \hat{\mathcal{U}}(x, z) + \hat{\mathcal{U}}(z, y) - \hat{\mathcal{U}}(x, y) - \hat{\mathcal{U}}(x, z) \hat{\mathcal{U}}(z, y) \right\}$$

I. B. (1996), Yu. Kovchegov (1999)

Alternative approach: JIMWLK (1997-2000)

LLA for DIS in pQCD \Rightarrow BFKL (LLA: $\alpha_s \ll 1, \alpha_s \eta \sim 1$)


LLA for DIS in sQCD \Rightarrow BK eqn (LLA: $\alpha_s \ll 1, \alpha_s \eta \sim 1, \alpha_s A^{1/3} \sim 1$)

(s for semiclassical)

Why NLO correction?

- To check that high-energy OPE works at the NLO level.
- To check conformal invariance of the NLO BK equation(in $\mathcal{N}=4$ SYM)
- To determine the argument of the coupling constant of the BK equation(in QCD).
- To get the region of application of the leading order evolution equation.

Expansion of the amplitude in color dipoles in the NLO



The high-energy operator expansion is



$$\mathcal{O} \equiv \text{Tr}\{Z^2\}$$

$$T\{\hat{\mathcal{O}}(x)\hat{\mathcal{O}}(y)\} = \int d^2 z_1 d^2 z_2 I^{\text{LO}}(z_1, z_2) \text{Tr}\{\hat{U}_{z_1}^\eta \hat{U}_{z_2}^{\dagger\eta}\} \\ + \int d^2 z_1 d^2 z_2 d^2 z_3 I^{\text{NLO}}(z_1, z_2, z_3) \left[\frac{1}{N_c} \text{Tr}\{T^n \hat{U}_{z_1}^\eta \hat{U}_{z_3}^{\dagger\eta} T^n \hat{U}_{z_3}^\eta \hat{U}_{z_2}^{\dagger\eta}\} - \text{Tr}\{\hat{U}_{z_1}^\eta \hat{U}_{z_2}^{\dagger\eta}\} \right]$$

In the leading order - conf. invariant impact factor

$$I_{\text{LO}} = \frac{x_+^{-2} y_+^{-2}}{\pi^2 \mathcal{Z}_1^2 \mathcal{Z}_2^2}, \quad \mathcal{Z}_i \equiv \frac{(x - z_i)_\perp^2}{x_+} - \frac{(y - z_i)_\perp^2}{y_+} \quad CCP, 2007$$

NLO impact factor (in $\mathcal{N} = 4$ SYM)



$$I^{\text{NLO}}(x, y; z_1, z_2, z_3; \eta) = -I^{\text{LO}} \times \frac{\lambda}{\pi^2} \frac{z_{13}^2}{z_{12}^2 z_{23}^2} \left[\ln \frac{\sigma s}{4} \mathcal{Z}_3 - \frac{i\pi}{2} + C \right]$$

The NLO impact factor is not Möbius invariant \Rightarrow the color dipole with the cutoff η is not invariant

However, if we define a composite operator (a - analog of μ^{-2} for usual OPE)

$$\begin{aligned} [\text{Tr}\{\hat{U}_{z_1}^\eta \hat{U}_{z_2}^{\dagger\eta}\}]^{\text{conf}} &= \text{Tr}\{\hat{U}_{z_1}^\eta \hat{U}_{z_2}^{\dagger\eta}\} \\ &+ \frac{\lambda}{2\pi^2} \int d^2 z_3 \frac{z_{12}^2}{z_{13}^2 z_{23}^2} [\text{Tr}\{T^n \hat{U}_{z_1}^\eta \hat{U}_{z_3}^{\dagger\eta} T^n \hat{U}_{z_3}^\eta \hat{U}_{z_2}^{\dagger\eta}\} - N_c \text{Tr}\{\hat{U}_{z_1}^\eta \hat{U}_{z_2}^{\dagger\eta}\}] \ln \frac{az_{12}^2}{z_{13}^2 z_{23}^2} + O(\lambda^2) \end{aligned}$$

the impact factor becomes conformal in the NLO.

Operator expansion in conformal dipoles in $\mathcal{N} = 4$ SYM

Conformal composite dipole

$$\begin{aligned} [\text{Tr}\{\hat{U}_{z_1}^\eta \hat{U}_{z_2}^{\dagger\eta}\}]^{\text{conf}} &= \text{Tr}\{\hat{U}_{z_1}^\eta \hat{U}_{z_2}^{\dagger\eta}\} \\ &+ \frac{\lambda}{2\pi^2} \int d^2 z_3 \frac{z_{12}^2}{z_{13}^2 z_{23}^2} [\text{Tr}\{T^n \hat{U}_{z_1}^\eta \hat{U}_{z_3}^{\dagger\eta} T^n \hat{U}_{z_3}^\eta \hat{U}_{z_2}^{\dagger\eta}\} - N_c \text{Tr}\{\hat{U}_{z_1}^\eta \hat{U}_{z_2}^{\dagger\eta}\}] \ln \frac{az_{12}^2}{z_{13}^2 z_{23}^2} + O(\lambda^2) \end{aligned}$$

High-energy OPE:

$$\begin{aligned} T\{\hat{\mathcal{O}}(x)\hat{\mathcal{O}}(y)\} &= \int d^2 z_1 d^2 z_2 I^{\text{LO}}(z_1, z_2) \text{Tr}\{\hat{U}_{z_1}^\eta \hat{U}_{z_2}^{\dagger\eta}\}^{\text{conf}} \\ &+ \int d^2 z_1 d^2 z_2 d^2 z_3 I^{\text{NLO}}(z_1, z_2, z_3) \left[\frac{1}{N_c} \text{Tr}\{T^n \hat{U}_{z_1}^\eta \hat{U}_{z_3}^{\dagger\eta} T^n \hat{U}_{z_3}^\eta \hat{U}_{z_2}^{\dagger\eta}\} - \text{Tr}\{\hat{U}_{z_1}^\eta \hat{U}_{z_2}^{\dagger\eta}\} \right] \end{aligned}$$

I^{LO} and I^{NLO} are Möbius invariant.

We think that one can construct the composite conformal dipole operator order by order in perturbation theory.

Analogy: when the UV cutoff does not respect the symmetry of a local operator, the composite local renormalized operator in must be corrected by finite counterterms order by order in perturbation theory.

NLO BK equation in $\mathcal{N} = 4$ SYM

Define


$$\begin{aligned} \hat{\mathcal{U}}_{\text{conf}}^a(z_1, z_2) \\ = \hat{\mathcal{U}}^\eta(z_1, z_2) + \frac{\alpha_s N_c}{4\pi^2} \int d^2 z \frac{z_{12}^2}{z_{13}^2 z_{23}^2} \ln \frac{ae^{2\eta} z_{12}^2}{z_{13}^2 z_{23}^2} [\text{Tr}\{T^a \hat{U}_{z_1}^\eta \hat{U}_{z_3}^{\dagger\eta} T^a \hat{U}_{z_3} \hat{U}_{z_2}^{\dagger\eta}\} - N_c \text{Tr}\{\hat{U}_{z_1}^\eta \hat{U}_{z_2}^{\dagger\eta}\}] \end{aligned}$$

such that $\frac{d}{d\eta} \hat{\mathcal{U}}_{\text{conf}}^a(z_1, z_2) = 0$.

⇒ The evolution can be rewritten in terms of a

$$\begin{aligned} & 2a \frac{d}{da} [\text{Tr}\{\hat{U}_{z_1}^\eta \hat{U}_{z_2}^{\dagger\eta}\}]^{\text{conf}} \\ &= \frac{\alpha_s}{\pi^2} \int d^2 z_3 \frac{z_{12}^2}{z_{13}^2 z_{23}^2} \left[1 - \frac{\alpha_s N_c}{4\pi} \frac{\pi^2}{3} \right] [\text{Tr}\{T^a \hat{U}_{z_1}^\eta \hat{U}_{z_3}^{\dagger\eta} T^a \hat{U}_{z_3} \hat{U}_{z_2}^{\dagger\eta}\} - N_c \text{Tr}\{\hat{U}_{z_1}^\eta \hat{U}_{z_2}^{\dagger\eta}\}]^{\text{conf}} \\ & - \frac{\alpha_s^2}{4\pi^4} \int d^2 z_3 d^2 z_4 \frac{z_{12}^2}{z_{13}^2 z_{24}^2 z_{34}^2} \left\{ 2 \ln \frac{z_{12}^2 z_{34}^2}{z_{14}^2 z_{23}^2} + \left[1 + \frac{z_{12}^2 z_{34}^2}{z_{13}^2 z_{24}^2 - z_{14}^2 z_{23}^2} \right] \ln \frac{z_{13}^2 z_{24}^2}{z_{14}^2 z_{23}^2} \right\} \\ & \times \text{Tr}\{[T^a, T^b] \hat{U}_{z_1}^\eta T^{a'} T^{b'} \hat{U}_{z_2}^{\dagger\eta} + T^b T^a \hat{U}_{z_1}^\eta [T^{b'}, T^{a'}] \hat{U}_{z_2}^{\dagger\eta}\} [(\hat{U}_{z_3}^\eta)^{aa'} (\hat{U}_{z_4}^\eta)^{bb'} - (z_4 \rightarrow z_3)] \end{aligned}$$

NLO Amplitude in $\mathcal{N}=4$ SYM theory




$$(x-y)^4(x'-y')^4 \langle T\{\hat{O}(x)\hat{O}^\dagger(y)\hat{O}(x')\hat{O}^\dagger(y')\} \rangle \\ = \int d^2z_{1\perp} d^2z_{2\perp} d^2z'_{1\perp} d^2z'_{2\perp} \text{IF}^{a_0}(x, y; z_1, z_2) [\text{DD}]^{a_0, b_0}(z_1, z_2; z'_1, z'_2) \text{IF}^{b_0}(x', y'; z'_1, z'_2)$$

Result :

(G.A. Chirilli and I.B.)

$$F(\nu) = \frac{N_c^2}{N_c^2 - 1} \frac{4\pi^4 \alpha_s^2}{\cosh^2 \pi \nu} \left\{ 1 + \frac{\alpha_s N_c}{\pi} \left[-\frac{2\pi^2}{\cosh^2 \pi \nu} + \frac{\pi^2}{2} - \frac{8}{1+4\nu^2} \right] + O(\alpha_s^2) \right\}$$

NLO high-energy OPE in QCD



DIS structure function $F_2(x)$: photon impact factor + evolution of color dipoles+ initial conditions for the small- x evolution

Photon impact factor in the LO

$$(x-y)^4 T\{\bar{\psi}(x)\gamma^\mu \hat{\psi}(x)\bar{\psi}(y)\gamma^\nu \hat{\psi}(y)\} = \int \frac{d^2 z_1 d^2 z_2}{z_{12}^4} I_{\mu\nu}^{\text{LO}}(z_1, z_2) \text{tr}\{\hat{U}_{z_1}^\eta \hat{U}_{z_2}^{\dagger\eta}\}$$

$$I_{\mu\nu}^{\text{LO}}(z_1, z_2) = \frac{\mathcal{R}^2}{\pi^6 (\kappa \cdot \zeta_1)(\kappa \cdot \zeta_2)} \frac{\partial^2}{\partial x^\mu \partial y^\nu} [(\kappa \cdot \zeta_1)(\kappa \cdot \zeta_2) - \frac{1}{2} \kappa^2 (\zeta_1 \cdot \zeta_2)].$$

$$\kappa \equiv \frac{1}{\sqrt{s}x^+} \left(\frac{p_1}{s} - x^2 p_2 + x_\perp \right) - \frac{1}{\sqrt{s}y^+} \left(\frac{p_1}{s} - y^2 p_2 + y_\perp \right)$$

$$\zeta_i \equiv \left(\frac{p_1}{s} + z_{i\perp}^2 p_2 + z_{i\perp} \right), \quad \mathcal{R} \equiv \frac{\kappa^2 (\zeta_1 \cdot \zeta_2)}{2(\kappa \cdot \zeta_1)(\kappa \cdot \zeta_2)}$$

Composite “conformal” dipole $[\text{tr}\{\hat{U}_{z_1} \hat{U}_{z_2}^\dagger\}]_{a_0}$ - same as in $\mathcal{N} = 4$ case.

$$\begin{aligned}
 & (x-y)^4 T\{\bar{\psi}(x)\gamma^\mu \hat{\psi}(x) \bar{\psi}(y)\gamma^\nu \hat{\psi}(y)\} \\
 &= \int \frac{d^2 z_1 d^2 z_2}{z_{12}^4} \left\{ I_{\text{LO}}^{\mu\nu}(z_1, z_2) \left[1 + \frac{\alpha_s}{\pi} \right] [\text{tr}\{\hat{U}_{z_1} \hat{U}_{z_2}^\dagger\}]_{a_0} \right. \\
 &+ \int d^2 z_3 \left[\frac{\alpha_s}{4\pi^2} \frac{z_{12}^2}{z_{13}^2 z_{23}^2} \left(\ln \frac{\kappa^2 (\zeta_1 \cdot \zeta_3)(\zeta_1 \cdot \zeta_3)}{2(\kappa \cdot \zeta_3)^2 (\zeta_1 \cdot \zeta_2)} - 2C \right) I_{\text{LO}}^{\mu\nu} + I_2^{\mu\nu} \right] \\
 &\quad \times [\text{tr}\{\hat{U}_{z_1} \hat{U}_{z_3}^\dagger\} \text{tr}\{\hat{U}_{z_3} \hat{U}_{z_2}^\dagger\} - N_c \text{tr}\{\hat{U}_{z_1} \hat{U}_{z_2}^\dagger\}]_{a_0} \Big\}
 \end{aligned}$$

$$\begin{aligned}
 (I_2)_{\mu\nu}(z_1, z_2, z_3) &= \frac{\alpha_s}{16\pi^8} \frac{\mathcal{R}^2}{(\kappa \cdot \zeta_1)(\kappa \cdot \zeta_2)} \left\{ \frac{(\kappa \cdot \zeta_2)}{(\kappa \cdot \zeta_3)} \frac{\partial^2}{\partial x^\mu \partial y^\nu} \left[-\frac{(\kappa \cdot \zeta_1)^2}{(\zeta_1 \cdot \zeta_3)} \right. \right. \\
 &+ \frac{(\kappa \cdot \zeta_1)(\kappa \cdot \zeta_2)}{(\zeta_2 \cdot \zeta_3)} + \frac{(\kappa \cdot \zeta_1)(\kappa \cdot \zeta_3)(\zeta_1 \cdot \zeta_2)}{(\zeta_1 \cdot \zeta_3)(\zeta_2 \cdot \zeta_3)} - \frac{\kappa^2 (\zeta_1 \cdot \zeta_2)}{(\zeta_2 \cdot \zeta_3)} \Big] \\
 &+ \left. \left. \frac{(\kappa \cdot \zeta_2)^2}{(\kappa \cdot \zeta_3)^2} \frac{\partial^2}{\partial x^\mu \partial y^\nu} \left[\frac{(\kappa \cdot \zeta_1)(\kappa \cdot \zeta_3)}{(\zeta_2 \cdot \zeta_3)} - \frac{\kappa^2 (\zeta_1 \cdot \zeta_3)}{2(\zeta_2 \cdot \zeta_3)} \right] + (\zeta_1 \leftrightarrow \zeta_2) \right\}
 \end{aligned}$$

With two-gluon (NLO BFKL) accuracy

$$\frac{1}{N_c} (x-y)^4 T \{ \bar{\psi}(x) \gamma^\mu \hat{\psi}(x) \bar{\psi}(y) \gamma^\nu \hat{\psi}(y) \} = \frac{\partial \kappa^\alpha}{\partial x^\mu} \frac{\partial \kappa^\beta}{\partial y^\nu} \int \frac{dz_1 dz_2}{z_{12}^4} \hat{U}_{a_0}(z_1, z_2) [\mathcal{I}_{\alpha\beta}^{\text{LO}} \left(1 + \frac{\alpha_s}{\pi}\right) + \mathcal{I}_{\alpha\beta}^{\text{NLO}}]$$

$$\mathcal{I}_{\text{LO}}^{\alpha\beta}(x, y; z_1, z_2) = \mathcal{R}^2 \frac{g^{\alpha\beta}(\zeta_1 \cdot \zeta_2) - \zeta_1^\alpha \zeta_2^\beta - \zeta_2^\alpha \zeta_1^\beta}{\pi^6 (\kappa \cdot \zeta_1)(\kappa \cdot \zeta_2)}$$

$$\begin{aligned} \mathcal{I}_{\text{NLO}}^{\alpha\beta}(x, y; z_1, z_2) = & \frac{\alpha_s N_c}{4\pi^7} \mathcal{R}^2 \left\{ \frac{\zeta_1^\alpha \zeta_2^\beta + \zeta_1 \leftrightarrow \zeta_2}{(\kappa \cdot \zeta_1)(\kappa \cdot \zeta_2)} \left[4\text{Li}_2(1-\mathcal{R}) - \frac{2\pi^2}{3} + \frac{2 \ln \mathcal{R}}{1-\mathcal{R}} + \frac{\ln \mathcal{R}}{\mathcal{R}} \right. \right. \\ & - 4 \ln \mathcal{R} + \frac{1}{2\mathcal{R}} - 2 + 2 \left(\ln \frac{1}{\mathcal{R}} + \frac{1}{\mathcal{R}} - 2 \right) \left(\ln \frac{1}{\mathcal{R}} + 2C \right) - 4C - \frac{2C}{\mathcal{R}} \Big] \\ & + \left(\frac{\zeta_1^\alpha \zeta_1^\beta}{(\kappa \cdot \zeta_1)^2} + \zeta_1 \leftrightarrow \zeta_2 \right) \left[\frac{\ln \mathcal{R}}{\mathcal{R}} - \frac{2C}{\mathcal{R}} + 2 \frac{\ln \mathcal{R}}{1-\mathcal{R}} - \frac{1}{2\mathcal{R}} \right] - \frac{2}{\kappa^2} \left(g^{\alpha\beta} - 2 \frac{\kappa^\alpha \kappa^\beta}{\kappa^2} \right) \\ & + \left[\frac{\zeta_1^\alpha \kappa^\beta + \zeta_1^\beta \kappa^\alpha}{(\kappa \cdot \zeta_1)\kappa^2} + \zeta_1 \leftrightarrow \zeta_2 \right] \left[-2 \frac{\ln \mathcal{R}}{1-\mathcal{R}} - \frac{\ln \mathcal{R}}{\mathcal{R}} + \ln \mathcal{R} - \frac{3}{2\mathcal{R}} + \frac{5}{2} + 2C + \frac{2C}{\mathcal{R}} \right] \\ & + \frac{g^{\alpha\beta}(\zeta_1 \cdot \zeta_2)}{(\kappa \cdot \zeta_1)(\kappa \cdot \zeta_2)} \left[\frac{2\pi^2}{3} - 4\text{Li}_2(1-\mathcal{R}) \right. \\ & \left. \left. - 2 \left(\ln \frac{1}{\mathcal{R}} + \frac{1}{\mathcal{R}} + \frac{1}{2\mathcal{R}^2} - 3 \right) \left(\ln \frac{1}{\mathcal{R}} + 2C \right) + 6 \ln \mathcal{R} - \frac{2}{\mathcal{R}} + 2 + \frac{3}{2\mathcal{R}^2} \right] \right\} \end{aligned}$$

5 tensor structures (CCP, 2009)

Photon Impact Factor at NLO

Reminder

$$\begin{aligned}\kappa^\mu &= \frac{1}{\sqrt{s}x^+} \left(\frac{p_1^\mu}{s} - x^2 p_2^\mu + x_\perp^\mu \right) - \frac{1}{\sqrt{s}y^+} \left(\frac{p_1^\mu}{s} - y^2 p_2^\mu + y_\perp^\mu \right) \\ \zeta_1^\mu &= \left(\frac{p_1^\mu}{s} + z_{1\perp}^2 p_2^\mu + z_{1\perp}^\mu \right), \quad \zeta_2^\mu = \left(\frac{p_1^\mu}{s} + z_{2\perp}^2 p_2^\mu + z_{2\perp}^\mu \right)\end{aligned}$$

DIS photon impact factor is a linear combination of the following tensor basis

$$\mathcal{I}_1^{\mu\nu} = g^{\mu\nu} \quad \mathcal{I}_2^{\mu\nu} = \frac{\kappa^\mu \kappa^\nu}{\kappa^2}$$

$$\mathcal{I}_3^{\mu\nu} = \frac{\kappa^\mu \zeta_1^\nu + \kappa^\nu \zeta_1^\mu}{\kappa \cdot \zeta_1} + \frac{\kappa^\mu \zeta_2^\nu + \kappa^\nu \zeta_2^\mu}{\kappa \cdot \zeta_2}$$

$$\mathcal{I}_4^{\mu\nu} = \frac{\kappa^2 \zeta_1^\mu \zeta_1^\nu}{(\kappa \cdot \zeta_1)^2} + \frac{\kappa^2 \zeta_2^\mu \zeta_2^\nu}{(\kappa \cdot \zeta_2)^2} \quad \mathcal{I}_5^{\mu\nu} = \frac{\zeta_1^\mu \zeta_2^\nu + \zeta_2^\mu \zeta_1^\nu}{\zeta_1 \cdot \zeta_2}$$

Cornalba, Costa, Penedones (2010)

Mellin representation of the LO impact factor

$$\int \frac{d^2 z_1 d^2 z_2}{z_{12}^2} I_{LO}^{\mu\nu}(z_1, z_2) \left(\frac{z_{12}^2}{z_{10}^2 z_{20}^2} \right)^\gamma = \frac{1}{\pi^4} B(1-\gamma) \Gamma(\gamma+2) \Gamma(3-\gamma) \\ \times \left\{ \frac{\gamma(1-\gamma) D_1^{\mu\nu}}{12(1+\gamma)(2-\gamma)} + \frac{D_2^{\mu\nu}}{2(1+\gamma)(2-\gamma)} - \frac{D_3^{\mu\nu}}{8(1+\gamma)(2-\gamma)} \right. \\ \left. - \frac{\gamma(1-\gamma) D_4}{16(1+2\gamma)(3-2\gamma)(1+\gamma)(2-\gamma)} - \frac{D_1^{\mu\nu} + D_2^{\mu\nu}}{8} \right\}_{\mu\nu} \left(\frac{\kappa^2}{(\kappa \cdot \zeta_0)^2} \right)^\gamma$$

where

$$(D_1 + D_2)^{\mu\nu} = -2\Delta^2 x^+ y^+ \kappa^{-2} \partial_x^\mu \partial_y^\nu \kappa^2$$

$$D_2^{\mu\nu} = -\Delta^2 x^+ y^+ \partial_x^\mu (\ln \kappa^2) \partial_y^\nu \ln \kappa^2$$

$$D_3^{\mu\nu} = 4\gamma \Delta^2 x^+ y^+ [(\partial_x^\mu \ln \kappa^2) \partial_y^\nu \ln(\kappa \cdot \zeta_0) + (\partial_y^\nu \ln \kappa^2) \partial_x^\mu \ln(\kappa \cdot \zeta_0) - (\partial_x^\mu \ln \kappa^2) \partial_y^\nu \ln \kappa^2]$$

$$D_4^{\mu\nu} = 4\gamma(1+2\gamma) \Delta^2 x^+ y^+ [-\frac{1}{3} \partial_x^\mu \partial_y^\nu \ln \kappa^2 - \partial_x^\mu (\ln \kappa^2) \partial_y^\nu \ln \kappa^2]$$

$$+ (\partial_x^\mu \ln \kappa^2) \partial_y^\nu \ln(\kappa \cdot \zeta_0) + (\partial_y^\nu \ln \kappa^2) \partial_x^\mu \ln(\kappa \cdot \zeta_0) - 2\partial_x^\mu \ln(\kappa \cdot \zeta_0) \partial_y^\nu \ln(\kappa \cdot \zeta_0)]$$

$$B(a, b) = \frac{\Gamma(a)\Gamma(b)}{\Gamma(a+b)}, C = -\psi(1) \text{ is the Euler constant, and } \psi'(a) = \frac{d}{da} \ln \Gamma(a)$$

Mellin representation of the photon impact factor (I.B. and G. A. C.)

$$\begin{aligned}
 & \int \frac{d^2 z_1 d^2 z_2}{z_{12}^4} [I_{LO}^{\mu\nu}(z_1, z_2) + {}_{NLO}^{\mu\nu}(z_1, z_2)] \left(\frac{z_{12}^2}{z_{10}^2 z_{20}^2} \right)^\gamma = \frac{N_c}{4\pi^6 \Delta^4} \frac{\Gamma(\gamma+1)\Gamma(2-\gamma)}{\Delta^2 x^+ y^+} \\
 & \times \left[\frac{\bar{\gamma}\gamma D_1}{3} \left\{ 1 + \frac{\alpha_s N_c}{4\pi} \left[\frac{\pi^2}{3} - \frac{\pi^2}{\sin^2 \pi\gamma} - C\chi_\gamma - \frac{1}{\gamma\bar{\gamma}} + \frac{1}{2} - \frac{\chi_\gamma}{\gamma\bar{\gamma}} \right] \right\} \right. \\
 & + 2D_2 \left\{ 1 + \frac{\alpha_s N_c}{4\pi} \left[\frac{\pi^2}{3} - \frac{\pi^2}{\sin^2 \pi\gamma} C\chi_\gamma - \frac{3}{4\gamma\bar{\gamma}} + \frac{1}{2}\chi_\gamma + \frac{\chi_\gamma}{2\gamma\bar{\gamma}} \right] \right\} \\
 & - \frac{D_3}{2} \left\{ 1 + \frac{\alpha_s N_c}{4\pi} \left[\frac{\pi^2}{3} - \frac{\pi^2}{\sin^2 \pi\gamma} - C\chi_\gamma + \frac{1}{2} - \frac{1}{\gamma\bar{\gamma}} + \frac{\chi_\gamma}{4} + \frac{\chi_\gamma}{2\gamma\bar{\gamma}} \right] \right\} \\
 & + \frac{\bar{\gamma}\gamma D_4}{4(3+4\bar{\gamma}\gamma)} \left\{ 1 + \frac{\alpha_s N_c}{4\pi} \left[\frac{\pi^2}{3} - \frac{\pi^2}{\sin^2 \pi\gamma} - C\chi_\gamma + \frac{1}{2} - \frac{4}{\gamma\bar{\gamma}} + \frac{3}{2\gamma^2\bar{\gamma}^2} - \frac{\chi_\gamma}{2\gamma\bar{\gamma}} \right] \right\} \\
 & - \frac{D_1 + D_2}{2} (2 + \bar{\gamma}\gamma) \left\{ 1 + \frac{\alpha_s N_c}{4\pi} \left[\frac{\pi^2}{3} - \frac{\pi^2}{\sin^2 \pi\gamma} - C\chi_\gamma + \frac{1}{2} \right. \right. \\
 & \left. \left. - \frac{4\gamma\bar{\gamma} + 3}{2\gamma\bar{\gamma}(2 + \bar{\gamma}\gamma)} + \frac{1 + 2\gamma\bar{\gamma}}{\gamma\bar{\gamma}(2 + \bar{\gamma}\gamma)} \chi_\gamma \right] \right\}^{\mu\nu} \left(\frac{\kappa^2}{(2\kappa \cdot \zeta_0)^2} \right)^\gamma \frac{\Gamma^2(\bar{\gamma})}{\Gamma(2\bar{\gamma})} \quad \bar{\gamma} \equiv 1 - \gamma
 \end{aligned}$$

Mellin representation of the impact factor for polarized DIS

Contribution of spin 2 in t-channel:

$$\int \frac{d^2 z_1 d^2 z_2}{z_{12}^4} (I_{LO}^{\mu\nu}(z_1, z_2) + I_{NLO}^{\mu\nu}(z_1, z_2)) \left(\frac{z_{12}}{z_{10} z_{20}} \right)^{\gamma+1} \left(\frac{\bar{z}_{12}}{\bar{z}_{10} \bar{z}_{20}} \right)^{\gamma+1} = B(2 - \gamma) \Gamma(3 - \gamma) \Gamma(2 + \gamma)$$
$$\times \left[1 + \frac{\alpha_s N_c}{4\pi} \left\{ 4\psi'(1 + \gamma) + 4\psi'(2 - \gamma) - 8\psi'(3) - \frac{6\chi(2, \gamma)}{(1 + \gamma)(2 - \gamma)} + 6 + 4C\chi(2, \gamma) \right. \right.$$
$$\left. \left. - \frac{6C}{(2 - \gamma)(1 + \gamma)} - \frac{6}{(1 + \gamma)(2 - \gamma)} \right\} \right] \left(\frac{\kappa^2}{(\kappa \cdot \zeta_0)^2} \right)^\gamma \left(\partial_\mu^x \frac{X^2 \bar{Y} - Y^2 \bar{X}}{x+y_+(\kappa \cdot \zeta_0)} \right) \left(\partial_\nu^y \frac{X^2 \bar{Y} - Y^2 \bar{X}}{x+y_+(\kappa \cdot \zeta_0)} \right)$$

$$\chi(2, \gamma) = 2\psi(1) - \psi(2 - \gamma) - \psi(1 + \gamma) \quad X \equiv x - z_0, Y \equiv y - z_0$$

In general

$$\frac{d}{d\eta} \text{Tr}\{\hat{U}_x \hat{U}_y^\dagger\} = \alpha_s K_{\text{LO}} \text{Tr}\{\hat{U}_x \hat{U}_y^\dagger\} + \alpha_s^2 K_{\text{NLO}} \text{Tr}\{\hat{U}_x \hat{U}_y^\dagger\} + O(\alpha_s^3)$$

In general

$$\frac{d}{d\eta} \text{Tr}\{\hat{U}_x \hat{U}_y^\dagger\} = \alpha_s K_{\text{LO}} \text{Tr}\{\hat{U}_x \hat{U}_y^\dagger\} + \alpha_s^2 K_{\text{NLO}} \text{Tr}\{\hat{U}_x \hat{U}_y^\dagger\} + O(\alpha_s^3)$$

$$\alpha_s^2 K_{\text{NLO}} \text{Tr}\{\hat{U}_x \hat{U}_y^\dagger\} = \frac{d}{d\eta} \text{Tr}\{\hat{U}_x \hat{U}_y^\dagger\} - \alpha_s K_{\text{LO}} \text{Tr}\{\hat{U}_x \hat{U}_y^\dagger\} + O(\alpha_s^3)$$

In general

$$\frac{d}{d\eta} \text{Tr}\{\hat{U}_x \hat{U}_y^\dagger\} = \alpha_s K_{\text{LO}} \text{Tr}\{\hat{U}_x \hat{U}_y^\dagger\} + \alpha_s^2 K_{\text{NLO}} \text{Tr}\{\hat{U}_x \hat{U}_y^\dagger\} + O(\alpha_s^3)$$

$$\alpha_s^2 K_{\text{NLO}} \text{Tr}\{\hat{U}_x \hat{U}_y^\dagger\} = \frac{d}{d\eta} \text{Tr}\{\hat{U}_x \hat{U}_y^\dagger\} - \alpha_s K_{\text{LO}} \text{Tr}\{\hat{U}_x \hat{U}_y^\dagger\} + O(\alpha_s^3)$$

We calculate the “matrix element” of the r.h.s. in the shock-wave background

$$\langle \alpha_s^2 K_{\text{NLO}} \text{Tr}\{\hat{U}_x \hat{U}_y^\dagger\} \rangle = \frac{d}{d\eta} \langle \text{Tr}\{\hat{U}_x \hat{U}_y^\dagger\} \rangle - \langle \alpha_s K_{\text{LO}} \text{Tr}\{\hat{U}_x \hat{U}_y^\dagger\} \rangle + O(\alpha_s^3)$$

In general

$$\frac{d}{d\eta} \text{Tr}\{\hat{U}_x \hat{U}_y^\dagger\} = \alpha_s K_{\text{LO}} \text{Tr}\{\hat{U}_x \hat{U}_y^\dagger\} + \alpha_s^2 K_{\text{NLO}} \text{Tr}\{\hat{U}_x \hat{U}_y^\dagger\} + O(\alpha_s^3)$$

$$\alpha_s^2 K_{\text{NLO}} \text{Tr}\{\hat{U}_x \hat{U}_y^\dagger\} = \frac{d}{d\eta} \text{Tr}\{\hat{U}_x \hat{U}_y^\dagger\} - \alpha_s K_{\text{LO}} \text{Tr}\{\hat{U}_x \hat{U}_y^\dagger\} + O(\alpha_s^3)$$

We calculate the “matrix element” of the r.h.s. in the shock-wave background

$$\langle \alpha_s^2 K_{\text{NLO}} \text{Tr}\{\hat{U}_x \hat{U}_y^\dagger\} \rangle = \frac{d}{d\eta} \langle \text{Tr}\{\hat{U}_x \hat{U}_y^\dagger\} \rangle - \langle \alpha_s K_{\text{LO}} \text{Tr}\{\hat{U}_x \hat{U}_y^\dagger\} \rangle + O(\alpha_s^3)$$


Subtraction of the (LO) contribution (with the rigid rapidity cutoff)

$\Rightarrow \left[\frac{1}{v} \right]_+$ prescription in the integrals over Feynman parameter v


Typical integral

$$\int_0^1 dv \frac{1}{(k-p)_\perp^2 v + p_\perp^2 (1-v)} \left[\frac{1}{v} \right]_+ = \frac{1}{p_\perp^2} \ln \frac{(k-p)_\perp^2}{p_\perp^2}$$


Gluon part of the NLO BK kernel: diagrams




Diagrams for $1 \rightarrow 3$ dipoles transition




Diagrams for $1 \rightarrow 3$ dipoles transition



"Running coupling" diagrams



$1 \rightarrow 2$ dipole transition diagrams



NLO evolution of composite “conformal” dipoles in QCD

I. B. and G. Chirilli

$$\begin{aligned}
 a \frac{d}{da} [\text{tr}\{U_{z_1} U_{z_2}^\dagger\}]_a^{\text{conf}} &= \frac{\alpha_s}{2\pi^2} \int d^2 z_3 \left([\text{tr}\{U_{z_1} U_{z_3}^\dagger\} \text{tr}\{U_{z_3} U_{z_2}^\dagger\} - N_c \text{tr}\{U_{z_1} U_{z_2}^\dagger\}]_a^{\text{conf}} \right. \\
 &\times \frac{z_{12}^2}{z_{13}^2 z_{23}^2} \left[1 + \frac{\alpha_s N_c}{4\pi} \left(b \ln z_{12}^2 \mu^2 + b \frac{z_{13}^2 - z_{23}^2}{z_{13}^2 z_{23}^2} \ln \frac{z_{13}^2}{z_{23}^2} + \frac{67}{9} - \frac{\pi^2}{3} \right) \right] \\
 &+ \frac{\alpha_s}{4\pi^2} \int \frac{d^2 z_4}{z_{34}^4} \left\{ \left[-2 + \frac{z_{23}^2 z_{23}^2 + z_{24}^2 z_{13}^2 - 4 z_{12}^2 z_{34}^2}{2(z_{23}^2 z_{23}^2 - z_{24}^2 z_{13}^2)} \ln \frac{z_{23}^2 z_{23}^2}{z_{24}^2 z_{13}^2} \right] \right. \\
 &\times [\text{tr}\{U_{z_1} U_{z_3}^\dagger\} \text{tr}\{U_{z_3} U_{z_4}^\dagger\} \{U_{z_4} U_{z_2}^\dagger\} - \text{tr}\{U_{z_1} U_{z_3}^\dagger U_{z_4} U_{z_2}^\dagger U_{z_3} U_{z_4}^\dagger\} - (z_4 \rightarrow z_3)] \\
 &+ \frac{z_{12}^2 z_{34}^2}{z_{13}^2 z_{24}^2} \left[2 \ln \frac{z_{12}^2 z_{34}^2}{z_{23}^2 z_{23}^2} + \left(1 + \frac{z_{12}^2 z_{34}^2}{z_{13}^2 z_{24}^2 - z_{23}^2 z_{23}^2} \right) \ln \frac{z_{13}^2 z_{24}^2}{z_{23}^2 z_{23}^2} \right] \\
 &\times [\text{tr}\{U_{z_1} U_{z_3}^\dagger\} \text{tr}\{U_{z_3} U_{z_4}^\dagger\} \text{tr}\{U_{z_4} U_{z_2}^\dagger\} - \text{tr}\{U_{z_1} U_{z_4}^\dagger U_{z_3} U_{z_2}^\dagger U_{z_4} U_{z_3}^\dagger\} - (z_4 \rightarrow z_3)] \Big\} \\
 b &= \frac{11}{3} N_c - \frac{2}{3} n_f
 \end{aligned}$$

$K_{\text{NLO BK}}$ = Running coupling part + Conformal "non-analytic" (in j) part
+ Conformal analytic ($\mathcal{N} = 4$) part

Linearized $K_{\text{NLO BK}}$ reproduces the known result for the forward NLO BFKL kernel.

k_T -factorization in the NLO

With two-gluon (one-dipole) accuracy

$$\int d^4x e^{iqx} \int d^4z \delta(z_-) \langle p_B | T\{\hat{j}_\mu(x+z)\hat{j}_\nu(z)\} | p_B \rangle = \int d^2k_\perp I_{\mu\nu}(q, k_\perp) \langle\langle p_B | \mathcal{U}(k_\perp) | p_B \rangle\rangle$$

$$\langle p_B | \mathcal{U}(k) | p_B + \beta p_B \rangle = 2\pi\delta(\beta) \langle\langle p_B | \mathcal{U}(k) | p_B \rangle\rangle$$

$$\langle\langle p_B | \mathcal{U}(k) | p_B \rangle\rangle = \int d^2z e^{-i(k,z)_\perp} \langle\langle p_B | \mathcal{U}(z) | p_B \rangle\rangle, \quad \mathcal{U}(z) \equiv 1 - \frac{1}{N_c} \text{Tr}\{U_z U_0^\dagger\}$$

k_T -factorization in the NLO

With two-gluon (one-dipole) accuracy

$$\int d^4x e^{iqx} \int d^4z \delta(z_-) \langle p_B | T\{\hat{j}_\mu(x+z)\hat{j}_\nu(z)\} | p_B \rangle = \int d^2k_\perp I_{\mu\nu}(q, k_\perp) \langle\langle p_B | \mathcal{U}(k_\perp) | p_B \rangle\rangle$$

$$\langle p_B | \mathcal{U}(k) | p_B + \beta p_B \rangle = 2\pi\delta(\beta) \langle\langle p_B | \mathcal{U}(k) | p_B \rangle\rangle$$

$$\langle\langle p_B | \mathcal{U}(k) | p_B \rangle\rangle = \int d^2z e^{-i(k,z)_\perp} \langle\langle p_B | \mathcal{U}(z) | p_B \rangle\rangle, \quad \mathcal{U}(z) \equiv 1 - \frac{1}{N_c} \text{Tr}\{U_z U_0^\dagger\}$$

$$I^{\mu\nu}(q, k_\perp) = \frac{N_c}{32} \int \frac{d\nu}{\pi\nu} \frac{\sinh \pi\nu}{(1+\nu^2) \cosh^2 \pi\nu} \left(\frac{k_\perp^2}{Q^2}\right)^{\frac{1}{2}-i\nu} \\ \times \left[\left(\frac{9}{4} + \nu^2 \right) \left(1 + \frac{\alpha_s}{\pi} + \frac{\alpha_s N_c}{2\pi} \mathcal{F}_1(\nu) \right) \color{red} P_1^{\mu\nu} + \left(\frac{11}{4} + 3\nu^2 \right) \left(1 + \frac{\alpha_s}{\pi} + \frac{\alpha_s N_c}{2\pi} \mathcal{F}_2(\nu) \right) \color{blue} P_2^{\mu\nu} \right]$$

$$P_1^{\mu\nu} = g^{\mu\nu} - \frac{q_\mu q_\nu}{q^2} \quad P_2^{\mu\nu} = \frac{1}{q^2} \left(q^\mu - \frac{p_2^\mu q^2}{q \cdot p_2} \right) \left(q^\nu - \frac{p_2^\nu q^2}{q \cdot p_2} \right)$$

$$\mathcal{F}_{1(2)}(\nu) = \Phi_{1(2)}(\nu) + \chi_\gamma \Psi(\nu),$$

$$\Psi(\nu) \equiv \psi(\bar{\gamma}) + 2\psi(2-\gamma) - 2\psi(4-2\gamma) - \psi(2+\gamma), \quad \gamma \equiv \frac{1}{2} + i\nu$$

k_T -factorization in the NLO

$$\Phi_1(\nu) = F(\gamma) + \frac{3\chi_\gamma}{2 + \bar{\gamma}\gamma} + 1 + \frac{25}{18(2 - \gamma)} + \frac{1}{2\bar{\gamma}} - \frac{1}{2\gamma} - \frac{7}{18(1 + \gamma)} + \frac{10}{3(1 + \gamma)^2}$$

$$\Phi_2(\nu) = F(\gamma) + \frac{3\chi_\gamma}{2 + \bar{\gamma}\gamma} + 1 + \frac{1}{2\bar{\gamma}\gamma} - \frac{7}{2(2 + 3\bar{\gamma}\gamma)} + \frac{\chi_\gamma}{1 + \gamma} + \frac{\chi_\gamma(1 + 3\gamma)}{2 + 3\bar{\gamma}\gamma}$$

$$F(\gamma) = \frac{2\pi^2}{3} - \frac{2\pi^2}{\sin^2 \pi\gamma} - 2C\chi_\gamma + \frac{\chi_\gamma - 2}{\bar{\gamma}\gamma}$$

Evolution equation for color dipole in momentum representation

$$\mathcal{V}_a(z) \equiv z^{-2} \mathcal{U}_a(z)$$

$\mathcal{V}_a(k) \equiv \int dz e^{-i(k,z)_\perp} \mathcal{V}_a(z)$ (sometimes called “unintegrated gluon TMD”)

$$\begin{aligned} 2a \frac{d}{da} \mathcal{V}_a(k) &= \frac{\alpha_s N_c}{2\pi^2} \int \frac{d^2 k'}{(k - k')^2} \left\{ \left(2\mathcal{V}(k') - \frac{k^2}{k'^2} \mathcal{V}_a(k) \right) \right. \\ &+ \frac{\alpha_s b}{4\pi} \left[\left(2\mathcal{V}(k') - \frac{k^2}{k'^2} \mathcal{V}_a(k) \right) \left(\ln \frac{\mu^2}{k^2} + \left(\frac{67}{9} - \frac{\pi^2}{3} \right) - \frac{10n_f}{9N_c} \right) \right. \\ &- 2 \left(\mathcal{V}_a(k') \ln \frac{(k - k')^2}{k'^2} - \mathcal{V}_a(k) \frac{k^2}{k'^2} \ln \frac{(k - k')^2}{k^2} \right) + \mathcal{V}_a(k') \frac{4(k', k - k')}{k'^2} \ln \frac{(k - k')^2}{k^2} \left. \right] \} \\ &+ \frac{\alpha_s^2 N_c^2}{4\pi^3} \int d^2 k' \left[-\frac{1}{(k - k')^2} \ln^2 \frac{k^2}{k'^2} + F(k, k') + \Phi(k, k') \right] \mathcal{V}_a(k') + 3 \frac{\alpha_s^2 N_c^2}{2\pi^2} \zeta(3) \mathcal{V}_a(k) \end{aligned}$$

$$\begin{aligned} F(k, k') &= \left(1 + \frac{n_f}{N_c^3} \right) \frac{3(k, k')^2 - 2k^2 k'^2}{16k^2 k'^2} \left(\frac{2}{k^2} + \frac{2}{k'^2} + \frac{k^2 - k'^2}{k^2 k'^2} \ln \frac{k^2}{k'^2} \right) \\ &- \left[3 + \left(1 + \frac{n_f}{N_c^3} \right) \left(1 - \frac{(k^2 + k'^2)^2}{8k^2 k'^2} + \frac{3k^4 + 3k'^4 - 2k^2 k'^2}{16k^4 k'^4} (k, k')^2 \right) \right] \int_0^\infty \frac{dt}{k^2 + t^2 k'^2} \ln \frac{1+t}{|1-t|}, \end{aligned}$$

$$\begin{aligned} \Phi(k, k') &= \frac{(k^2 - k'^2)}{(k - k')^2 (k + k')^2} \left[\ln \frac{k^2}{k'^2} \ln \frac{k^2 k'^2 (k - k')^4}{(k^2 + k'^2)^4} \right. \\ &\left. + 2 \text{Li}_2 \left(-\frac{k'^2}{k^2} \right) - 2 \text{Li}_2 \left(-\frac{k^2}{k'^2} \right) \right] - \left(1 - \frac{(k^2 - k'^2)^2}{(k - k')^2 (k + k')^2} \right) \left[\int_0^1 - \int_1^\infty \right] \frac{du}{(k - k'u)^2} \ln \frac{u^2 k'^2}{k^2} \end{aligned}$$

Agrees with NLO BFKL

Argument of coupling constant

$$\frac{d}{d\eta} \hat{\mathcal{U}}(z_1, z_2) =$$


$$\frac{\alpha_s(?_\perp) N_c}{2\pi^2} \int dz_3 \frac{z_{12}^2}{z_{13}^2 z_{23}^2} \left\{ \hat{\mathcal{U}}(z_1, z_3) + \hat{\mathcal{U}}(z_3, z_2) - \hat{\mathcal{U}}(z_1, z_2) - \hat{\mathcal{U}}(z_1, z_3) \hat{\mathcal{U}}(z_3, z_2) \right\}$$

Argument of coupling constant

$$\frac{d}{d\eta} \hat{\mathcal{U}}(z_1, z_2) =$$

$$\frac{\alpha_s(?_\perp) N_c}{2\pi^2} \int dz_3 \frac{z_{12}^2}{z_{13}^2 z_{23}^2} \left\{ \hat{\mathcal{U}}(z_1, z_3) + \hat{\mathcal{U}}(z_3, z_2) - \hat{\mathcal{U}}(z_1, z_2) - \hat{\mathcal{U}}(z_1, z_3) \hat{\mathcal{U}}(z_3, z_2) \right\}$$

Renormalon-based approach: summation of quark bubbles



$$-\frac{2}{3} n_f \rightarrow b = \frac{11}{3} N_c - \frac{2}{3} n_f$$

Argument of coupling constant

$$\begin{aligned} \frac{d}{d\eta} \text{Tr}\{\hat{U}_{z_1} \hat{U}_{z_2}^\dagger\} &= \frac{\alpha_s(z_{12}^2)}{2\pi^2} \int d^2 z [\text{Tr}\{\hat{U}_{z_1} \hat{U}_{z_3}^\dagger\} \text{Tr}\{\hat{U}_{z_3} \hat{U}_{z_2}^\dagger\} - N_c \text{Tr}\{\hat{U}_{z_1} \hat{U}_{z_2}^\dagger\}] \\ &\times \left[\frac{z_{12}^2}{z_{13}^2 z_{23}^2} + \frac{1}{z_{13}^2} \left(\frac{\alpha_s(z_{13}^2)}{\alpha_s(z_{23}^2)} - 1 \right) + \frac{1}{z_{23}^2} \left(\frac{\alpha_s(z_{23}^2)}{\alpha_s(z_{13}^2)} - 1 \right) \right] + \dots \end{aligned}$$

I.B.; Yu. Kovchegov and H. Weigert (2006)

When the sizes of the dipoles are very different the kernel reduces to:

$$\frac{\alpha_s(z_{12}^2)}{2\pi^2} \frac{z_{12}^2}{z_{13}^2 z_{23}^2} \quad |z_{12}| \ll |z_{13}|, |z_{23}|$$

$$\frac{\alpha_s(z_{13}^2)}{2\pi^2 z_{13}^2} \quad |z_{13}| \ll |z_{12}|, |z_{23}|$$

$$\frac{\alpha_s(z_{23}^2)}{2\pi^2 z_{23}^2} \quad |z_{23}| \ll |z_{12}|, |z_{13}|$$

⇒ the argument of the coupling constant is given by the size of the smallest dipole.

Conclusions

- High-energy operator expansion in color dipoles works at the NLO level.

- High-energy operator expansion in color dipoles works at the NLO level.
- The NLO BK kernel in for the evolution of conformal composite dipoles in $\mathcal{N} = 4$ SYM is Möbius invariant in the transverse plane.
- The NLO BK kernel agrees with NLO BFKL equation.
- The correlation function of four Z^2 operators is calculated at the NLO order.
- NLO photon impact factor is calculated.

Outlook: relation to conformal light-ray operators

Gluon parton density $\mathcal{D}(x_B, \mu^2)$ is proportional to matrix element of the light-ray operator

$$\mathcal{O}(x_B, \mu^2) = \int d\lambda e^{i\lambda x_B} \text{Tr}\{G_{+i}(\lambda e^+) [\lambda e^+, 0] G_{+i}(0) [0, \lambda e^+] \}^\mu$$

Conformal light-ray operator \mathcal{O}_j (j - conformal spin in $SL(2, R)$ group)

$$\mathcal{O}_j^\mu = \int d\lambda \lambda^{1-j} \text{Tr}\{G_{+i}(\lambda e^+) [\lambda e^+, 0] G_{+i}(0) [0, \lambda e^+] \}^\mu$$

Anomalous dimension

$$\mu \frac{d}{d\mu} \mathcal{O}_j = \gamma_j(\alpha_s) \mathcal{O}_j$$

At $j = n$ γ_n is an anomalous dimension of the local twist-2 operator

$$G^{+i} (D^+)^{n-2} G_i^+$$

Outlook: relation to conformal light-ray operators

Gluon parton density $\mathcal{D}(x_B, \mu^2)$ is proportional to matrix element of the light-ray operator

$$\mathcal{O}(x_B, \mu^2) = \int d\lambda e^{i\lambda x_B} \text{Tr}\{G_{+i}(\lambda e^+) [\lambda e^+, 0] G_{+i}(0) [0, \lambda e^+]\}^\mu$$

Conformal light-ray operator \mathcal{O}_j (j - conformal spin in $SL(2, R)$ group)

$$\mathcal{O}_j^\mu = \int d\lambda \lambda^{1-j} \text{Tr}\{G_{+i}(\lambda e^+) [\lambda e^+, 0] G_{+i}(0) [0, \lambda e^+]\}^\mu$$

Anomalous dimension

$$\mu \frac{d}{d\mu} \mathcal{O}_j = \gamma_j(\alpha_s) \mathcal{O}_j$$

At $j = n$ γ_n is an anomalous dimension of the local twist-2 operator

$$G^{+i} (D^+)^{n-2} G_i^+$$

Expansion of conformal dipoles in conformal light-ray operators - ?

Outlook: relation to conformal light-ray operators

In the leading order relation this expansion is trivial: x_\perp^2 is the normalization point of gluon light-ray operator and $x_B = e^{-\eta}$:

$$\begin{aligned}\text{Tr}\{\partial_i U_x \partial^i U_0\}^\eta &= \mathcal{D}_{x_B=e^{-\eta}}^{\mu^2=x_\perp^{-2}} + O(x_\perp^2) = \int_{\frac{1}{2}-i\infty}^{\frac{1}{2}+i\infty} \frac{dj}{2\pi i} \frac{\Gamma(j-1)}{x_B^{j-1}} (x_\perp^2 \mu^2)^{-\gamma_j} \mathcal{O}_j^{\mu^2} \\ &= \int_{-\frac{1}{2}-i\infty}^{-\frac{1}{2}+i\infty} \frac{d\omega}{2\pi i} \Gamma(\omega) e^{\omega\eta} (x_\perp^2 \mu^2)^{-\gamma_\omega} \mathcal{O}_\omega^{\mu^2}\end{aligned}$$

This should be compared to LO rapidity evolution of color dipole
 $\omega_{\gamma=\frac{1}{2}+i\nu} = \omega(\nu)$ - pomeron intercept)

$$\text{Tr}\{\partial_i U_x \partial^i U_0\}^\eta = \int_{\frac{1}{2}-i\infty}^{\frac{1}{2}+i\infty} \frac{d\gamma}{2\pi i} e^{\omega_\gamma(\eta-\eta_0)} (x_\perp^2 \mu^2)^{-\gamma} \int d^2 z (z_\perp^2)^{1-\gamma} \mathcal{U}(z_\perp)^{\eta_0}$$

Outlook: relation to conformal light-ray operators

In the leading order relation this expansion is trivial: x_\perp^2 is the normalization point of gluon light-ray operator and $x_B = e^{-\eta}$:

$$\begin{aligned}\text{Tr}\{\partial_i U_x \partial^i U_0\}^\eta &= \mathcal{D}_{x_B=e^{-\eta}}^{\mu^2=x_\perp^{-2}} + O(x_\perp^2) = \int_{\frac{1}{2}-i\infty}^{\frac{1}{2}+i\infty} \frac{dj}{2\pi i} \frac{\Gamma(j-1)}{x_B^{j-1}} (x_\perp^2 \mu^2)^{-\gamma_j} \mathcal{O}_j^{\mu^2} \\ &= \int_{-\frac{1}{2}-i\infty}^{-\frac{1}{2}+i\infty} \frac{d\omega}{2\pi i} \Gamma(\omega) e^{\omega\eta} (x_\perp^2 \mu^2)^{-\gamma_\omega} \mathcal{O}_\omega^{\mu^2}\end{aligned}$$

This should be compared to LO rapidity evolution of color dipole
 $\omega_{\gamma=\frac{1}{2}+i\nu} = \omega(\nu)$ - pomeron intercept)

$$\text{Tr}\{\partial_i U_x \partial^i U_0\}^\eta = \int_{\frac{1}{2}-i\infty}^{\frac{1}{2}+i\infty} \frac{d\gamma}{2\pi i} e^{\omega_\gamma(\eta-\eta_0)} (x_\perp^2 \mu^2)^{-\gamma} \int d^2 z (z_\perp^2)^{1-\gamma} \mathcal{U}(z_\perp)^{\eta_0}$$

⇒

$$\omega = \omega(\gamma, \alpha_s) \Leftrightarrow \gamma = \gamma(\omega, \alpha_s) \simeq \sum \frac{\alpha_s^n}{\omega^n} = \frac{\alpha_s}{\omega} + \frac{\alpha_s^3}{\omega^3} + \dots$$

BFKL gives the anomalous dimensions in all orders as $\omega \rightarrow 0$ which corresponds to the non-physical point $j = n = 1$ for γ_n of local operators

Outlook: relation to conformal light-ray operators

In the NLO the expansion of conformal dipoles in conformal light-ray operators is not straightforward due to mismatch of *UV* and rapidity regularizations.

$$\tilde{\omega}(\alpha_s, \gamma) = \omega\left(\alpha_s, \gamma + \frac{1}{2}\omega\right) \Rightarrow \gamma = \gamma(\tilde{\omega}, \alpha_s)$$

$\omega(\alpha_s, \gamma)$ is the pomeron intercept which enters stands in the formula for the amplitude in terms of conformal ratios.

$\tilde{\omega}(\alpha_s, \gamma)$ determines anomalous dimensions of conformal light-ray operators.

The difficulty is probably due to the fact that conformal dipoles are invariant under $SL(2, C)$ and light-ray operators under $SL(2, R)$

Gluon TMDs may serve as a bridge between these two approaches

Outlook: rapidity evolution of gluon TMD's. $\mathcal{N} = 4$ for simplicity.

$$\text{Gluon TMD (without subtractions)} : \quad D(x_B, \eta, k_\perp, \mu^2) \sim \int d^2 k_\perp e^{ik_\perp \cdot z_\perp} \\ \times \int du dv e^{i(u-v)x_B \frac{s}{2}} \langle [-\infty, u]_z G_{+i}(z_\perp + up_1)[u, -\infty]_z [-\infty, u]_0 G_{+i}(vp_1)[u, -\infty]_0 \rangle^\eta$$

Two evolutions: η and $\mu^2 \Rightarrow$ double logs.

At $x_B = 0$ we get ($U_i \equiv U_i^\dagger i \partial_i U$)

$$D(x_B, \eta, k_\perp) = \mathcal{V}^\eta(k) = \int d^2 k_\perp e^{ik_\perp \cdot z_\perp} \langle \text{Tr}\{U_i(z_\perp)U_i(0_\perp)\} \rangle^\eta \\ = \int d^2 k_\perp e^{ik_\perp \cdot z_\perp} \int du dv \langle [-\infty, u]_z G_{+i}(z_\perp + up_1)[u, -\infty]_z [-\infty, u]_0 G_{+i}(vp_1)[u, -\infty] \rangle^\eta$$

No μ dependence (dipole amplitudes are UV finite) \Rightarrow rapidity evolution only.

Evolution of gluon TMD probably depends on the interplay between x_B and η