SIDIS Program at 12 GeV Jefferson Lab with SoLID

Radiative Corrections 2016 @ JLab Newport News, VA
May 16-19th 2016

Tianbo Liu
(SoLID Collaboration)
Duke University and Duke Kunshan University

Thanks Xuefei Yan for inputs with some slides

Duke UNIVERSITY

昆山杜克大学 DUKE KUNSHAN UNIVERSITY

Overview of SoLID

Solenoidal **L**arge **I**ntensity **D**evice



Full exploitation of JLab 12 GeV upgrade

- The capability to handle high luminosity $L \sim 10^{37\text{-}39}\ cm^{-2}\ s^{-1}$.
- Large acceptance with full 2π azimuthal angle coverage.

  Polar angle coverage: 8°~24°. (SIDIS)

Five highly rated approved experiments

- Nucleon structure: three-dimensional imaging of the nucleon in momentum space in valence quark region.
  E12-10-006, E12-11-007, E12-11-008 (SIDIS)
- Fundamental symmetries: new physics in the 10~20 TeV region, complementary to the reach of LHC.
  E12-10-007 (PVDIS)
- J/ψ: probe the color field in the nucleon, access to QCD conformal anomaly.
  E12-12-006 (J/ψ)

SIDIS @ SoLID

Approved SIDIS experiments 11/8.8 GeV

E12-10-006: Single Spin Asymmetry on Transversely polarized ^{3}He, 90 days.

E12-11-007: Single and Double Spin Asymmetry on Longitudinally polarized ^{3}He, 35 days.

E12-10-008: Single Spin Asymmetry on Transversely polarized proton (NH_3), 120 days.

Run group:
E12-10-006A, E12-11-108A
Dihadron process
Target single spin asymmetry A_y



$P_n = 86\%$, $P_p = -2.8\%$

Total about 1400 ^{3}He bins
and 650 proton bins in x, z, Q^2, P_T

High statistics (example)



Projected data of E12-10-006

PVDIS @ SoLID

Approved PVDIS experiment @ 11 GeV

E12-10-007: Parity violating asymmetry in DIS with LH_2 and LD_2 targets.



SoLID projected and final Qweak results



J/ψ @ SoLID

Approved J/ψ near threshold production @ 11 GeV

E12-12-006: measure J/ψ near threshold production cross section on proton (LH_2).

Run group:
E12-12-006A Timelike Compton Scattering (TCS).



Imaginary part: total cross section through the optical theorem.
Real part: contains the conformal anomaly.



Proton mass:



$\overline{MS}$ and $\mu = 2$ GeV

H. Gao *et al.*, The Universe 3, no.2, 18 (2015).

Probing the strong color field in the nucleon, conformal anomaly (has impact on understanding of the proton mass).

Nucleon Spin Decomposition



Proton spin puzzle

$$\Delta\Sigma = \Delta u + \Delta d + \Delta s \sim 0.3$$

Quark spin only contributes a small fraction to nucleon spin.

J. Ashman *et al.*, PLB 206, 364 (1988); NP B328, 1 (1989).

Spin decomposition

$$J = \frac{1}{2}\Delta\Sigma + \Delta G + L_q + L_g$$



JAM Collaboration, PR D 93, 074005 (2016).



K.-F. Liu, NP A928, 99 (2014).

Access to $L_{q/g}$

It is necessary to have transverse information.

Coordinate space: GPDs
Momentum space: TMDs

3*D* imaging of the nucleon.

Unified View of Nucleon Structure



Unified View of Nucleon Structure



Structure Functions

SIDIS differential cross section

18 structure functions $F(x, z, Q^2, P_T)$,
model independent. (one photon exchange approximation)



[Diehl&Sapeta EPJC2005]

$$
\begin{aligned}
&\frac{d\sigma}{dx\,dy\,dz\,dP_T^2\,d\phi_h\,d\phi_S} \\
&= \frac{\alpha^2}{xyQ^2}\frac{y^2}{2(1-\epsilon)}\left(1+\frac{\gamma^2}{2x}\right) \\
&\times \Big\{ F_{UU,T} + \epsilon F_{UU,L} + \sqrt{2\epsilon(1+\epsilon)}F_{UU}^{\cos\phi_h}\cos\phi_h + \epsilon F_{UU}^{\cos 2\phi_h}\cos 2\phi_h + \lambda_e\sqrt{2\epsilon(1-\epsilon)}F_{LU}^{\sin\phi_h}\sin\phi_h \\
&+ S_L\left[\sqrt{2\epsilon(1+\epsilon)}F_{UL}^{\sin\phi_h}\sin\phi_h + \epsilon F_{UL}^{\sin 2\phi_h}\sin 2\phi_h\right] + \lambda_e S_L\left[\sqrt{1-\epsilon^2}F_{LL} + \sqrt{2\epsilon(1-\epsilon)}F_{LL}^{\cos\phi_h}\cos\phi_h\right] \\
&+ S_T\Big[\left(F_{UT,T}^{\sin(\phi_h-\phi_S)} + \epsilon F_{UT,L}^{\sin(\phi_h-\phi_S)}\right)\sin(\phi_h-\phi_S) + \epsilon F_{UT}^{\sin(\phi_h+\phi_S)}\sin(\phi_h+\phi_S) + \epsilon F_{UT}^{\sin(3\phi_h-\phi_S)}\sin(3\phi_h-\phi_S) \\
&\qquad + \sqrt{2\epsilon(1+\epsilon)}F_{UT}^{\sin\phi_S}\sin\phi_S + \sqrt{2\epsilon(1+\epsilon)}F_{UT}^{\sin(2\phi_h-\phi_S)}\sin(2\phi_h-\phi_S)\Big] \\
&+ \lambda_e S_T\Big[\sqrt{1-\epsilon^2}F_{LT}^{\cos(\phi_h-\phi_S)}\cos(\phi_h-\phi_S) \\
&\qquad + \sqrt{2\epsilon(1-\epsilon)}F_{LT}^{\cos\phi_S}\cos\phi_S + \sqrt{2\epsilon(1-\epsilon)}F_{LT}^{\cos(2\phi_h-\phi_S)}\cos(2\phi_h-\phi_S)\Big]\Big\}
\end{aligned}
$$

SoLID:
4D bins in (x, z, Q^2, P_T)

In parton model, $F(x, z, Q^2, P_T)$s are expressed as the convolution of TMDs.

Leading Twist TMDs

Nucleon Polarization	Quark Polarization: U	Quark Polarization: L	Quark Polarization: T
U	f_1  **unpolarized**		$h_1^\perp$  **Boer-Mulders**
L		g_{1L}  **helicity**	$h_{1L}^\perp$  **longi-transversity (worm-gear)**
T	$f_{1T}^\perp$  **Sivers**	g_{1T}  **trans-helicity (worm-gear)**	h_1  **transversity** $h_{1T}^\perp$  **pretzelosity**

Unpolarized Quark in $p\uparrow$

$$f_{q/p\uparrow}(x, \mathbf{k}_\perp) = f_1^q(x, k_\perp) - f_{1T}^{\perp q}(x, k_\perp)\frac{\hat{\mathbf{P}} \times \mathbf{k}_\perp \cdot \mathbf{S}}{M}$$

Sivers distribution



naively time-reversal odd.

$$f_{1T}^{\perp q}(x, k_\perp)\Big|_{\text{SIDIS}} = -f_{1T}^{\perp q}(x, k_\perp)\Big|_{\text{DY}}$$

Measurement in SIDIS

Single spin asymmetry (Sivers asymmetry)

$$A_{UT}^{\sin(\phi_h - \phi_S)} \sim f_{1T}^{\perp}(x, k_\perp) \bigotimes D_1(z, p_\perp)$$



6 GeV JLab E06-010, X. Qian *et al.*, PRL 107, 072003 (2011).

Model Calculation





Bacchetta, Conti, Radici
PR D 78, 074010 (2008).

Transverse Spin Structure

Transversity

h_1  (Collinear & TMD)

A transverse counter part to the longitudinal spin structure: helicity g_{1L}

They are NOT the same due to relativity.

Chiral-odd

Unique for the quarks.
No mixing with gluons.
Simpler evolution effect.

NOT accessible via inclusive DIS process.
Must couple to another chiral-odd function.
(*e.g.* Collins function $H_1^{\perp}$)

Measured via
SIDIS (E12-10-006, E12-11-008), Drell-Yan
Di-hadron (approved as run group with E12-10-006)

Measurement in SIDIS

Single spin asymmetry
(Collins asymmetry)

$$A_{UT}^{\sin(\phi_h+\phi_S)} \sim h_1(x,k_\perp) \bigotimes H_1^{\perp}(z,p_\perp)$$

$H_1^{\perp}(z,p_\perp)$ Collins fragmentation function



6 GeV JLab E06-010, X. Qian *et al.*, PRL 107, 072003 (2011).

Soffer's Inequality

Soffer's bound

$$|h_1(x)| \leq \frac{1}{2}\left[f_1(x) + g_{1L}(x)\right]$$

Derived by using the positivity constraint on the forward scattering helicity amplitude.

Global fits of transversity



M. Anselmino *et al.*, PR D 92, 114023 (2015).



Z.-B. Kang *et al.*, PR D 93, 014009 (2016).

Test Soffer's inequality @ SoLID

Pretzelosity

Pretzelosity distribution



Chiral-odd. NO gluon analogy.
Interference of light-front wave functions differing by $\Delta L = 2$.
Measuring the difference between helicity and transversity, and hence relativistic effects. (spherically symmetric models)

Relation to OAM (canonical)

$$L_z^q = -\int \mathrm{d}x \mathrm{d}^2\mathbf{k}_\perp \frac{\mathbf{k}_\perp^2}{2M^2} h_{1T}^{\perp q}(x, k_\perp) = -\int \mathrm{d}x h_{1T}^{\perp(1)q}(x)$$

(model dependent)

Measurement in SIDIS

Single spin asymmetry

$$A_{UT}^{\sin(3\phi_h - \phi_S)} \sim h_{1T}^{\perp}(x, k_\perp) \otimes H_1^{\perp}(z, p_\perp)$$

A global fit to 175 data from COMPASS, HERMES, and JLab found comparable with null signal hypothesis at 72% C.L..

C. Lefky, A. Prokudin, PR D 91, 034010 (2015).



6 GeV JLab E06-010, Y. Zhang *et al.*, PR C 90, 055209 (2014).

Radiative Correction @ 6 GeV

Transversity experiment at Hall A: E06-010



by Xuefei Yan

HAPRAD built in MC.

I. Akushvich et al., PL B 672, 35 (2009).

Radiative Correction @ SoLID

Transverse polarized ^{3}He: E12-10-006 (proposal: a simple estimation)



Estimated by Pythia with RADGEN.

RC under Development @ SoLID

- Internal radiative correction (HAPRAD, etc.) built in MC
  - internal and external RC combination
  - full end-to-end MC under development

- Cross section model adjustability: for iterative processes
  - reduce model dependence

- Capability for bin-by-bin and unfolding

- Exclusive tail: there will be data at the same kinematic for model updating

- Collaborating with theorists and other groups to develop full procedure

Present Status On TMD Extractions

Sivers



Anselmino et al, EPJA39, 89 (2009)

Transversity



Anselmino et al, PRD92, 114023 (2015)

Pretzelosity



Lefky et al, PRD91, 034010 (2015)

Collins fragmentation

Anselmino et al, PRD92, 114023 (2015)
PRD93, 034025 (2016)





SoLID Impact on Sivers

$$f_{1T}^{\perp(1)}(x) = \int \mathrm{d}^2\mathbf{k}_\perp \frac{\mathbf{k}_\perp^2}{2M^2} f_{1T}^{\perp}(x, k_\perp)$$





95% C.L.

parametrization by M. Anselmino *et al.*, EPJ A 39, 89 (2009).

SoLID projection with transversely polarized neutron and proton data.

Quark Transverse Momentum in $p\uparrow$





95% C.L.

parametrization by M. Anselmino *et al.*, EPJ A 39, 89 (2009).

SoLID projection with transversely polarized neutron and proton data.

Transverse momentum:

$$\langle \mathbf{k}_\perp \rangle = -M \int \mathrm{d}x\, f_{1T}^{\perp(1)}(x)\, (\mathbf{S} \times \hat{\mathbf{P}})$$



SoLID Impact on Transversity





95% C.L.

parametrization by M. Anselmino *et al.*, PR D 87, 094019 (2013).

SoLID projection with transversely polarized neutron and proton data.

SoLID Impact on Pretzelosity





95% C.L.

parametrization by C. Lefky *et al.*, PR D 91, 034010 (2015).

SoLID projection with transversely polarized neutron and proton data.

OAM:

$$L_z^q = -\int \mathrm{d}x\mathrm{d}^2\mathbf{k}_\perp \frac{\mathbf{k}_\perp^2}{2M^2} h_{1T}^{\perp q}(x, k_\perp) = -\int \mathrm{d}x h_{1T}^{\perp(1)q}(x)$$



Tensor Charge

Definition

$$\langle P,S|\bar{\psi}_q i\sigma^{\mu\nu}\psi_q|P,S\rangle = \delta_T q\bar{u}(P,S)i\sigma^{\mu\nu}u(P,S) \qquad \delta_T q = \int_0^1 \left[h_1^q(x) - h_1^{\bar{q}}(x)\right] \mathrm{d}x$$

A fundamental QCD quantity. Matrix element of local operators.
Moment of transversity distribution. Valence quark dominant.
Calculable in lattice QCD.

SoLID impact



Tensor Charge and Neutron EDM

Electric **D**ipole **M**oment

Tensor charge and EDM

$$d_n = \delta_T u \, d_u + \delta_T d \, d_d + \delta_T s \, d_s$$



current neutron EDM limit $|d_n| < 2.9 \times 10^{-26} \, e \cdot \mathrm{cm}$

Summary

- Lepton scattering is a powerful tool to probe the internal structure of the nucleon.
- Unprecedented precision with high luminosity and large acceptance at JLab 12-GeV with SoLID.
- SoLID-SIDIS program: multi-dimensional mapping in valence quark region with high precision.
- TMD: transverse imaging of the nucleon, access to orbital angular momentum.
- Tensor charge and neutron EDM, constraint on new physics.
- Developing full procedure for RC and physics extraction.

Thank you!

Backup

Exclusive tail @ 6 GeV

Transversity experiment at Hall A: E06-010



by Xuefei Yan

Exclusive model. R. Asaturyan et al., PR C 85, 015202 (2012).

SoLID Impact on Transversity TMD





95% C.L.

parametrization by M. Anselmino *et al.*, PR D 87, 094019 (2013).

SoLID projection with transversely polarized neutron and proton data.

What Else in SIDIS @ SoLID

Worm-gears

g_{1T}



Trans-helicity (worm-gear).
Interference of light-front wave functions differing by $\Delta L = 1$.

Measured by DSA $A_{LT}^{\cos(\phi_h - \phi_S)} \sim g_{1T}(x, k_\perp) \bigotimes D_1(z, p_\perp)$

(The other worm-gear from $A_{UL}^{\sin 2\phi_h} \sim h_{1L}^\perp(x, k_\perp) \bigotimes H_1^\perp(z, p_\perp)$.)

Subleading twist effect

Beam spin asymmetry $A_{LU}^{\sin \phi_h}$ from twist-3 TMDs.

Other subleading twist asymmetries, *e.g.* $A_{UT}^{\sin(2\phi_h - \phi_S)}$, $A_{UT}^{\sin \phi_S}$, $A_{LT}^{\cos \phi_S}$ $\cdots$

Unpolarized process

Multiplicity or differential cross section.

Cahn effect $f_1 \otimes D_1$ Boer-Mulders effect $h_1^\perp \otimes H_1^\perp$