

NLO($O(\alpha_s^2)$) SIDIS at large transverse momentum

Bowen Wang



Collaborators:

J. Gonzalez, T. Rogers, A. Signori, N. Sato


JLab
February 22 2017

Role of perturbative term

Some details on NLO calculation


SIDIS at small and large q_T

$$q_T \sim m$$




TMD Factorization

$$q_T \gtrsim Q$$



Coll. Factorization

Unified: Transverse Momentum:



Why collinear term is important


TMD term

- contains TMD PDFs and TMD FFs
- describes intrinsic structure of nucleons
- process independent

Collinear term

- Calculable in pQCD
- process dependent


Theory designed for fragmentation of the struck quark (current fragmentation)



M. Boglion, J.Collins, L.Gamberg, J.O.Gonzalez, T.C.Rogers, N.Sato, arXiv:1611.10329

Are all the data in the current region?


- Need to isolate a current region to apply TMD factorization.



M. Boglion, J.Collins, L.Gamberg, J.O.Gonzalez, T.C.Rogers, N.Sato, arXiv:1611.10329


What to do with data not in the current region?

- No problem for large enough q_T — collinear factorization



M. Boglion, J.Collins, L.Gamberg, J.O.Gonzalez, T.C.Rogers, N.Sato, arXiv:1611.10329


FO at $O(\alpha_s)$




TMD and FO cross section in SIDIS. P. Nadolsky, *etal* (1999)

$O(\alpha_s)$ and $O(\alpha_s^2)$

Perturbative cross section in SIDIS. A. Daleo, *etal* (2004)



$O(\alpha_s)$ and $O(\alpha_s^2)$



Kinematical reason: with extra radiations, it is more likely to produce low energy jets at the central rapidity region.

Calculation of hard scattering coefficients

$$\frac{d\Gamma}{dx dy dz dP_T^2} = \frac{2\pi^2 \alpha_{EM}^2}{Q^4 y(1-\epsilon)} \left[Q^2(y^2 - 2y + 2) g^{\mu\nu} W_{\mu\nu} \right. \\ \left. + 4x^2(y^2 - 6y + 6 + 4\epsilon(y-1)) p^\mu p^\nu W_{\mu\nu} \right] \quad \left(y = \frac{Q^2}{xs} \right)$$

$$g^{\mu\nu} W_{\mu\nu} = \int |M|^2 dPS_3(\overline{z, P_T^2}) \equiv \frac{d\sigma}{dz dP_T^2}$$

or change to Mandelstam's variables

$$\frac{d\sigma}{du dt} \equiv \int |M|^2 dPS_3(\overline{u, t}), \quad \frac{d\sigma}{dz dP_T^2} = \frac{d\sigma}{du dt} \left| \frac{\partial(u, t)}{\partial(z, P_T^2)} \right|$$

similar definitions are used for $p^\mu p^\nu W_{\mu\nu}$

Calculation of hard scattering coefficients

$$\frac{d\sigma}{dudt} = \int d\xi d\zeta \frac{d\hat{\sigma}}{dudt} f(\xi) d(\zeta)$$

$$\begin{aligned} \frac{d\sigma^{(1)}}{dudt} + \frac{d\sigma^{(2)}}{dudt} + \dots &= \int d\xi d\zeta \left(\frac{d\hat{\sigma}^{(1)}}{dudt} + \frac{d\hat{\sigma}^{(2)}}{dudt} + \dots \right) \\ &\quad \times (f^{(0)}(\xi) + f^{(1)}(\xi) + \dots)(d^{(0)}(\zeta) + d^{(1)}(\zeta) + \dots) \end{aligned}$$

$$\frac{d\hat{\sigma}^{(1)}}{dudt} = \frac{d\sigma^{(1)}}{dudt}$$

$$\frac{d\hat{\sigma}^{(2)}}{dudt} = \frac{d\sigma^{(2)}}{dudt} - \int d\xi \frac{d\sigma^{(1)}}{dudt} f^{(1)}(\xi) - \int d\zeta \frac{d\sigma^{(1)}}{dudt} d^{(1)}(\zeta)$$

...

Calculation of hard scattering coefficients

$$f^{(0)}(\xi) = \delta(1 - \xi)$$


$$d^{(0)}(\zeta) = \delta(1 - \zeta)$$

$$f_{qq}^{(1)\overline{MS}}(\xi) = C_F \left(\frac{1 + \xi^2}{(1 - \xi)_+} + \frac{3}{2} \delta(1 - \xi) \right) \left(-\frac{1}{\epsilon} - \ln 4\pi + \gamma_E + \ln \left(\frac{M^2}{\mu^2} \right) \right)$$


$$d_{qq}^{(1)\overline{MS}}(\zeta) = f_{qq}^{(1)\overline{MS}}(\zeta)$$

...

Calculation of hard scattering coefficients



Three particle phase space



$$PS_3 = \int \frac{d^n k_1}{(2\pi)^{n-1} 2E_1} \frac{d^n k_2}{(2\pi)^{n-1} 2E_2} \frac{d^n k_3}{(2\pi)^{n-1} 2E_3} \\ \times (2\pi)^n \delta^{(n)}(p + q - k_1 - k_2 - k_3) \\ n = 4 - 2\epsilon$$

In the CM frame of $k_2 + k_3$,

$$k_2 = E_2(1, \dots, \cos \theta_2 \sin \theta_1, \cos \theta_1)$$

$$k_3 = E_2(1, \dots, -\cos \theta_2 \sin \theta_1, -\cos \theta_1)$$

the above phase space simplifies

to

$$\int \frac{-1}{128(s + Q^2)\Gamma(1 - 2\epsilon)} \left(\frac{16\pi^2(s + Q^2)^2}{s_{23}u(st + Q^2 s_{23})} \right)^\epsilon \\ \times \sin^{1-2\epsilon} \theta_1 \sin^{-2\epsilon} \theta_2 d\theta_1 d\theta_2 du dt$$

$$s \equiv (p + q)^2$$

$$t_i \equiv (q - k_i)^2$$

$$u_i \equiv (p - k_i)^2$$


$$s_{ij} \equiv (k_i + k_j)^2$$

$$i, j = 1, 2, 3$$

$$u_1 \equiv u$$

$$t_1 \equiv t$$

Partial fraction



Need to reduce the number of Mandelstam variables in each term in the squared matrix element to less than two. Example:

$$\frac{u_2}{u_3 t_2^2 s_{13}}$$

Apply the relation $u_1 = t_2 - u_3 - s_{13}$ to get

$$\begin{aligned} \frac{u_2}{u_3 t_2^2 s_{13}} &= \frac{u_2}{u_3 t_2^2 s_{13}} \frac{t_2 - u_3 - s_{13}}{u_1} \\ &= -\frac{u_2}{s_{13} t_2^2 u_1} - \frac{u_2}{t_2^2 u_3 u_1} + \frac{u_2}{s_{13} t_2 u_3 u_1} \\ \frac{u_2}{s_{13} t_2 u_3 u_1} &= -\frac{u_2}{s_{13} t_2 u_1^2} + \frac{u_2}{s_{13} u_3 u_1^2} - \frac{u_2}{t_2 u_3 u_1^2} \end{aligned}$$

$$s \equiv (p + q)^2$$

$$t_i \equiv (q - k_i)^2$$

$$u_i \equiv (p - k_i)^2$$


$$s_{ij} \equiv (k_i + k_j)^2$$

$$i, j = 1, 2, 3$$

$$u_1 \equiv u$$

$$t_1 \equiv t$$

Partial fraction



There are six relevant variables: $t_2, t_3, u_2, u_3, s_{12}, s_{13}$. (s_{23} is constrained by the momentum conservation $s + u + t = -Q^2 + s_{23}$). Four relations between them

$$t_1 + t_2 + t_3 + s + 2Q^2 = 0$$

$$u_1 + u_2 + u_3 + s + Q^2 = 0$$

$$s_{12} = s + t_3 + u_3 + Q^2$$

$$s_{13} = s + t_2 + u_2 + Q^2$$

reduce the number of independent variables to two. Therefore any variable in numerator can be expressed in terms of the two variables in denominator.

$$s \equiv (p + q)^2$$

$$t_i \equiv (q - k_i)^2$$

$$u_i \equiv (p - k_i)^2$$

$$s_{ij} \equiv (k_i + k_j)^2$$

$$i, j = 1, 2, 3$$

$$u_1 \equiv u$$

$$t_1 \equiv t$$


Phase space integration

$$\int_0^\pi d\theta_1 \sin^{1-2\epsilon} \theta_1 \int_0^\pi d\theta_2 \sin^{-2\epsilon} \theta_2$$
$$\times \frac{1}{(1 - \cos \theta_1)^j (1 - \cos \theta_1 \cos \chi - \sin \theta_1 \cos \theta_2 \sin \chi)^l}$$
$$= 2\pi \frac{\Gamma(1 - 2\epsilon)}{\Gamma(1 - \epsilon)^2} 2^{-j-l} B(1 - \epsilon - j, 1 - \epsilon - l) {}_2F_1\left(j, l, 1 - \epsilon, \cos^2 \frac{\chi}{2}\right)$$

(Van Neervan 1985)


If the variables in integrand depend on momentum of massive particles, e.g. t_2 or t_3 , the above identity cannot be used.

Calculation of FO term at NLO (e.g. $\gamma q \rightarrow qgg$)




Status

$$\frac{d\hat{\sigma}^{(1)}}{dudt} :$$




$$\frac{d\hat{\sigma}^{(2)}}{dudt} :$$




almost + ...

+




+




+ ...

-



✓ -



✓ + ...