

94-005

# Determination of the $N\Delta$ Axial Vector Transition

## Form Factor $G_A^{N\Delta}$ from the

### $ep \rightarrow e'\Delta^{++}\pi^-$ Reaction

P. Degtyarenko, L. Elouadrhiri, T. Hemmert, R. Hicks, R. Miskimen,  
G. Peterson, K. Wang

*University of Massachusetts, Amherst, MA 01003*

D. Doughty, D. Heddle, Zh. Li

*Christopher Newport University, Newport News, Va. 23606*

W. Brooks, V. Burkert, D. Cords, B. Mecking, B. Niczyporuk,  
E.S. Smith, A. Yegneswaran

*CEBAF, Newport News, Va. 23606*

N. Bianchi, G.P. Capitani, E. De Sanctis, P. Levi Sandri,  
V. Muccifora, E. Poli, A.R. Reolon, P. Rossi

*INFN, Frascati, Italy*

M. Anghinolfi, P. Corvisiero, G. Gervino, L. Mazzaschi, V.I. Mokeev,  
G. Ricco, M. Ripani, M. Sanzone, M. Taiuti, A. Zucchiatti

*INFN, Genova, Italy*

P.Y. Bertin


*Universite Blaise-Pascal, Clermont-Ferrand, France*

and the

CLAS Collaboration

Spokespersons: L. Elouadrhiri (contact), D. Heddle, R. Hicks, Zh. Li


- Goal of hadronic physics at intermediate energies
  - Describe the structure of the nucleon in terms of its fundamental constituents.
- Extended experimental program at CEBAF to study the electromagnetic structure of the hadronic current
  - Elastic form factors of the Nucleon  $eN \rightarrow eN$
  - Resonance transition form factors  $eN \rightarrow eN^*, e\Delta^*$
- Axial structure of hadronic current of similar importance
  - Independent information about nucleon structure
  - Test QCD motivated models, QCD lattice calculations



$$\begin{aligned}
 \langle \Delta^{++}(\vec{p}') | \underline{A}_\mu^\Delta | p(\vec{p}) \rangle = & \bar{\Delta}^{++\nu}(\vec{p}') \left[ \underline{d}_1(q^2) g_{\mu\nu} + \frac{d_2(q^2)}{M^2} P^\alpha (q_\alpha g_{\mu\nu} - q_\nu g_{\alpha\mu}) \right. \\
 & \left. - \frac{d_3(q^2)}{M^2} p_\nu q_\mu + i \frac{d_4(q^2)}{M^2} \epsilon_{\mu\nu\alpha\beta} P^\alpha q^\beta \gamma_5 \right] u(\vec{p})
 \end{aligned}$$

**Weak interaction**

$\nu, \bar{\nu}$  scattering



**Leptonic current**

$$\gamma^\mu (1 + \gamma^5)$$

**Direct access to the axial part  $A_\mu$**




**But: low statistics requires model dependent fit, e.g.**

$$G_A^{N\Delta}(Q^2) = \frac{G_A^{N\Delta}(0)}{(1 + M_A^{*2}/Q^2)^2}$$

**e.m. interaction**

**pion electroproduction at the threshold**



**Leptonic current**

$$\gamma^\mu$$

**Access to  $A_\mu$  only if pion production**

**Chiral symmetry and PCAC**



$$G_A(Q^2), G_P(Q^2), G_A^{N\Delta}(Q^2)$$









Figure 3: Processes contributing to  $\Delta\pi$  production off the nucleon<sup>7</sup>

$$Q^2 = 0.6 \text{ GeV}^2$$





# DESY Data



## Projected Results for CLAS Experiment

- Cross section and Event generator
  - Gauge invariant formalism of Stichel and Scholz
  - Form factor parametrization with  $G_A^\Delta = (1 + Q^2/M_\rho^2)^{-1}$ .
- Experimental Method
  - Detect  $e'$ ,  $p$ ,  $\pi^+$  in CLAS
  - Identify  $ep \rightarrow e'\Delta^{++}\pi^-$  using missing mass method
- Analysis Method
  - Identify  $\Delta^{++}$  in invariant mass distribution of  $p\pi^+$
  - Separate  $\Delta^{++}\pi^-$  from  $\Delta^0\pi^+$  and  $(p\pi^+\pi^-)_{ps}$  in fit to Dalitz density distributions (tested in trial analysis)
  - Test  $\pi^-$  angular distribution for different cuts on  $W_{\Delta\pi}^{max}$  and  $p_{\pi^-}^{max}$
  - Identify 'contact' term in angular distribution (s-wave)
- Statistics and Systematic Error on  $(G_A^\Delta)^2$ 
  - Statistics: 0.5% - 5% (high  $Q^2$ )     $\mathcal{L} = (3-10) 10^{33} \text{ cm}^{-2} \text{ sec}^{-1}$
  - Acceptance: 3% - 5%, Analysis: 2% - 5%
  - Total estimated error in  $G_A^\Delta$ : 2.2% to 7%.





CLAS


SDR Event Reconstruction



$$E = 4 \text{ GeV}, \quad 1.5 < Q^2 < 3 \text{ (GeV}^2\text{)}, \quad W < 1.5 \text{ GeV}$$


Reconstructed missing mass to p and  $\pi^+$







Parameter	Input value	Fitted value
$a_{\Delta^{++}}$	30000	$30218 \pm 625$
$a_{\Delta^0}$	6000	$5720 \pm 558$
$a_{p_s}$	15000	$15525 \pm 1018$



$Q = 1.5 \text{ GeV}^2$ ;  $W = 1.4 \text{ GeV}$





# Projected Errors for CLAS Experiment




# Projected Error for CLAS Experiment



# Projected Error for CLAS Experiment



# Projected Errors for CLAS Experiment



## CONCLUSIONS

- Cross section measurements of  $ep \rightarrow e'\Delta^{++}\pi^-$  near threshold allow determination of the axial vector transition form factor  $G_A^{N\Delta}$ 
  - data on  $ep \rightarrow e'\Delta^{++}\pi^-$  of unprecedented quality
  - high statistics  $\rightarrow$  measurement close to threshold where axial vector contribution dominates the cross section
  - large range in  $Q^2$
- Opens a window for new physics to be explored at CEBAF
- Tests of Chiral Perturbation Theory at low  $Q^2$

### Beam time request

Target	Energy	time	comment
H <sub>2</sub> gas	E = 1.6 GeV	150 hrs	parallel to $N^*$ program
H <sub>2</sub> gas	E = 2.4 GeV	250 hrs	parallel to $N^*$ program
H <sub>2</sub> gas	E = 4.0 GeV	800 hrs	parallel to $N^*$ program
empty	1.6, 2.4, 4.0 GeV	72 hrs	<u>new beam time</u>