

FROST Experiment

Helicity Amplitude and EM coupling constants
in Exclusive Pion Photo-production


Hideko Iwamoto

The George Washington University

HUGS 2008
June 19th, 2008

Outline

1. *Motivation*
2. *Observables*
3. *Helicity Amplitudes*
4. *g_9 Experiment*
5. *Polarized Beam and Target*
6. *Electromagnetic Coupling Constants*
7. *Summary*




Baryon Resonances

Missing resonances

Parameters (mass, width, and couplings to various decay modes) are poorly known



Many broad overlapping resonances

$I = \frac{1}{2}$



Overlap of Partial Waves

To determine each amplitude
of partial wave at resonance regions
--> need to determine the strength
of contribution of each partial wave



$d\sigma/d\Omega$:
 unpolarized observable
 Σ, T, P :
 single polarized observables
 G, H :
 double polarized observables

Observables


Complete set of observables for helicity amplitudes -->8 experiments

Photon beam		Target			Recoil			Target + Recoil			
		x	y	z	x'	y'	z'	x'	x'	z'	z'
unpolarized	σ_0		T			P		Tx'	-Lx'	Tz'	Lz'
linearly	$-\Sigma$	H	(-P)	-G	O_x	(-T)	O_z	(-Lz')	(Tz')	(-Lx')	(-Tx')
circularly		F		-E	$-Cx'$		$-Cz'$				

For Beam and Target double polarization (fixed energy and fixed angle)

$$\frac{d\sigma}{d\Omega} = \left(\frac{d\sigma}{d\Omega}\right)_{un} \left(1 - P_L \Sigma \cos(2\phi) + P_x \{ P_c F - P_L H \sin(2\phi) \} \right. \\ \left. - P_y \{ -T + P_L P \cos(2\phi) \} - P_z \{ P_c E - P_L G \sin(2\phi) \} \right)$$

- $P_{x,y}$: transversely polarized target
- P_z : longitudinally polarized target
- $P_L(\phi)$: linearly polarized beam
at an angle Φ to the reaction plane
- P_c : right circularly polarized beam




Helicity Amplitudes and Observables for $\gamma p \rightarrow \pi \mathcal{N}$

$$\frac{d\sigma(\theta)}{d\Omega} = \frac{q}{k} \sum_{spins} |A_{fi}|^2 = \frac{q}{k} \sum_{i=1}^4 |H_i|^2$$

	λ_i	λ_f
H_1	+3/2	+1/2
H_2	+1/2	+1/2
H_3	+3/2	-1/2
H_4	+1/2	-1/2

λ_i : total initial helicity
 $+1 \pm 1/2 = +1/2, +3/2$
 λ_f : total final helicity
 $+1/2, -1/2$

Observables	Helicity representation
$(d\sigma/d\Omega)_{un}$	$1/2(H_1 ^2 + H_2 ^2 + H_3 ^2 + H_4 ^2)$
Σ	$\text{Re}(-H_1 H_4^* + H_2 H_3^*)$
T	$\text{Im}(H_1 H_2^* + H_3 H_4^*)$
P	$\text{Im}(-H_1 H_3^* - H_2 H_4^*)$
G	$\text{Im}(H_1 H_4^* - H_3 H_2^*)$
H	$\text{Im}(-H_2 H_4^* + H_1 H_3^*)$
E	$1/2(H_1 ^2 - H_2 ^2 + H_3 ^2 - H_4 ^2)$
F	$\text{Re}(-H_2 H_1^* - H_4 H_3^*)$







center-of-momentum frame

G9a Experiment at JLab

Experiments at Jlab

Photon beam		Proton Target		
		X	Y	Z
unpolarized	σ_0		T	
linearly	$-\Sigma$	H	(-P)	-G
circularly		F		-E

	g1	E_γ 0.5 ~ 2.9 GeV
	g8	0.9 ~ 2.1 GeV
	g9a	0.5 ~ 2.4 GeV
	g9b	

g1 : unpolarized beam and target --> σ_0


g8 : linearly polarized beam --> Σ

**g9a : linearly polarized beam and longitudinally polarized target
circularly polarized beam and longitudinally polarized target**

g9b : linearly polarized beam and transversely polarized target

(2010) circularly polarized beam and transversely polarized target

CLAS Detector




Polarized Target

Butanol (C_4H_9OH)

High polarizations (longitudinal)
Long spin-relaxation times
Short re-polarization time
Holding mode (0.5T)

Design goal
80%
500hours
few hours
< 50mK

Result
82 ~85%
1600~2800 hours
few hours
28 ~ 30mK



Circularly Polarized Photon Beam



Bremsstrahlung

linearly polarized electron beam

use radiator of gold

depend on polarization of initial electron beam

and the ratio of photon energy to the initial electron energy



$$P(\gamma) = P(e) \frac{4x - x^2}{4 - 4x + 3x^2}$$

$$x = \frac{k}{\epsilon_1} = \frac{(\text{photon energy})}{(\text{incident electron energy})}$$

H. Olson, L.C. Maximon

Phys. Rev. 114 p887 (1959)

Amplitudes from \mathcal{N}^* and Δ^* for direct resonance term

Example S_{11} (Spin $1/2$, Isospin $1/2$)

Lagrangians

$$L_{piNN^*} = \frac{-H}{m} [\bar{N} \gamma_\alpha \tau_i N^* + \bar{N}^* \gamma_\alpha \tau_i N] \partial^\alpha \pi_i$$

$$L_{em} = \frac{-F^{\alpha\beta}}{4M} [\bar{N} \sigma_{\alpha\beta} \gamma_5 (G_S + \tau_3 G_V) N^* - \bar{N}^* \sigma_{\alpha\beta} \gamma_5 (G_S + \tau_3 G_V) N]$$

$G_S (G_p + G_n), G_V (G_p - G_n)$: EM coupling constants ($Q^2=0$)

Propagator (spin $1/2$ resonances)

$$G(P^2, P) = \frac{\not{P} + M^*}{P^2 - M^{*2} + i M^* \Gamma(P^2)}$$


M: nucleon mass

$P^2 = M^{*2}$ (for $Q^2=0$)

Calculated amplitude (in the case of initial and final particles are proton)

$$A_f = \frac{G_p}{2M} \frac{H}{m} \bar{u}_2 q G (\not{\epsilon} k - k \not{\epsilon}) \gamma u_1$$

H : Strong coupling constant ($Q^2=0$)



Electromagnetic Coupling Constants

Partial Wave Analysis (PWA)

--> determine resonance couplings, $A_{\lambda}^I(I : isospin, \lambda : spin)$

$$A_{\lambda}^I \propto \frac{iB_{\lambda}^I \cdot \Gamma}{\sqrt{\Gamma_{\pi}}}$$

B_{λ}^I : *Imaginary Part of H_i (helicity amplitudes)*

Γ : *Full Width*

Γ_{π} : *Partial Width for the decay $N^* \rightarrow \pi N$*

The case of S_{11} (in the case of initial and final particles are proton)

$$A_{1/2}^p(S_{11}) = \frac{-e G_p}{\sqrt{2} M} \sqrt{\frac{k}{M(E_1 + M)}} (M + M^*)$$

--> Determine the coupling constant G_p

Summery

1. Double polarized measurement is important
2. g9a experiment
 - > double polarization, beam and target
 - > part of a series of experiments to determine the invariant helicity amplitudes and electromagnetic coupling constants