




JOHANNES
GUTENBERG
UNIVERSITÄT
MAINZ



Hypernuclei at MAMI-C and PANDA


- ▶ Introduction
- ▶ Hypernuclei at MAMI-C: KAOS
- ▶ The GSI project: PANDA
- ▶ Search for the $\Xi(1860)$ in WA89

Birth, life and death of a hypernucleus



Reminder: momentum distributions



- ▶ semi inclusive measurements of one-nucleon knockout reactions provide information on the momentum distribution of the nucleon and hence on the nucleon wave function



$$E_x = W - T_p - T_{A-1} - E_s$$

$$p_m = q - p'$$

- ▶ Example: $^{12}\text{C}(e, e' p)^{11}\text{B}$
- ▶ Steenhoven et al., (1988)



Double Hypernuclei

- Multi-Hypernuclei are terra incognita, but they exist !

1963: Danysz et al. $\Lambda\Lambda^{10}\text{Be}$

1966: Prowse $\Lambda\Lambda^6\text{He}$

1991: KEK-E176 $\Lambda\Lambda^{13}\text{B}$ (or $\Lambda\Lambda^{10}\text{Be}$)

2001: AGS-E906 $\Lambda\Lambda^4\text{H}$ (~ 15);
no binding energies


2001: KEK-E373 $\Lambda\Lambda^6\text{He}$

"

(Nagara)

$\Lambda\Lambda^{10}\text{Be}$


(Demachi-Yanagi)



H. Takahashi et al., PRL 87, 212502-1
(2001)

2. ($e, e' K$) at MAMI-C


MAMI-C



- ▶ Electron energy: 0.855 GeV → 1.5 GeV
- ▶ first beam end of 2005

Existing data and experiments


- ▶ photoproduction
 - ▶ 1970: Cornell, CalTech, Bonn, DESY, Orsay
 - ▶ 2000: SAPHIR (Bonn), GRAAL (Grenoble), CLAS (Jlab, Hall B)
- ▶ electroproduction
 - ▶ 1970: Cornell, DESY, CEA
 - ▶ 2000: E93-018 (Jlab, Hall C), E98-108 (Jlab, Hall A)



- ▶ for $Q^2 < 0.5 \text{ (GeV/c)}^2$ only few precise data

New Experimental Opportunities

- ▶ Crystal ball + TAPS (A2)
 - ▶ real photons
 - ▶ calorimetry of neutral and charged particles



- ▶ KAOS (A1)
 - ▶ virtual photons
 - ▶ high resolution spectroscopy of charged particles




(→ Volker Kochs talk)

Hypernucleus production

- ▶ cross section for $A(e, e' K^+) \Lambda B$ in e.g. relativistic impulse approximation

$$ds \propto (2p)^4 d^4(p_e + p_A - p_{e'} - p_B - p_K) \\ \times \frac{d^3 p_e m_e}{(2p)^3 E_{e'}} \cdot \frac{d^3 p_K}{(2p)^3 E_K} \cdot \frac{d^3 p_B (2m_p)(2m_e)}{(2p)^3 \sqrt{(p_e \cdot p_A)^2 - p_e^2 p_A^2}} \\ \times \sum_{M_f M_i} \left| \sum_{a a'} \langle J_f M_f T_f N_f | C_a^\dagger C_a | J_i M_i T_i N_i \rangle \cdot \langle a' | t | a \rangle \right|^2$$


matrix elements \times elementary operator for $p(g, K^+) \Lambda$





- ▶ key ingredient to the elementary operator $p(\gamma, K^+) \Lambda$
 - ▶ nucleon wave function and spectroscopic factor (e.g. from $(e, e' p)$)
 - ▶ K^+ optical potential (e.g. from K^+ elastic scattering)
 - ▶ Λ wave function
- ▶ we can map out the Λ wave function by varying the kaon angle

Kinematics

- ▶ qualitative example
 - ▶ $\langle p_{K_+} \rangle = 0.8 \text{ GeV}/c$
 - ▶ $\langle p_e \rangle = 0.4 \text{ GeV}/c$
 - ▶ $\langle \theta_{K_+} \rangle = 6.8^\circ$
 - ▶ averaged over all kaon momenta




Kinematics $e + {}^A Z \rightarrow e' + K^+ + {}^A_{\Lambda} (Z - 1)$



- ▶ strong correlation between kaon and scattered electron momentum
 - ▶ large momentum acceptance for both particles
- ▶ both particles are produced at small angle
 - ▶ forward double spectrometer


KAOS at 0°

- ▶ large count rate
- ▶ chicane for primary beam



KaoS@MAMI

- ▶ arrival of KaoS at Mainz on 11 June 2003



Fiberdetectors for KAOS




► requirements

- ▶ high count rate capability
- ▶ good position resolution (150 mm)
- ▶ timing information (needed for trigger)

► realization


- ▶ 0.83 mm fiber diameter
- ▶ 4 fibers=1 PMT channel
- ▶ 4000 channels
- ▶ multi-anode phototube (32 ch/tube)
- ▶ 32-ch discriminator based on double threshold discriminator chip GSI-3

► funded by HBFG



KAOS @ A1


- ▶ planing for setting up KaoS is ongoing
 - ▶ electrical power, water cooling...
 - ▶ parking position, vertical and horizontal moving
 - ▶ at 0°: chicane for beam dump (**background!**)
 - ▶ trigger
 - ▶ ...



3. The GSI Future Project


The GSI Future Project

- ▶ origin of confinement ?
- ▶ hierarchy of quark masses ?
- ▶ atomic nucleus and nuclear matter as quark-gluon systems ?




Antiproton Storage Ring

- ▶ Antiproto momentum 1.5 – 15 GeV/c
- ▶ Luminosity $2 \cdot 10^{32} \text{ cm}^{-2}\text{s}^{-1}$
- ▶ beam diameter 10-100 μm
- ▶ $\Delta p/p = 10^{-4} \dots 3 \cdot 10^{-5}$



The PANDA Project

- ▶ Anti-proton storage ring HESR
 - ▶ internal target
- ▶ Physics program
 - ▶ quark-quark interaction
⇒ charmonium spectroscopy
 - ▶ gluonische degress of freedom
⇒ glueballs, hybrids
 - ▶ hadrons in nuclear matter
⇒ mesons in nuclei
⇒ Hyperkerne
- ▶ Far future
 - ▶ CP violation?
 - ▶ inverse DVCS ?



Double hypernuclei: what is known?

$$B_{\Lambda\Lambda}({}_{\Lambda\Lambda}^A Z) = B_\Lambda({}_{\Lambda\Lambda}^A Z) + B_\Lambda({}_{\Lambda\Lambda}^{A-1} Z)$$

$$\Delta B_{\Lambda\Lambda}({}_{\Lambda\Lambda}^A Z) = B_\Lambda({}_{\Lambda\Lambda}^A Z) - B_\Lambda({}_{\Lambda\Lambda}^{A-1} Z)$$

| Hyperkern | $B_{\Lambda\Lambda}$ (MeV) | $\Delta B_{\Lambda\Lambda}$ (MeV) | |
|-------------------------------|------------------------------------|------------------------------------|------------------------------|
| ${}_{\Lambda\Lambda}^6 He$ | 10.9 ± 0.5 | 4.7 ± 0.6 | Prowse (1966) |
| ${}_{\Lambda\Lambda}^6 He$ | $7.25 \pm 0.19 {}^{+0.18}_{-0.11}$ | $1.01 \pm 0.20 {}^{+0.18}_{-0.11}$ | KEK-E373 (2001) |
| ${}_{\Lambda\Lambda}^{10} Be$ | 17.7 ± 0.4 | 4.3 ± 0.4 | Danysz (1963) |
| ${}_{\Lambda\Lambda}^{10} Be$ | 8.5 ± 0.7 | -4.9 ± 0.7 | KEK-E176 (1991) |
| ${}_{\Lambda\Lambda}^{13} B$ | 27.6 ± 0.7 | 4.8 ± 0.7 | KEK-E176 (1991) |
| ${}_{\Lambda\Lambda}^{10} Be$ | $12.33 {}^{+0.35}_{-0.21}$ | | KEK-E373 (2001, unpublished) |

same event



- ▶ Interpreting $\Delta B_{\Lambda\Lambda}$ as $\Lambda\Lambda$ bond energy one has to consider e.g.
 - ▶ dynamical change of the core nucleus
 - ▶ ΛN spin-spin interaction for non-zero spin of core
 - ▶ excited states possible ...
- ▶ if $\Lambda\Lambda$ - or intermediate Λ -nuclei are produced in excited states
 - ▶ Q-value difficult to determine (particularly for heavy nuclei)
 - ▶ nuclear fragments difficult to identify with usual emulsion technique
- ▶ new concept required $\Rightarrow \gamma$ -spectroscopy!

Production of $\Lambda\Lambda$ -Hypernuclei


- Ξ^- conversion in 2 Λ




- Ξ^- production
 - $p(K^-, K^+) \Xi^-$
 - needs K^- beam ($c \cdot \tau = 3.7 \text{ cm}$)
 - recoil momentum $> 460 \text{ MeV}/c$
 - KEK-E176: 10^2 stopped Ξ
 - E373: 10^3 stopped Ξ
 - AGS-E885: 10^4 stopped Ξ
- at HESR



- few times 10^5 stopped Ξ per day
- γ -spectroscopy possible




Production of Double Hypernuclei





General Idea

- ▶ Use $\bar{p}p$ Interaction to produce a hyperon "beam" ($t \sim 10^{-10}$ s) which is tagged by the antihyperon or its decay products



Ξ^- properties



- Ξ^- mean lifetime 0.164 ns



- minimal distance production \Rightarrow capture
- initial momentum $100\text{-}500 \text{ MeV/c} \Rightarrow$ range $\sim \text{few g/cm}^2$

Ξ^- capture

- ▶ Ξ^- -atoms: x-rays
- ▶ conversion
 - ▶ $\Xi^-(dss) \rightarrow \Lambda(uud) \Lambda(uds)$
 - ▶ $\Delta Q = 28 \text{ MeV}$
- ▶ Conversion probability approximately 5-10%



Expected Count Rate

► Ingredients (golden events: Ξ^+ trigger)

- luminosity $2 \cdot 10^{32} \text{ cm}^{-2}\text{s}^{-1}$
- $\Xi^+\Xi^-$ cross section 2mb for pp
- $p(100\text{-}500 \text{ MeV/c})$
- Ξ^+ reconstruction probability
- stopping and capture probability
- total captured Ξ^-

P 700 Hz
 $p_{500} \approx 0.0005$
0.5
 $p_{\text{CAP}} \approx 0.20$
 P 3000 / day

- Ξ^- to $\Lambda\Lambda$ -nucleus conversion probability
- total $\Lambda\Lambda$ hyper nucleus production

$p_{\Lambda\Lambda} \approx 0.05$
 P 4500 / month

- gamma emission/event,
- γ -ray peak efficiency

$p_\gamma \approx 0.5$
 $p_{\text{GE}} \approx 0.1$


- ~7/day „golden“ γ -ray events
- ~ several 100/day with KK trigger

high resolution γ -spectroscopy of double hypernuclei will be feasible

Competition

| experiment | reaction | device | beam/ target | status |
|-------------------|--|--|---|---|
| BNL-AGS E885 | $(\Xi^-, {}^{12}\text{C}) \rightarrow {}^{12}\text{B} + n$ LL | neutron detector arrays | K- beam, diamond target | 20000 stopped Ξ^- |
| BNL-AGS E906 | 2π decays | Cylindrical Detector System | K- beam line | few tens 2π decays of ${}^4_{\Lambda\Lambda}\text{H}$ |
| KEK-PS E373 | $(K^-, K^+) \Xi$ | emulsion | (K^-, K^+) | several hundreds stopped Ξ^- |
| facility | reaction | device | beam/ target | Captured X- per day |
| JHF | $(K^-, K^+) \Xi$ | spectrometer, $\Delta\Omega = 30$ msr | $8 \cdot 10^6/\text{sec}$ $5 \text{ cm } {}^{12}\text{C}$ | <7000 |
| cold anti-protons | $p \bar{p} \rightarrow K^* \bar{K}^*$ $\bar{K}^* N \rightarrow X K$ | vertex detector | 10^6 stopped \bar{p} per sec | 2000 |
| GSI-HESR | $p p \rightarrow \Xi \Xi$ | vertex detector + γ -spectrometer | $L = 2 \cdot 10^{32}$, thin target, production vertex \neq decay vertex | 3000 „golden events“ ~ 100000 KK trigger (numbers incl. trigger) |

Production of Ω -Atoms



Fundamental Properties of Baryons


- ▶ Contributions to intrinsic quadrupole moment of baryons
 - ▶ One-gluon exchange
 - ▶ Meson exchange

$$Q_i = \int d^3r r(r)(3z^2 - r^2)$$


- ▶ $J=1/2$ baryons have no spectroscopic quadrupole moment

$$Q_s \propto (3J_z^2 - J(J+1)) \xrightarrow[J_z=1/2]{J=1/2} 0$$

- ▶ Ω^- Baryon:
 - ▶ $J=3/2$
 - ▶ long mean lifetime $0.82 \cdot 10^{-10}$ s
 - ▶ only one-gluon contributions to quadrupole moment
- (A.J. Buchmann Z. Naturforsch. 52 (1997) 877-940)



A very strange Atom



- ▶ Ω atoms by $\Omega\bar{\Omega}$ production
 - ▶ hyperfine splitting in Ω -atom
⇒ electric quadrupole moment of Ω

| | |
|------------|--|
| spin-orbit | $\Delta E_{ls} \sim (aZ)^4 l \cdot m_\Omega$ |
| quadrupole | $\Delta E_\Theta \sim (aZ)^4 Q m^3 \Omega$ |

- ▶ prediction $Q_\Omega = (0 - 3.1) \text{ } 10^{-2} \text{ fm}^2$
 - ▶ $E(n=11, l=10 \rightarrow n=10, l=9) \sim 520 \text{ keV}$
 - ▷ calibration with 511keV line!
 - ▶ $\Delta E_\Theta \sim \text{few tenth of keV for Pb}$

close to
511keV



R.M. Sternheimer, M. Goldhaber
PRA 8, 2207 (1973)

M.M. Giannini, M.I. Krivoruchenko
 Phys. Lett. B 291, 329 (1992)

with high resolution γ -spectroscopy difficult
but feasible if statistics sufficient

The PANDA Detector

- ▶ hermetic (4π)
- ▶ high rate
- ▶ PID ($\gamma, e, \mu, \pi, K, p$)
- ▶ trigger (e, μ, K, D, Λ)
- ▶ compact (ϵ)
- ▶ modular



- ▶ Solid state-micro-tracker
 - ▶ thickness ~ 3 cm
- ▶ High rate germanium detector

4. Ξ(1860) search in WA89

NA49: Ξ^{--} Pentaquark

Observation of an Exotic $S = -2$, $Q = -2$ Baryon Resonance in Proton-Proton Collisions at the CERN SPS

C. Alt,⁹ T. Anticic,²⁰ B. Baatar,⁸ D. Barna,⁴ J. Bartke,⁶ M. Behler,¹³ L. Betev,^{10,9} H. Biakowska,¹⁸ A. Billmeier,⁹ C. Blume,^{7,9} B. Boimska,¹⁸ M. Botje,¹ J. Bracinik,³ R. Bramm,⁹ R. Brun,¹⁰ P. Bunčić,^{9,10} V. Černý,³ P. Christakoglou,² O. Chvala,¹⁵ J.G. Cramer,¹⁶ P. Csató,⁴ N. Darmenov,¹⁷ A. Dimitrov,¹⁷ P. Dinkelaker,⁹ V. Eckardt,¹⁴ G. Farantatos,² P. Filip,¹⁴ D. Flierl,⁹ Z. Fodor,⁴ P. Foka,⁷ P. Freund,¹⁴ V. Friese,^{7,13} J. Gál,⁴ M. Gaździcki,⁹ G. Georgopoulos,² E. Gladysz,⁶ S. Hegyi,⁴ C. Höhne,¹³ K. Kadija,²⁰ A. Karev,¹⁴ S. Kniege,⁹ V.I. Kolesnikov,⁸ T. Kollegger,⁹ R. Korus,¹² M. Kowalski,⁶ I. Kraus,⁷ M. Kreps,³ M. van Leeuwen,¹ P. Lévai,⁴ L. Litov,¹⁷ M. Makariev,¹⁷ A.I. Malakhov,⁸ C. Markert,⁷ M. Mateev,¹⁷ B.W. Mayes,¹¹ G.L. Melkumov,⁸ C. Meurer,⁹ A. Mischke,⁷ M. Mitrofski,⁹ J. Molnár,⁴ St. Mrówczyński,¹² G. Pálka,⁴ A.D. Panagiotou,² D. Panayotov,¹⁷ K. Perl,¹⁹ A. Petridis,² M. Pikna,³ L. Pinsky,¹¹ F. Pühlhofer,¹³ J.G. Reid,¹⁶ R. Renfordt,⁹ W. Retyk,¹⁹ C. Roland,⁵ G. Roland,⁵ M. Rybczyński,¹² A. Rybicki,^{6,10} A. Sandoval,⁷ H. Sann,^{7,*} N. Schmitz,¹⁴ P. Seyboth,¹⁴ F. Siklér,⁴ B. Sitar,³ E. Skrzypeczak,¹⁹ G. Stefanek,¹² R. Stock,⁹ H. Ströbele,⁹ T. Susa,²⁰ I. Szentpétery,⁴ J. Sziklai,⁴ T.A. Trainor,¹⁶ D. Varga,⁴ M. Vassiliou,² G.I. Veres,^{4,5} G. Vesztergombi,⁴ D. Vranić,⁷ A. Wetzler,⁹ Z. Włodarczyk,¹² I.K. Yoo,⁷ J. Zaranek,⁹ and J. Zimányi⁴

(NA49 Collaboration)

¹NIKHEF, Amsterdam, Netherlands.

²Department of Physics, University of Athens, Athens, Greece.

³Comenius University, Bratislava, Slovakia.

⁴KFKI Research Institute for Particle and Nuclear Physics, Budapest, Hungary.

⁵MIT, Cambridge, MA, USA.

⁶Institute of Nuclear Physics, Cracow, Poland.

⁷Gesellschaft für Schwerionenforschung (GSI), Darmstadt, Germany.

⁸Joint Institute for Nuclear Research, Dubna, Russia.

⁹Fachbereich Physik der Universität, Frankfurt, Germany.

¹⁰CERN, Geneva, Switzerland.

¹¹University of Houston, Houston, TX, USA.

¹²Świetokrzyska Academy, Kielce, Poland.


¹³Fachbereich Physik der Universität, Marburg, Germany.

¹⁴Max-Planck-Institut für Physik, Munich, Germany.


¹⁵Institute of Particle and Nuclear Physics, Charles University, Prague, Czech Republic.

Ξ^- in NA49

- ▶ distance between primary and secondary vertex 12cm
- ▶ additional cuts on track impact position and angle
- ▶ 1640 Ξ^- events
- ▶ 551 anti- Ξ^+ events



- ▶ Ξ^- combined with primary π^-
- ▶ $\theta(\pi^-, \Xi^-) > 4.4^\circ$
- ▶ $p(\pi^+) > 3\text{GeV}/c$
- ▶ ...several additional cuts



The WA89 Experiment

- ▶ Σ^- and π^- beam of 340 GeV/c, n-beam of 260 GeV/c
- ▶ 1993, 1994 data taking
- ▶ $4 \cdot 10^8$ interactions

WA89
Hyperon beam
1993 layout



TRD: beam identification

Si- μ -strip vertex near target

MWPC: tracking


RICH: π/K separation

Calorimeter: e, γ , n




Cross sections

- ▶ more than 20 different strange hadrons are analyzed under identical conditions
- ▶ typical statistical distribution with slope ~ 150 MeV



Ξ^- Mass Spectrum

► $\Xi^- \rightarrow \Lambda^0 \pi^-$
 $\Xi^- \rightarrow p \pi^-$



$X^{*0}(1690) \rightarrow X^- p^+$

► $\Xi^{*0}(1690) \rightarrow \Xi^- \pi^+$

$$\begin{array}{c} \Xi^- \Lambda^0 \pi^+ \\ \Xi^- p \pi^- \end{array}$$

► $M(\Xi^{*0}) = 1685 \pm 4 \text{ MeV}/c^2$

► $\Gamma = 10 \pm 6 \text{ MeV}/c^2$

► $\sigma \cdot BR = 6.8 \pm 0.2 \mu\text{b}$



► $M(\Xi^*) - M(\Xi^{*0}) =$

► $\Delta(\Xi_{1690}) = 6 \pm 5 \text{ MeV}/c^2$

► $\Delta(\Xi_{1530}) = 6.4 \pm 0.6 \text{ MeV}/c^2$

► $\Delta(\Xi_{1320}) = 3.2 \pm 0.6 \text{ MeV}/c^2$

Euro. Phys. J. C 5, 621 (1998)





$\Xi^*(1820)$ and $\Xi^*(1960)$ Production

- $\Xi^* \rightarrow \Xi^{*0}(1530) \pi^-$
- $\hat{\Xi} \Xi^- \pi^+$
- $\hat{\Xi} \Lambda^0 \pi^-$
- $\hat{\Xi} p \pi$


Eur. Phys. J. C11, 271 (1999)

$\Xi^*(1820)$

$\Xi^*(1960?)$



$M(\Xi^{*0}(1.53)+\pi^-), \text{GeV}/c^2$



- ▶ further cuts on p_t , x_F , θ_{cm}
- ▶ additional (justified) cuts?
- ▶ ...more to come

Summary

- ▶ Hypernuclear Physics is getting mature after 50 years
- ▶ MAMI-C is on track
- ▶ PANDA and HESR will enable high statistics double hypernuclear physics
 - ▶ many questions to answer, many problems to solve
 - ▶ comments, suggestions, help... are very welcome



See you all
in Mainz
2006



JOHANNES
GUTENBERG
UNIVERSITÄT
MAINZ

