


## Part IV: General Formalism of PQCD

- Factorization Theorem to all orders of the perturbative expansion (with non-zero quark masses);
- The importance of Scales, Factorization and Renormalization;
- General definition of Parton Distribution Functions (PDF);
- The Three Faces of the Magical Factorization Master Formula;
- Scale dependence of PDFs and QCD Evolution;
- Scale dependence (and independence) of Physical Predictions.

What happens when we consider the other NLO subprocess  $\gamma q \rightarrow gq$ ?

Square:



Compared to the  $\gamma^*g$  case:

- Order  $\alpha_s$  terms are divergent: needs UV renormalization;
- Combine real and virtual diagrams to ensure cancellation of infra-red divergences (cf.  $e^+e^-$  case);
- Remaining infra-red unsafe terms are associated with long-distance physics: they are universal, and factorizable into a parton distribution function (—in this case, quark/quark).

The IRS hard scattering cross section is the remainder, after the long-distance piece is subtracted (at the factorization scale  $\mu_f$ ).

## What Happens at Higher Orders in $\alpha_s$ and for Other Hard Processes?

The same pattern of behavior is repeated order-by-order for most hard processes which has been calculated so far, to 1- or 2-loop level.

To prove the validity of the factorization formula *to all orders of PQCD* is a highly theoretical and technical matter (very sophisticated mathematical induction methods).


Serious proofs exist only for a limited number of processes such as  $e^+e^-$ , DIS and DY. Faith in general factorization rests on existing concrete 1- or 2-loop calculations, and on the suggestive space-time picture.

Traditional proofs of factorization assume  $m=0$  as the starting point. J. Collins showed that factorization in DIS (hence probably other processes as well) is independent of the value of quark masses. (hep-ph/9806259) Our sample calculation with finite quark mass illustrates why this is the case. (The hard cross sections contain IR mass effects.)

## What does a Factorization Thm Proof Consist of?

- It involves the demonstration that, to all orders of  $\alpha_s$  in PQCD, a set of *universal parton distribution functions* exists, which will absorb all the collinear and soft singularities encountered in the calculation of all relevant partonic scattering processes.
- Two points should immediately come to mind:
  - This situation is very similar to renormalization, where all (UV) singularities are shown to be absorbed into a set of finite number of renormalization constants; (next two slides)
  - If such universal functions (PDF) exist, there must be a precise definition of them - in a *process-independent* way. (topic after that)

# The importance of **Scales** -- Renormalization and Factorization



Renormalization Group Equations (RGE) relates physics at different scales

Ultra-violet  
Renormalization  
hides / summarizes  
our ignorance of  
physics at huge scale in  
 $\alpha_s(\mu_R), m_i(\mu_R), \dots$

# “Renormalization” and “Factorization”

UV renormalization		Collinear/soft factorization	
A: Bare Green Func.	$G_0(\alpha_0, m_0, ..)$	Partonic X-sect	$F_a$
B: Ren. constants	$Z_i(\mu)$	Pert. parton dist.	$f_a^b(\mu)$
C: Ren. Green Fun.	$G_R = G_0/Z$	Hard X-sect	$\hat{F} = F / f$
D: Anomalous dim.	$\gamma = \frac{\mu}{Z} \frac{d}{d\mu} Z$	Splitting fun.	$P = \frac{\mu}{f} \frac{d}{d\mu} f$
E: Phys. para. $\alpha, m$	$\alpha_0 Z_i \dots$	Had. parton dist. $f_A$	resummed
F: Phys sc. amp.	$\alpha(\mu) G_R(m, \mu)$	Hadronic S.F.'s $F_A$	$f_A(\mu) \times \hat{F}(\mu)$

Some common features:

A : divergent; but, independent of “scheme” and scale  $\mu$ ;

B : divergent; scale and scheme dependent;  
universal; absorbs all ultra-violet/soft/collinear divergences;

C & D : finite; scheme-dependent;  
D controls the  $\mu$  dependence of E & F;

E : physical parameters to be obtained from experiment;

F : Theoretical “prediction”;  $\mu$ -indep. to all orders,  
but  $\mu$ -dep. at finite order  $n$ ;  $\mu \frac{d}{d\mu} \sim \mathcal{O}(\alpha^{n+1})$

Note: “Renormalization” is factorization (of UV divergences);  
“factorization” is renormalization (of soft/collinear div.)

$$F = f \otimes \hat{F}$$

## $\overline{\text{MS}}$ definition of parton distribution functions

Factorization Theorem requires a precise definition of the parton distribution functions (pdf's).

Note:  $F_A = f_A^a \otimes \hat{F}_a \implies \hat{F}_a = \frac{F_A}{f_A^a}$

$\implies$  the definition of hard scattering cross-sections is determined by that of pdf's.

Historical definition of pdf's: use "operator product expansion" – towers of local operators.

Equivalent modern definition:  
use bi-local operators on the light cone:

## Definition of PDF's

The distribution of quark “ $a$ ” in a parent “ $A$ ” (either hadron or another parton) is:

$$f_A^a(\xi) \equiv \frac{1}{2} \int \frac{dy^-}{2\pi} e^{-i\xi p^+ y^-} \langle A | \bar{\psi}_a(0, y^-, \mathbf{0}) \gamma^+ F \psi_a(0) | A \rangle.$$

$|A\rangle$  represents the parent state  $A$ , with momentum  $p^\mu$  aligned so that  $p_T = 0$  and  $p_a^+ / p_A^+ = \xi$ . This is the matrix element of the appropriate number operator for finding “ $a$ ” in “ $A$ ”.

The operator

$$F = \mathcal{P} \exp \left( -ig \int_0^{y^-} dz^- A_a^+(0, z^-, \mathbf{0}) t_a \right)$$

ensures the definition gauge invariant.

There is a similar definition of the gluon distribution inside the hadron  $A$ , using the gluon field operator.


## Definition of PDF's

- The  $\overline{\text{MS}}$  definition in terms of operators is process independent, re-affirming that PDF's are *universal*.
- Sum rules are automatic. Eg.

$$\sum_a \int_0^1 d\xi \xi f_A^a(\xi, \mu) = 1.$$

Space-time picture of the definition of PDF's is very similar to that of DIS structure functions:

$\langle p | \psi(x) \bar{\psi}(0) | p \rangle$  vs.  $\langle p | J(x) J^\dagger(0) | p \rangle$ :



# What does factorization of partonic cross sections have to do with hadronic cross sections.?

From the fact that

- the factorization theorem is proven for partonic cross-sections to all orders in PQCD;
- the parton distributions are universal; and are defined for hadron as well as parton parents.
- the hard cross-section is infra-red safe;

it is natural to assume,

Hadronic cross-sections, such as  $F_A^\lambda(x, Q)$ , satisfy the same factorization formula as the partonic structure functions:


$$F_A^\lambda(x, Q) = \sum_a f_A^a(x, \mu) \otimes \hat{F}_a^\lambda(x, \frac{Q}{\mu}) \quad (a = q, g)$$

and that *the hadronic parton distributions  $f_A^a(x, \mu)$  are finite.*

# The three faces of the Magical Factorization Formula

## #1 (seen in illustrative sample calculation)

Choose  $A$  to be a parton (say,  $A = b$ ):



1. Calculate the partonic cross section  $F_b$  perturbatively (contains collinear and soft singularities).

2. Calculate the perturbative PDFs  $f_b^a$  (also contains collinear and soft singularities)



Subtract (2) from (1), and derive the (IRS) hard cross sections  $\hat{F}_a$  (Wilson coefficients).

# The three faces of the Magical Factorization Formula

#2

Choose  $F_A$  to be a set of established experimental cross sections

1. Input: the well-measured data for all  $\{F_A\}$ .




2. Input: known perturbatively calculated hard cross sections  $\hat{F}_a$

Perform a Global QCD Analysis to determine the universal (but perturbatively non-calculable) hadronic parton distributions  $f_A^a(x, \mu)$ .

# The three faces of the Magical Factorization Formula

#3

Choose  $F_A$  to be some physical cross section of interest:



1. Input **hadronic PDFs** determined by global QCD analysis .

2. Input: known perturbatively calculated hard cross sections  $\hat{F}_a$

Predict cross sections for SM or New Physics Processes of interest!

# The QCD Evolution Equation & Scale dependence of PDFs

Parton distributions represent long-distance physics,

- for parton targets,  $f_c^a$  are calculable, but contain collinear/soft “divergences”;
- for hadron targets,  $f_A^a$  are finite, but not calculable in perturbation theory.

Since,  $f_c^a$  are calculable, one can examine their  $\mu$ -dependence:

$$\frac{d}{d \ln \mu} f_c^a(x, \mu) = \sum_b \int_x^1 \frac{d\xi}{\xi} P_{ab}\left(\frac{x}{\xi}, \alpha_s(\mu)\right) f_c^b(\xi, \mu).$$

where  $P_{ab}$  on the RHS (the splitting functions) has the perturbative form

$$P_{ab}(z, \alpha_s(\mu)) = P_{ab}^{(1)}(z) \frac{\alpha_s(\mu)}{\pi} + P_{ab}^{(2)}(z) \left(\frac{\alpha_s(\mu)}{\pi}\right)^2 + \dots$$

Note, the splitting functions  $P_{ab}(x)$  are independent of the parent  $c$ , *because of factorization*.

Hence, the hadronic (i.e. physical) structure functions satisfy the *same QCD evolution equation*, with the *same evolution kernels* (splitting function) that are calculated using partonic amplitudes!

$$\frac{d}{d \ln \mu} f_{\mathbf{A}}^a(x, \mu) = \sum_b \int_x^1 \frac{d\xi}{\xi} P_{ab}\left(\frac{x}{\xi}, \alpha_s(\mu)\right) f_{\mathbf{A}}^b(\xi, \mu)$$

Compare with the more familiar evolution equation for scale dependence of renormalization constants (e.g.  $\alpha_s(\mu)$ ),

$$\mu \frac{d}{d\mu} Z_i(\mu) = \gamma_i Z_i(\mu)$$

$P_{ab}(x)$  (calculable order by order in  $\alpha_s$ ) is the analogue of the *anomalous dimension* parameters  $\gamma_i$  that are characteristic of the underlying theory (QCD).

# The Scale Dependence of Physical Predictions


$$F_A^\lambda(x, Q, m) = \sum_a f_A^a(x, \mu_F, m) \otimes \hat{F}_a^\lambda(x, \frac{Q}{\mu_F}, \frac{m}{\mu_F}, \alpha_x(\mu_R))$$

In principle, the choice of factorization scale  $\mu_f$  should not affect physical predictions:

graphically,

analytically, in the perturbative approach:

$$\begin{aligned} \mu \frac{d}{d\mu} F &= \mu \frac{d}{d\mu} f \otimes \hat{F} + f \otimes \mu \frac{d}{d\mu} \hat{F} \\ &= O(\alpha_s^{N+1}) \end{aligned}$$




i.e. the physical  $F$ 's are *scale independent*, to the order of the pert. calculation.

# How to Choose the Renormalization/Factorization Scale?

- Similar considerations apply to  $\mu_R$  and  $\mu_F$ ; to simplify the discussion, let  $\mu_R = \mu_F = \mu$ .
- In order to apply perturbative expansion, we need  $\alpha_s(\mu)/\pi \ll 1 \Rightarrow \mu \gg \Lambda_{QCD}$ .
- Since  $\hat{F}_A$  contains terms of the form  $\alpha_s^n(\mu) \ln^n(Q/\mu)$  (or even  $\ln^{2n}(Q/\mu)$  in some cases), we cannot have  $\mu \gg Q$  without spoiling the perturbative approach.
- Therefore,  $\mu$  must be chosen to be of the order order as the hard scale  $Q$ , i.e.  $\mu = cQ$  with  $c \sim 1$ .
- To estimate the uncertainty of a  $N$ th order calculation, we often take  $\Delta F_A$  to be the range of variation of  $F_A$  calculated with  $\mu = cQ$  and  $\frac{1}{2} < c < 1$ —keeping in mind that  $\mu \frac{d}{d\mu} F_A = O(\alpha_s^{N+1})$ .

# Homework assignment

An exotic boson  $Z'$  with mass  $Q = 2$  TeV is produced similarly to SM  $Z$  bosons, but only via the  $s\bar{s} \rightarrow Z'$  vertex ( $Z'$  does not interact with non-strange (anti-)quarks).



$Z'$  couples only to  $s, \bar{s}$

You need to compute  $\sigma(pp \rightarrow Z'X)$  at the LHC  $\sqrt{s} = 13000$  GeV, but for that you need to precisely know the strange (anti-)quark PDFs,  $s(x, Q)$  and  $\bar{s}(x, Q)$ . Propose one or two scattering processes to constrain  $s(x, Q)$  and  $\bar{s}(x, Q)$  at the relevant  $\{x, Q\}$ . Specify  $\sqrt{s}$  and other kinematic parameters of these processes. Can you use non-LHC measurements to constrain  $s(x, Q)$  at the LHC? Why or why not?