

# Sigma terms and currents in $SU(3)$ BChPT $\times 1/N_c$

Physics Letters B 781 (2018) 719–722

Phys. Rev. D 97, 054010 (2018)

arXiv:1911.00987 (recently submitted to PRD)

## Ishara Fernando

*Collaborators*

**José Goity, José Manuel Alarcon**

**JLab Theory Seminar**

**January 15, 2020**



This work was supported by  
National Science Foundation PHY-1307413 and No. PHY-1613951  
&  
Department of Energy Contract No. DE-AC0506OR23177,



## Outline

- 1) Motivation for the BChPT  $\times 1/N_c$  expansion
- 2) Introduction to the BChPT  $\times 1/N_c$  expansion (combined) approach
- 3) Baryon masses
- 4) Baryon sigma terms
- 5) Vector currents
- 6) Axial-vector currents
- 7) Summary



Non relativistic version of the BChPT or HBChPT is based on the expansion in terms of the “baryon mass”

Derivative expansion for both mesons and baryons becomes an expansion in powers of  $(k/\Lambda_\chi)$

The issue of experiencing a slower rate of convergence compared to the Goldstone Boson Sector

$$p_\mu = m_B v_\mu + k_\mu$$

$$\frac{1}{p^2 - m_B^2} \rightarrow \frac{1}{2m_B} \frac{1}{(v.k)} + \mathcal{O}(1/m_B^2)$$

$$v^\mu v_\mu = 1$$

$$v.k \ll m_B$$


Solution : Inclusion of the decuplet baryons in one-loop corrections to physical observables, has been showing a great improvement!

These improvements are due to cancellations between octet and decuplet contributions in loops

On the other hand, studying the baryons in the large Nc limit of QCD emerges a dynamical symmetry called “spin-flavor symmetry” which requires the possibility of having degenerate baryon multiplets of higher spin in the intermediate state/s.

# 1.1 Spin-flavor symmetry of Baryons in large $N_c$

Gervais & Sakita; Dashen & Manohar



Since,  $\pi N$  amplitude is  $\mathcal{O}(N_c^0)$

$$A = -i \frac{k^i k^j}{k_0} \frac{N_c^2 g^2}{f_\pi^2} [X^{ia}, X^{ib}] \Rightarrow [X^{ia}, X^{ib}] \leq \mathcal{O}(1/N_c)$$

Large  $N_c$  consistency condition  
 $[X_0^{ia}, X_0^{ib}] = 0$


At large  $N_c$ , QCD has contracted spin-flavor symmetry  $SU_c(2N_f)$  in baryon sector

$$X_0^{ia} = \lim_{N_c \rightarrow \infty} \frac{G^{ia}}{N_c}$$

This symmetry is broken at sub-leading orders in  $1/N_c$

This spin-flavor symmetry requires the existence of degenerate baryon multiplets with different spins (a dynamical symmetry) : leads to the consideration of both octet and decuplet contributions in the intermediate state

$$L(\text{Lagrangian}) = x^0 L_{LO} + x^1 L_{NLO} + x^2 L_{NNLO} + x^3 L_{NNNLO} + \dots$$



Spin-flavor Symmetry

+

Chiral Symmetry

imposes constraints in  
the Chiral Lagrangian


Combined approach

Combining the HBChPT with 1/N<sub>c</sub> provides a  
well behaved expansion  
in the low energy phenomenology  
(because one cannot expand them  
independently in low energy)

Link between the Chiral and 1/N<sub>c</sub> expansion

$$\xi - \text{expansion} : \mathcal{O}(1/N_c) = \mathcal{O}(p) = \mathcal{O}(\xi)$$

Chiral Symmetry + Spin-flavor Symmetry



$$= \int \frac{d^d k}{(2\pi)^d} \frac{i}{k^2 - M_\pi^2} \frac{i}{p^0 + k^0 - \underbrace{(m_{B'} - m_B)}_{Q \sim \mathcal{O}(1/N_c)}} \times \text{vertex factors}$$

Intermediate Octet and Decuplet baryon contributions are included

$\xi$  - expansion :  $\mathcal{O}(1/N_c) = \mathcal{O}(p) = \mathcal{O}(\xi)$

$$\begin{aligned} I_{1-loop}(Q, M_\pi) &= \int \frac{d^d k}{(2\pi)^d} \frac{\vec{k}^2}{k^2 - M_\pi^2 + i\epsilon} \frac{1}{k^0 - Q + i\epsilon} \\ &= \frac{i}{16\pi^2} \left\{ Q \left( (3M_\pi^2 - 2Q^2)(\lambda_\epsilon - \log \frac{M_\pi^2}{\mu^2}) + (5M_\pi^2 - 4Q^2) \right) \right. \\ &\quad \left. + 2\pi(M_\pi^2 - Q^2)^{3/2} + 4(Q^2 - M_\pi^2)^{3/2} \tanh^{-1} \frac{Q}{\sqrt{Q^2 - M_\pi^2}} \right\}, \end{aligned}$$

Contains both scales: therefore cannot be expanded independently

$$Q = \delta m_n - p^0, \quad \lambda_\epsilon = \frac{1}{\epsilon} - \gamma + \log 4\pi$$

## Building blocks :

➔ Goldstone bosons: pions, kaons, eta

$$u = \exp\left(\frac{i\Pi}{2F_\pi}\right) \quad \text{Meson Fields : } \pi^a T^a$$

➔ Baryons with spin 1/2, 3/2, ..., Nc/2

$$B = \begin{pmatrix} N \\ \Delta \\ \vdots \\ \vdots \end{pmatrix} \quad \text{Degrees of freedom : Hadrons}$$

➔ Leading order (Spin-flavor symmetry + chiral symmetry )

$$L_B = B^\dagger \left( iD_0 + g_A^\circ u^{ia} G^{ia} - \frac{C_{HF}}{N_c} \vec{S}^2 - \frac{c_1}{\Lambda} \hat{\chi}_+ \right) B$$

the axial coupling is at LO  $\dot{g}_A = \frac{6}{5}g_A$ , being  $g_A = 1.2732(23)$

$$(\partial_0 - i\Gamma_0)$$

$$\text{Tr} \langle (u^\dagger (i\partial_i + r_i) u - u (i\partial_i + l_i) u^\dagger) \lambda^a \rangle$$

$$\left( \frac{1}{2} (u^\dagger (i\partial_0 + r_0) u + u (i\partial_0 + l_0) u^\dagger) \right)$$

$$\hat{\chi}_+ \equiv \tilde{\chi}_+ + N_c \chi_+^0$$

$$\chi = 2B_0(s + ip),$$

$$\chi_\pm = u^\dagger \chi u^\dagger \pm u \chi^\dagger u,$$


$$\chi_\pm^0 = \langle \chi_\pm \rangle,$$

$$\tilde{\chi}_\pm \equiv \chi_\pm^a T^a, \quad \text{where } \chi_\pm^a \equiv \frac{1}{2} \langle \lambda^a \chi_\pm \rangle$$

## Baryon Masses to $\mathcal{O}(\xi^2)$ in SU(2)

$$m_{\mathbf{B}} = N_c M_0 + \frac{C_{HF}}{N_c} \hat{S}^2 + c_1 M_\pi^2 + \delta m_{\mathbf{B}}^{1\text{-loop}+CT}$$

A. Calle-Cordon & J.L. Goity (PHYSICAL REVIEW D 87, 016019 (2013))



## Baryon Masses to $\mathcal{O}(\xi^3)$ in SU(3)

I. P. FERNANDO and J. L. GOITY ( PHYS. REV. D 97, 054010 (2018) )

$$\begin{aligned} \mathcal{L}_B = \mathbf{B}^\dagger & \left( iD_0 + \hat{g}_A u^{ia} G^{ia} - \frac{C_{HF}}{N_c} \hat{S}^2 - \frac{1}{2\Lambda} c_2 \hat{\chi}_+ + \frac{c_3}{N_c \Lambda^3} \hat{\chi}_+^2 \right. \\ & \left. + \frac{h_1}{N_c^3} \hat{S}^4 + \frac{h_2}{N_c^2 \Lambda} \hat{\chi}_+ \hat{S}^2 + \frac{h_3}{N_c \Lambda} \chi_+^0 \hat{S}^2 + \frac{h_4}{N_c \Lambda} \chi_+^a \{S^i, G^{ia}\} + \alpha \hat{Q} + \beta \hat{Q}^2 \right) \mathbf{B} \end{aligned}$$

$$\begin{aligned} \hat{\chi}_+ &= N_c \chi_+^0 + \tilde{\chi}_+ \\ \chi_+^0 &\rightarrow 4B_0 m^0 \\ \tilde{\chi}_+^a &\rightarrow 8B_0 \delta^{a8} m^8 \\ \hat{\chi}_+ &\rightarrow 4B_0 (m^8 T^8 + N_c m^0) \end{aligned}$$

$$m^0 = (2\hat{m} + m_s)/3$$

$$m^8 = 2/\sqrt{3}(\hat{m} - m_s)$$



$$\begin{aligned}
m_B = & M_0 + \frac{C_{HF}}{N_c} \hat{S}^2 - \frac{c_1}{\Lambda} 2B_0(\sqrt{3}m_8 Y + N_c m_0) - \frac{c_2}{\Lambda} 4B_0 m_0 \\
& - \frac{c_3}{N_c \Lambda^3} \left( 4B_0(\sqrt{3}m_8 Y + N_c m_0) \right)^2 \\
& - \frac{h_1}{N_c^2 \Lambda} \hat{S}^4 - \frac{h_2}{N_c \Lambda} 4B_0(\sqrt{3}m_8 Y + N_c m_0) \hat{S}^2 - \frac{h_3}{N_c \Lambda} 4B_0 m_0 \hat{S}^2 \\
& - \frac{h_4}{N_c \Lambda} \frac{4B_0 m_8}{\sqrt{3}} \left( 3\hat{I}^2 - \hat{S}^2 - \frac{1}{12} N_c(N_c + 6) \right. \\
& \left. + \frac{1}{2}(N_c + 3)Y - \frac{3}{4}Y^2 \right) + \delta m_B^{\text{loop}},
\end{aligned}$$

$$\delta m_B^{1-loop} = i \frac{\overset{\circ}{g}_A^2}{F_\pi^2} \frac{1}{d-1} \sum_n G^{ia} \mathcal{P}_n G^{ia} I_{1-loop}(\delta m_n - p^0, M_\pi)$$

$$\Delta_{GMO} \equiv 3m_\Lambda + m_\Sigma - 2(m_N + m_\Xi)$$

The breaking to the GMO relation is only coming through the loop corrections and it behaves like  $1/N_c$  in the strict large  $N_c$  limit

$$\begin{aligned}
\Delta_{GMO} = & - \left( \frac{\overset{\circ}{g}_A}{4\pi F_\pi} \right)^2 \left( \frac{2\pi}{3} \left( M_K^3 - \frac{1}{4} M_\pi^3 - \frac{2}{\sqrt{3}} \left( M_K^2 - \frac{1}{4} M_\pi^2 \right)^{\frac{3}{2}} \right) \right. \\
& + \frac{C_{HF}}{2N_c} \left( 4M_K^2 \log \left( \frac{4M_K^2 - M_\pi^2}{3M_K^2} \right) - M_\pi^2 \log \left( \frac{4M_K^2 - \frac{1}{3} M_\pi^2}{3M_\pi^2} \right) \right) \Bigg) \\
& + \mathcal{O}(1/N_c^3).
\end{aligned}$$



## 3.2

## Fit Results to Baryon Masses







## Fit results to Experimental & Lattice QCD masses

I. P. FERNANDO and J. L. GOITY

PHYS. REV. D **97**, 054010 (2018)

TABLE II. Results for LECs: the ratio  $\bar{g}_A/F_\pi = 0.0122 \text{ MeV}^{-1}$  is fixed by using  $\Delta_{\text{GMO}}$ . The first row is the fit to LQCD octet and decuplet baryon masses [48] including results for  $M_\pi \leq 303 \text{ MeV}$  (dof = 50), and second row is the fit including also the physical masses (dof = 58). Throughout the  $\mu = \Lambda = m_\rho$ .

$\chi^2_{\text{dof}}$	$m_0$ [MeV]	$C_{\text{HF}}$ [MeV]	$c_1$	$c_2$	$h_2$	$h_3$	$h_4$
0.47	221(26)	215(46)	-1.49(1)	-0.83(5)	0.03(3)	0.61(8)	0.59(1)
0.64	191(5)	242(20)	-1.47(1)	-0.99(3)	0.01(1)	0.73(3)	0.56(1)




$$\sigma_{fB}(m_f) = m_f \frac{\partial}{\partial m_f} m_B = \frac{m_f}{2m_B} \langle \mathbf{B} | \bar{q}_f q_f | \mathbf{B} \rangle$$


1) The value of the pion-Nucleon sigma term ranges from 45 MeV to 64 MeV

Eur. Phys. J. C (2018) 78:569

John Ellis, Natsumi Nagata, Keith A. Olive



$$\sigma_{\pi N} \equiv \frac{\hat{m}}{2m_N} \langle N | \bar{u}u + \bar{d}d | N \rangle$$



2) There is a long lasting “puzzle” associated with a combination of baryon masses (in SU(3) ) in the iso-spin symmetric limit, to obtain the pion-Nucleon sigma term, assuming the contribution by strange quark mass to the nucleon mass is negligible (OZI).

3) The connection between the pion-Nucleon sigma term and size of the correction to the Gell-Mann-Okubo relation

Can one explain these from the combined approach ?

Baryon matrix elements of scalar quark densities give us the information on the amount of baryon mass originates from the quark masses

## Feynman-Hellman theorem

$$\sigma_i(B) = m_i \frac{\partial}{\partial m_i} m_B$$

Baryon mass dependencies on quark masses

$$\sigma_{\pi N} = \hat{\sigma} + 2 \frac{\hat{m}}{m_s} \sigma_s$$

$m_i$  indicates a quark mass

$$\sigma_{\pi N} \equiv \frac{\hat{m}}{2m_N} \langle N | \bar{u}u + \bar{d}d | N \rangle$$

$$\hat{\sigma} = \frac{\hat{m}}{2m_N} \langle N | \bar{u}u + \bar{d}d - 2\bar{s}s | N \rangle$$

$$\sigma_s = \frac{m_s}{2m_N} \langle N | \bar{s}s | N \rangle$$

$$\sigma_{\pi N} \sim \hat{\sigma}$$

$$|\sigma_s| \lesssim 50 \text{ MeV}$$

A long lasting puzzle !

$$\sigma_{\pi N} = \hat{\sigma} + 2 \frac{\hat{m}}{m_s} \sigma_s$$

$$\hat{\sigma} \equiv \sqrt{3} \frac{\hat{m}}{m_8} \sigma_8$$

$$\sigma_8 = \frac{1}{3} (2m_N - m_\Sigma - m_\Xi)$$

$$m_3 = m_u - m_d$$

$$m_8 = \frac{1}{\sqrt{3}} (\hat{m} - m_s)$$

$$\hat{\sigma} = \underbrace{\frac{\hat{m}}{m_s - \hat{m}}}_{\Delta \hat{\sigma}} (m_\Xi + m_\Sigma - 2m_N) \sim 26 \text{ MeV}$$

There is a (hidden) large correction  $\sim 44 \text{ MeV}$  from non-analytic contributions from baryon self-energies

$$\sigma_8 = \frac{1}{2m_N} \langle N | \bar{u}u + \bar{d}d - 2\bar{s}s | N \rangle$$

$$\Delta \sigma_8 \equiv \sigma_8 - \frac{1}{3} (2m_N - m_\Sigma - m_\Xi)$$

$$\Delta \sigma_8 = \sigma_8 - \frac{1}{9} \left( \frac{5N_c - 3}{2} m_N - (2N_c - 3) m_\Sigma - \frac{N_c + 3}{2} m_\Xi \right)$$

$$\Delta_{GMO} \equiv 3m_\Lambda + m_\Sigma - 2(m_N + m_\Xi) \sim 25 \text{ MeV}$$

The dominant contributions to  $\Delta_{GMO}$  and  $\Delta \sigma_8$  are calculable non-analytic contributions:  $\Delta \sigma_8 / \Delta_{GMO}$  ( $\sim -13.5$  for  $N_c = 3$ )

$$\sigma_{8N} = \frac{1}{9} \left( \frac{5N_c - 3}{2} m_N - (2N_c - 3) m_\Sigma - \frac{N_c + 3}{2} m_\Xi \right)$$


$$\sigma_{8\Delta} = \frac{N_c}{3} (m_\Delta - m_{\Sigma^*}) - \frac{5(N_c - 3)}{12} (m_\Lambda - m_\Sigma)$$

	$\frac{\dot{g}_A}{F_\pi}$	$\frac{M_0}{N_c}$	$C_{HF}$	$c_1$	$c_2$	$h_2$	$h_3$	$h_4$	$\alpha$	$\beta$
Fit	MeV <sup>-1</sup>	MeV	MeV						MeV	MeV
1	0.0126(2)	364(1)	166(23)	-1.48(4)	0	0	0.67(9)	0.56(2)	-1.63(24)	2.16(22)
2	0.0126(3)	213(1)	179(20)	-1.49(4)	-1.02(5)	-0.018(20)	0.69(7)	0.56(2)	-1.62(24)	2.14(22)
3	0.0126*	262(30)	147(52)	-1.55(3)	-0.67(8)	0	0.64(3)	0.63(3)	-1.63*	2.14*
	$\Delta_{GMO}^{\text{phys}}$	$\sigma_{8N}$	$\Delta\sigma_{8N}$	$\hat{\sigma}_N$	$\sigma_{\pi N}$	$\sigma_{sN}$	$\sigma_{8\Delta}$	$\Delta\sigma_{8\Delta}$	$\hat{\sigma}_\Delta$	
	MeV	MeV	MeV	MeV	MeV	MeV	MeV	MeV	MeV	
1	25.6(1.1)	-583(24)	-382(13)	70(3)(6)	—	—	-496(46)	-348(16)	59(5)(6)	
2	25.5(1.5)	-582(55)	-381(20)	70(7)(6)	69(8)(6)	-3(32)	-511(52)	-352(22)	60(10)(6)	
3	25.8*	-615(80)	-384(2)	74(1)(6)	65(15)(6)	-121(15)	-469(26)	350(27)	56(4)(6)	

$$\sigma_{\pi N} = 69(10) \text{ MeV}$$

$$\sigma_{\pi\Delta} = 60(10)(6) \text{ MeV}$$

Hadronic weak currents possess V-A Lorentz structure of the weak interactions



It is important to know these axial and vector couplings in order to extract the standard model parameters for flavor mixings

$$J_\mu = V_\mu - A_\mu$$

$$V_\mu = V_{ud}\bar{u}\gamma_\mu d + V_{us}\bar{u}\gamma_\mu s$$

$$A_\mu = V_{ud}\bar{u}\gamma_\mu\gamma_5 d + V_{us}\bar{u}\gamma_\mu\gamma_5 s$$

$$\langle B_2 | V_\mu | B_1 \rangle = V_{CKM} \bar{u}_{B_2}(p_2) \left[ f_1(q^2) \gamma_\mu + \frac{f_2(q^2)}{M_{B_1}} \sigma_{\mu\nu} q^\nu \right] u_{B_1}(p_1)$$

$$\langle B_2 | A_\mu | B_1 \rangle = V_{CKM} \bar{u}_{B_2}(p_2) \left[ g_1(q^2) \gamma_\mu + \frac{g_2(q^2)}{M_{B_1}} \sigma_{\mu\nu} q^\nu \right] \gamma_5 u_{B_1}(p_1)$$

$$\bullet \quad \mathcal{L}_B^{(2)} = \mathbf{B}^\dagger \left( \frac{c_2}{\Lambda} \chi_+^0 + \frac{C_1^A}{N_c} u^{ia} S^i T^a + \frac{\kappa}{2\Lambda} B_+^{ia} G^{ia} + \dots \right) \mathbf{B}$$

where the flavor  $SU(3)$  electric and magnetic fields are denoted by  $E_+$  and  $B_+$  and given by  $E_+^i = F_+^{0i}$  and  $B_+^i = \frac{1}{2}\epsilon^{ijk} F_+^{jk}$

$$\bullet \quad \mathcal{L}_B^{(3)} = \mathbf{B}^\dagger \left( \frac{g_1}{\Lambda^2} D_i E_+^a T^a + \frac{\kappa_1}{2\Lambda N_c} B_+^{ia} S^i T^a + \dots \right) \mathbf{B}$$

$$\bullet \quad \mathcal{L}_B^{(4)} = \mathbf{B}^\dagger \left( \frac{1}{N_c \Lambda^2} (g_2 D_i E_+^a S^j G^{ja} + g_3 D_i E_+^a \{S^i, G^{ja}\}^{\ell=2}) + \frac{\kappa_r}{\Lambda^3} D^2 B_+^{ia} G^{ia} \right. \\ \left. + \frac{1}{2\Lambda^3} (\kappa_2 \chi_+^0 B_+^{ia} G^{ia} + i\kappa_F f^{abc} \chi_+^a B_+^{ib} G^{ic} + \kappa_D d^{abc} \chi_+^a B_+^{ib} G^{ic} + \kappa_3 \chi_+^a B_+^{ia} S^i) \right. \\ \left. + \frac{1}{2\Lambda N_c^2} (\kappa_4 B_+^{ia} \{\hat{S}^2, G^{ia}\} + \kappa_5 B_+^{ia} S^i S^j G^{ja}) + \dots \right) \mathbf{B}$$

- The LECs  $g_1$  and  $g_2$  will be determined by charge radii
- The term proportional to  $g_3$  gives electric quadrupole moment for  $\mathbf{10}_B$  and  $\mathbf{10}_B \rightarrow \mathbf{8}_B$  transitions.
- The term proportional to  $\kappa_r$  gives contribution to magnetic radii
- The renormalization of the magnetic moments is provided by LECs  $\kappa_{D,F,1,\dots,5}$



# 5.1


## Baryon vector currents

$$f_1 = f_1^{SU(3)} + \delta f_1$$

### ONE LOOP CORRECTIONS

RUBÉN FLORES-MENDIETA AND JOSÉ L. GOITY PHYSICAL REVIEW D **90**, 114008 (2014)

I. P. FERNANDO and J. L. GOITY PHYS. REV. D **97**, 054010 (2018)



$$p_0 \rightarrow \delta m_{in}$$

$$p_0 + q_0 \rightarrow \delta m_{out}.$$

$$\delta \hat{m} \equiv \frac{C_{HF}}{N_c} \hat{S}^2 - \frac{c_1}{2\Lambda} \hat{\chi}_+$$

$SU(3)$  breaking in  $\delta \hat{m}$  are disregarded

$q_0 = SU(3)$  breaking mass difference + kinetic energy

$$\Gamma^{\mu a} = \underbrace{g^{\mu 0} T^a}_{\text{charge}} + i \underbrace{\frac{\kappa}{\Lambda} \epsilon^{0\mu ij} f^{abc} f^{cbd} q^i G^{jd}}_{\text{magnetic moment}}$$

charge

magnetic moment

Ademollo-Gatto Theorem (AGT) is “satisfied” : The amplitude of vector currents in the  $q^2 \rightarrow 0$  limit are uniquely predicted up to first order in symmetry breaking.

## 5.2

## Baryon vector currents charges

$$f_1 = f_1^{SU(3)} + \delta f_1$$

At lowest order the charges are represented by the flavor generators  $T^a$

$$f_1^a(A_{1+2+3})^{\text{poly}} = \frac{\lambda_\epsilon - 3}{(4\pi)^2} \left( \frac{\dot{g}_A}{4F_\pi} \right)^2 Q^2 T^a$$

$$f_1^a(B_{1+2})^{\text{poly}} = -\frac{\lambda_\epsilon + 1}{(4\pi)^2} \frac{Q^2}{4F_\pi^2} T^a$$

● UV finite ;  $Q^2 \rightarrow 0$

● At  $Q^2 \rightarrow \text{finite}$  :  
UV divergent terms renormalized via  $g_1$  and  $g_2$  in the Lagrangian

## Results

SU(3) breaking corrections to the  $\Delta S = 1$  vector charges.

- $\delta f_1/f_1 = \mathcal{O}(1/N_c)$
- Dominant contribution to the Corrections are entirely by the non-analytic pieces

If one doesn't combine chiral and  $1/N_c$ , then the  $N_c$  power counting is violated

	$\frac{\delta f_1}{f_1}$		experimentally not enough precision from semi-leptonic hyperon decays to determine
	One-loop	LQCD	
$\Lambda p$	$-0.067(15)$	$-0.05(2)$	
$\Sigma^- n$	$-0.025(10)$	$-0.02(3)$	
$\Xi^- \Lambda$	$-0.053(10)$	$-0.06(4)$	
$\Xi^- \Sigma^0$	$-0.068(17)$	$-0.05(2)$	


(P. E. Shanahan et al)

## 5.3

## Baryon charge radii

$$\langle r^2 \rangle = -6 \frac{df_1(Q^2)}{dQ^2} \Big|_{Q^2 \rightarrow 0}$$

Only the following diagrams contribute



● Charge operator for generic  $N_c$

$$\hat{Q} = T^3 + \frac{1}{\sqrt{3}} T^8 + \frac{3-N_c}{6N_c} B$$

Some important observations (at strict large  $N_c$  limit)


loop contributions \ $T^a$	$T^3$	$T^8$
Diagram A3	$\mathcal{O}(N_c^0)$	$\mathcal{O}(N_c^0)$
Diagram B2	$\mathcal{O}(1/N_c)$	$\mathcal{O}(N_c^0)$

● Dominant non-analytic contributions to the radii are proportional to  $\log m_q$

## 5.3

## Baryon charge radii (Cont...)

$$\langle r^2 \rangle = -6 \frac{df_1(Q^2)}{dQ^2} \Big|_{Q^2 \rightarrow 0}$$



Contributions from Counter Terms (CT)  
Satisfies the following relation

$$a\Lambda + p + \Sigma^+ + \frac{1}{3}(a-4)(n + \Sigma^0 + \Xi^0) + \Sigma^- + \Xi^- = 0$$

resulting from the electric charge being a U-spin singlet

	$\langle r^2 \rangle [\text{fm}^2]$		
	Full	CT	Exp
p	0.707	0.596	0.7071(7)
n	-0.116	-0.049	-0.116(2)
$\Lambda$	-0.029	-0.024	...
$\Sigma^+$	0.742	0.596	...
$\Sigma^0$	0.029	0.024	...
$\Sigma^-$	0.683	0.548	0.608(156)
$\Xi^0$	-0.016	-0.049	...
$\Xi^-$	0.633	0.548	...

proton radius used is the one resulting from the muonic Hydrogen Lamb shift

## 5.4

## Baryon magnetic moments

LO magnetic moment is given by,  $\frac{\kappa}{2\Lambda} B_+^{ia} G^{ia}$  in the  $\mathcal{O}(\xi^2)$  Lagrangian

$$e \frac{\kappa}{2\Lambda} = \mu_p = 2.7928 \mu_N$$

LO magnetic moment operator  $G^{ia}$  is proportional to the LO axial currents

LO ratios of magnetic moments



Note that the experimentally available magnetic moment ratios and corresponding LO results shows that the combined approach can describe well these ratios at LO

NLO effects stem from quark masses and spin symmetry breaking

$SU(3)$  breaking corrections  $\mathcal{O}((m_s - \hat{m})N_c)$


spin symmetry breaking corrections  $\mathcal{O}(1/N_c)$

	Exp	LO
$p/n$	-1.46	-1.5
$\Sigma^+/\Sigma^-$	-2.12	-3
$\Lambda/\Sigma^+$	-0.25	$-\frac{1}{3}$
$p/\Sigma^+$	1.14	1
$\Xi^0/\Xi^-$	1.92	2
$p/\Xi^0$	-2.23	-1.5
$\Delta^{++}/\Delta^+$	1.4(2.8)	2
$\Omega^-/\Delta^+$	-0.75	-1
$p/\Delta^+$	1.03	1
$p/(\Delta^+ p)$	0.78	$\frac{3}{2\sqrt{2}}$
$p/(\Sigma^{*0}\Lambda)$	1.02	$\sqrt{\frac{3}{2}}$
$p/(\Sigma^{*+}\Sigma^+)$	-0.88	$-\frac{3}{2\sqrt{2}}$

## 5.4

## Baryon magnetic moments

One loop corrections to the magnetic moments are obtained from the spatial components of the following diagrams



Renormalization of magnetic moments

$$\begin{aligned}
 & \frac{1}{\Lambda} \kappa_1 \frac{1}{N_c} B_+^{ia} S^i T^a + \frac{1}{2\Lambda} \left\{ \frac{1}{\Lambda^2} (\kappa_2 \chi_+^0 B_+^{ia} G^{ia} + \kappa_3 d^{abc} \chi_+^a B_+^{ib} G^{ic} + \kappa_4 \chi_+^a B_+^{ia} S^i) \right. \\
 & + \left. \kappa_5 \frac{1}{N_c^2} B_+^{ia} \{\hat{S}^2, G^{ia}\} + \kappa_6 \frac{1}{N_c^2} B_+^{ia} S^i S^j G^{ja} \right\}
 \end{aligned}$$

Renormalization of magnetic radii  $\frac{\kappa_r}{\Lambda^3} D^2 B_+^{ia} G^{ia}$

## 5.4

## Baryon magnetic moments

As an input proton and neutron magnetic moments giving the following relation between LECs

$$\kappa_1 = -19.662 + 6.926 \kappa - 0.833 \left( \kappa_4 + \frac{\kappa_5}{2} \right) + 2.550 \kappa_D$$

$$\kappa_3 = -5.136 + 1.648 \kappa - 0.218 \left( \kappa_4 + \frac{\kappa_5}{2} \right) + \kappa_D.$$

LEC $\times \frac{m_N}{\Lambda}$	LO	NNLO		$\mu_{LO}$	$\mu_{NNLO}$	$\mu_{Exp}$		$\mu_{LO}$	$\mu_{NNLO}$	$\mu_{Exp}$
$\kappa$	2.80	2.887	p	2.691	Input	2.792847356(23)	$\Delta^{++}$	5.381	5.962	3.7 – 7.5
$\kappa_1$	0	3.29	n	-1.794	Input	-1.9130427(5)	$\Delta^+$	2.691	3.049	2.7(3.6)
$\kappa_2$	0	0.00	$\Sigma^+$	2.691	2.367	2.458(10)	$\Delta^0$	0	0.136	...
$\kappa_D$	0	0.397	$\Sigma^0$	0.897	0.869	...	$\Delta^-$	-2.691	-2.777	...
$\kappa_F$	0	...	$\Sigma^-$	-0.897	-0.629	-1.160(25)	$\Sigma^{*+}$	2.691	3.151	...
$\kappa_3$	0	0.53	$\Lambda$	-0.897	-0.611	-0.613(4)	$\Sigma^{*0}$	0	0.343	...
$\kappa_4$	0	-2.85	$\Xi^0$	-1.794	-1.275	-1.250(14)	$\Sigma^{*-}$	-2.691	-2.465	...
$\kappa_5$	0	1.05	$\Xi^-$	-0.897	-0.652	-0.6507(25)	$\Xi^{*0}$	0	0.490	...
			$\Delta^+ p$	2.537	3.65	3.58(10)	$\Xi^{*-}$	-2.691	-2.208	...
			$\Sigma^0 \Lambda$	1.553	1.57	1.61(8)	$\Omega$	-2.691	-2.005	-2.02(5)
			$\Sigma^{*0} \Lambda$	2.197	2.68	2.73(25) <sup>a</sup>				
			$\Sigma^{*+} \Sigma^+$	-2.537	-2.35	-3.17(36) <sup>b</sup>				

Coleman Glashow (CG) relation  $\mu_p - \mu_n - \mu_{\Sigma^+} + \mu_{\Sigma^-} + \mu_{\Xi^0} - \mu_{\Xi^-} = 0$

valid at tree level NNLO and receives only a finite correction from the one loop contributions.





Only the magnetic radii of proton and neutron are experimentally known

	$\kappa_r = -2.63$ $\langle r^2 \rangle [\text{fm}^2]$		
	Exp	Th	Loop
p	0.724	0.718	0.134
n	0.746	0.747	0.179
$\Sigma^+$	...	0.766	0.100
$\Sigma^0$	...	0.698	0.061
$\Sigma^-$	...	0.922	0.189
$\Lambda$	...	0.895	0.079
$\Xi^0$	...	0.872	0.081
$\Xi^-$	...	0.796	0.035
$\Delta^+ p$	...	0.875	0.226

LQCD can test  
these  
predictions

At tree level  $A^{\mu c} = \dot{g}_A G^{jc} \left( g_j^\mu - \frac{q^\mu q_j}{q^2 - M_b^2} \delta^{bc} \right) \xrightarrow{\text{In the large } N_c \text{ limit}} A^{ic}$



At the leading order, axial couplings are given in terms of  $\dot{g}_A$


Octet :  $F = \dot{g}_A/3 \quad D = \dot{g}_A/2$       Decuplet :  $\mathcal{H} = \dot{g}_A/6$

At vanishing 3-momentum  $\langle \mathbf{B}' | A^{ic} | \mathbf{B} \rangle = g_A^{\mathbf{B}\mathbf{B}'} \frac{6}{5} \langle \mathbf{B}' | G^{ic} | \mathbf{B} \rangle$

The calculations up to 1-loop corrections to the axial-currents are performed in SU(3)



$$A^{ic} + \delta A_{1\text{-loop}}^{ic}$$

Diagrams contributing to the 1-loop corrections to the axial vector currents in SU(3)



$$\begin{aligned}
 & + \frac{C_1^A}{N_c} u^{ia} S^i T^a + \frac{C_2^A}{N_c} \epsilon^{ijk} u^{ia} \{S^j, G^{ka}\} + \frac{C_3^A}{N_c^2} u^{ia} \{\hat{S}^2, G^{ia}\} + \frac{C_4^A}{N_c^2} u^{ia} S^i S^j G^{ja} \\
 & + \frac{D_1^A}{\Lambda^2} \chi_+^0 u^{ia} G^{ia} + \frac{D_2^A}{\Lambda^2} \chi_+^a u^{ia} S^i + \frac{D_3^A(d)}{\Lambda^2} d^{abc} \chi_+^a u^{ib} G^{ic} + \frac{D_4^f(f)}{\Lambda^2} f^{abc} \chi_+^a u^{ib} G^{ic}
 \end{aligned}$$



SU(2)



A. CALLE CORDÓN AND J.L. GOITY  
PHYSICAL REVIEW D **87**, 016019 (2013)

## 6

## Baryon axial-vector currents : Fits to LQCD




## Fit results


C. Alexandrou et al. Phys. Rev. **D94**, 034502 (2016)

Fit	$\chi^2_{\text{dof}}$	$\dot{g}_A$	$\delta \dot{g}_A$	$C_1^A$	$C_2^A$	$C_3^A$	$C_4^A$	$D_1^A$	$D_2^A$	$D_3^A$	$D_4^A$
LO	3.9	1.35	-	-	-	-	-	-	-	-	-
NLO Tree	0.91	1.42	-	-0.18	-	-	-	-	0.009	-	-
NLO Full	1.08	1.02	0.15	-1.11	0.	1.08	0.	-0.56	-0.02	-0.08	0.
	1.13	1.04	0.08	-1.17	0.	1.15	0.	-0.59	-0.02	-0.09	0.
	1.19	1.06	0.	-1.23	0.	1.21	0.	-0.62	-0.03	-0.09	0.



First LQCD  
calculations  
for baryon  
axial currents  
including  
hyperons and  
the decuplet :


 $g_A^3$ 

[LQCD from Alexandrou et al, (2016)]


 $g_A^8$

- The  $\sigma$  terms of nucleons were calculated using  $SU(3)$  BChPT  $\times 1/N_c$
- Our value for  $\sigma_{\pi N}$  is in agreement with similar determinations in calculations that included the decuplet baryons as explicit degrees of freedom
- The “ $\sigma$  term puzzle” is understood as the result of large non-analytic contributions to the mass combination, while the higher order corrections to the  $\sigma$  terms have natural magnitude.
- The intermediate spin  $3/2$  baryons play an important role in enhancing  $\hat{\sigma}$  and thus  $\sigma_{\pi N}$
- The analysis carried out here shows that there is compatibility in the description of  $GMO$  and the nucleon  $\sigma$  terms
- The value of  $\sigma_{\pi N} = 69 \pm 10$  MeV obtained here from fitting to Physical & LQCD baryon masses agrees with the more recent results from  $\pi N$  analyses



- $BChPT \times 1/N_c$  improves convergence by eliminating large  $N_c$  power power violating terms in loop corrections
- $SU(3) BChPT \times 1/N_c$  shows a great improvement in describing charge, charge-radii, magnetic moments, magnetic-radii
- Only two LECs are needed to determine charge-radii of baryons
- Only eight LECs are needed to determine magnetic moments of baryons
- Only one LEC is needed to determine magnetic radii of baryons
- Axial couplings are also an important test of this approach
- More LQCD calculations are welcome, and current predictions can be used to test experimentally as well as in LQCD



Thank you

