

Tolga Erboral¹, Joerg Reinhold¹
¹Florida International University, Miami, Florida




Introduction

We are studying the photoproduction mechanisms of η mesons at the GlueX experiment in Hall-D at Thomas Jefferson National Laboratory in Newport News, VA. These particles are produced by a linearly polarized photon beam at energies between 8.2 and 8.8 GeV incident on a liquid hydrogen target.

Azimuthal (ϕ) angular distributions for the η with respect to the direction of the polarized photon facilitate the extraction of the beam asymmetry Σ for the reaction $\gamma p \rightarrow \eta p$. Σ , derived as a function of four-momentum transfer $-t$, quantitatively denotes contributions from natural and unnatural parity exchanges in η photoproduction. Compared with previous GlueX results [1,2], new data using 100% of the allocated GlueX-I runtime produced approximately 6 times more statistics, thereby allowing to extend these measurements to values well beyond the previous limitation of $-t \leq 1.1$ (GeV/c)². Preliminary results will be shown for events reconstructed from the decay of $\eta \rightarrow \gamma\gamma$.



Experiment

An 11.6 GeV electron beam from the CEBAF accelerator is directed onto a diamond radiator, producing coherent, polarized bremsstrahlung. This photon beam is transported to the GlueX experiment. Drift chambers track charged particles through the magnetic field, providing momentum and angle measurements. The energy and direction of photons from the η decay are measured with calorimeters. Full reconstruction of the final state allows identification of η mesons in the $\gamma\gamma$ invariant mass.




Triplet polarimeter values (statistical errors only) for the analyzed data. Photons can be polarized either parallel or perpendicular to the floor in a 0/90 degree pair or a -45/45 degree pair.

Preliminary Distributions



Peak and sideband region Σ vs Invariant Mass (All $-t$)




Method

The azimuthal yields were used to obtain the yield asymmetry $f(\phi) = \frac{Y_{\perp} - F_R Y_{\parallel}}{Y_{\perp} + F_R Y_{\parallel}}$, where F_R is the photon flux ratio obtained from a given run period.

Σ was extracted from fits to $f(\phi) = \frac{(P_{\parallel} + P_{\perp})\Sigma \cos(2\phi - 2\phi_0)}{2 + (P_{\perp} - P_{\parallel})\Sigma \cos(2\phi - 2\phi_0)}$, where P_{\parallel} and P_{\perp} are corresponding polarization values, ranging from 32-35%.

Preliminary Results



Preliminary statistical errors only for our data (GlueX Phase-I analyzed), compared to results from previous measurements. Our data values are set arbitrarily at 1.0.

Outlook & Acknowledgements

We expect to further improve the statistical errors by performing a detailed analysis of the background of the $\eta \rightarrow \gamma\gamma$ decay as well as other systematic corrections. Also, the range in t is expected to improve with forthcoming GlueX data. We also expect to achieve similar progress in the $\eta \rightarrow \pi^+\pi^-\pi^0$ decay.

This work was partially supported by the U.S. Department of Energy, Office of Science, Office of Nuclear Physics under contracts DE-SC0013620 and DE-AC05-06OR23177.

References

- [1] S. Adhikari et al. Beam asymmetry Σ for the photoproduction of η and η' mesons at $E_{\gamma} = 8.8$ GeV. Physical Review C, 100(5), Nov 2019.
- [2] H. Al Ghoul et al. Measurement of the beam asymmetry Σ for π^0 and η photoproduction on the proton at $E_{\gamma} = 9$ GeV. Physical Review C, 95(4), Apr 2017.